

2020-04-30

# Groundnut rust disease epidemiology and potential sustainable management strategies

Francis, Magreth

INNSPUB

---

<https://dspace.nm-aist.ac.tz/handle/20.500.12479/1551>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*



## Groundnut rust disease epidemiology and potential sustainable management strategies

Magreth Francis<sup>\*1,2</sup>, Musa Chacha<sup>1</sup>, Ernest R. Mbega<sup>1</sup>, Patrick A. Ndakidemi<sup>1</sup>

<sup>1</sup>*School of Life Sciences and Bio-engineering The Nelson Mandela African Institution of Science and Technology (NM- AIST), Arusha, Tanzania*

<sup>2</sup>*Centre for Research, Agricultural Advancement, Teaching Excellence and Sustainability in Food and Nutrition Security (CREATES-FNS), Tanzania*

**Key words:** *Puccinia arachidis*, Epidemiology, Management, Pesticidal plants

<http://dx.doi.org/10.12692/ijb/16.4.524-536>

Article published on April 30, 2020

### Abstract

Groundnut is an important food and cash crop for the majority of the world's population. However, diseases have impeded groundnut cultivation in groundnut-growing regions. Of them, groundnut rust disease caused by *Puccinia arachidis* is one among the major diseases causing more than 50% economic yield losses. For decades, chemical pesticide application has been the most effective option for controlling groundnut in Tanzania rust disease. However, hazards associated with excessive usage of chemical pesticides and development of pathogen- resistance against pesticides has prompted intensive research on disease management alternatives that are effective and eco-friendly to the environment. Thus, this review will focus on groundnut rust disease etiology, epidemiology and expounds various management options with much emphasis on the potential of the selected pesticidal plant species (*Azadirachta indica*, *Jatropha curcas*, *Parthenium hysterophorus* and *Moringa oleifera*) in combatting the disease.

\* **Corresponding Author:** Magreth Francis ✉ [francism@nm-aist.ac.tz](mailto:francism@nm-aist.ac.tz)

## Introduction

Groundnut (*Arachis hypogaea* L.) is an important legume crop and it is among the most important oil and protein sources for smallholders in tropical and subtropical countries of the world (Pande *et al.*, 2003). It contains 48-50% of oil and 26-28% of protein and is a rich source of dietary fibres, minerals (zinc, iron, riboflavin, thiamine, phosphorus, magnesium, calcium and potassium) and vitamins (niacin, folate, thiamin, riboflavin, choline, Vitamin B<sub>6</sub> and E) (Pandey *et al.*, 2012; Izge *et al.*, 2007).

Despite the importance of groundnut in Tanzania, its production is still low with an average of 718kg/ha (MAFC, 2010) while expected potential yield is 1500kg/ha (Philipo and Nchimbi-Msolla, 2019; Mponda and Kafiriti, 2008). Groundnut production and economic returns are threatened yearly by destructive fungal diseases namely rust, early leaf spots and late leaf spots (USDA, 2003). When these foliar fungal diseases attack groundnut in combination they may cause about 70% yield loss. Nevertheless, in many regions of the world where groundnut is grown, groundnut rust disease caused by the basidiomycete fungus *Puccinia arachidis* Speg., which belongs to class Pucciniomycetes has been identified to cause up to 57% yield loss mainly during rainy season since the condition favour disease development (Mansoor, 2012; Patel *et al.*, 1987). Other losses associated are reductions in the amount and quality of foliage used for fodder (Subrahmanyam and McDonald, 1987). To mitigate the groundnut rust disease, so far synthetic fungicides have been recommended as the most effectual management option (Monyo *et al.*, 2009). However harmful impact of most of synthetic fungicides application on human and environment also development of pathogen resistance against pesticides has led to a growing demand of using pesticidal plants as protection materials.

Based on researches and many scientific works gives evident that pesticidal plant are biodegradable and have less risks on human and environment also are cost effective unlike widely used synthetic chemicals.

Nevertheless little is still known of the available resources of pesticidal plants and their great role on crop protection. Therefore investigating efficiency of more plant species/ resources available will enhance utilization of the local resources in order to produce more bio pesticides which are highly needed. Thus, this review will expounds various management options with much emphasis on the potential of the selected pesticidal plant species (*Azadirachta indica*, *Jatropha curcas*, *Parthenium hysterophorus* and *Moringa oleifera*) in combatting groundnut rust disease.

### *Groundnut rust disease*

The disease is caused by a pathogen *Puccinia arachidis* Speg., which is a member of the Urediniomycetes where the Uredinales, the rusts, is a single order, containing approximately 5000 host-specific, obligate parasites (Agrios, 2005; Subrahmanyam *et al.*, 1983). The diseases significantly reduces the pod and fodder yield and oil quality (Agrios, 2005; Subrahmanyam *et al.*, 1983).

### *Occurrence and distribution of Groundnut rust disease*

The disease was first recorded in 1827 from Surinam, in 1910 and in Soviet Union and Mauritius in 1914. Thereafter, groundnut rust was considered a disease of South America prior to 1970. In the recent years, the disease has spread around the world where the groundnut is grown in Asia (Brunei, Philippines, Indonesia, Japan, Korea, Thailand, India, Taiwan, Malaysia and Taiwan), Australasia and Oceania (Australia, Fiji, Solomon Island and New Guinea) and Africa (Burkina Faso, Sudan, Benin, Malawi, Botswana, Ethiopia, Uganda, Kenya, Tanzania, Mauritius, Mozambique, Nigeria, Zambia, Senegal, Zimbabwe and Republic of South Africa) (Subrahmanyam *et al.*, 1985; Bromfield, 1971). Since this pathogen is found wherever the crop is grown, there is need to seek for sustainable management option because the pathogen tend to spread very fast causing huge crop losses.

### *Hosts of Puccinia arachidis*

The pathogen *P. arachidis* is genus-specific, being limited to *Arachis* including *A. glabrata*, *A. burkartii*,

*A. marginata*, *A. cardenasii*, *A. helodes*, *A. stenosperma*, *A. nambyquarae* and *A. hagenbeckii* (Hennen *et al.*, 1987). It does not have any alternate host either in legume or non-legume plants (Hennen *et al.*, 1987; Subrahmanyam and McDonald 1983). The obligate nature of the pathogen has implications in management of the disease. For instance, practicing crop rotation could be a better solution since the pathogen cannot infect any non-*Arachis* genus. However, farmers may not be able to do crop rotation by substituting with crops that are not of interest to them and therefore searching for other management options for groundnut rust disease is of paramount importance to groundnut growing farmers.

#### *Biology and ecology of Puccinia arachidis*

The *Puccinia arachidis* pathogen is known to perpetuate and spread via urediniospores (Savary *et al.*, 1985; Subrahmanyam *et al.*, 1983). The urediniospores may disperse through wind movement, rain splash, insects and sometimes by the movement of the infected crop debris, pods or seeds (Subrahmanyam and McDonald, 1983). Moreover the presence of groundnut plants or volunteer groundnut plants in one or other parts of the country enabled the survival of the pathogen from season to season on self-sown (volunteer) groundnut plants (Subrahmanyam *et al.*, 1983). Furthermore, (Sunkad and Kulkarni, 2007) reported that the urediniospores on self-sown/voluntary peanut plants play an important role in survival and viability of the pathogen from season to season. The existing survival mechanism of the groundnut rust pathogen shows challenges in breaking the disease cycle hence need to find the best management option for the disease.

#### *Factors affecting P. arachidis survival*

##### *Temperature*

Urediniospores production is highly favored at temperature ranging between 20-28°C and subsequently leads to infection and disease development (Rao *et al.*, 1994; Subrahmanyam and McDonald, 1987). The temperatures below 20°C or above 30°C are highly detrimental to urediniospore production (Rao *et al.*, 1994). Consequently it is

important to seek for new management options to control the disease since the favourable weather condition for crop production also favors disease infection.

##### *Wind*

Wind as agent of urediniospore dispersal is responsible for spreading the groundnut rust disease. Normally distribution of wind speed near leaves is a bit more complex and is influenced by canopy density, height and roughness (Aylor, 1990). The urediniospore is disseminated long distance during dry windy weather by the movement of the infected crop debris or seeds surface-contaminated with uredospores moreover the infection depends on leaf wetness from rain or dew (Savary *et al.*, 1985). In view of the above, the wind factor can be controlled by planting tall trees to act as wind breakers and hence reducing the disease dispersal emanating from wind.

##### *Moisture*

Moisture in the plant environment comprises humidity, dew, rainfall or water from irrigation. This is critical factor in the spread of most plant diseases. Normally, rain is characterized by its duration and the amount of fallen rain [mm/h], with great variation in each rain event (Ivan, 2000). The distribution of rain drop diameters depends on the number of drops per unit volume and the rain intensity and different types of rain events (Ulbrich, 1983). Also according to Savary and Janeau, 1986; the light rains favor disease dispersal than heavy showers because the spore content of the lesions in the canopy are strongly reduced by heavy rain as compared with light rain. Moreover, urediniospore germination depends on the presence of fluid water (Mallaiah and Savary 1986; Rao, 1982). Thus taking measure against that factor is the most economical means of preventing the disease development.

##### *Humidity*

Normally, growth rate of spores of some fungi occurs at 100% relative humidity and sometimes could not grow below 93% relative humidity (Subrahmanyam and McDonald 1983; Rao *et al.*, 1994).

High relative humidity favour urediniospore production and subsequently favour rust disease development (Savary *et al.*, 1985). Since it is difficult to manage relative humidity in the ecosystem and thus there is need to seek for different management options in controlling this important disease of groundnut.

#### *Symptomatology of Groundnut rust disease*

The disease occurs in the field under warm conditions at temperature ranging between 15-28°C and when leaves are wet due to rain or dew the uredospores germinate and produce appressoria that penetrate the leaf through the stomata (Mallaiah and Rao, 1979). Always incubation period ranges between 9 to 20 days, depending on environmental factors and host reaction and spreads quickly by repeated infection cycles of urediniospores. Normally it starts with whitish flecks on the abaxial surface of leaves (Chiteka *et al.*, 1988; Subrahmanyam *et al.*, 1985). Thereafter, pustules rupture after about 2 days of appearance to expose circular or oval urediniospores which are dark-orange at first but become cinnamon-brown with maturity. Later on, epidermis ruptures and exposes a powdery mass of urediniospores. Under wet and warm weather, the spots may be formed on all aerial plant parts from flowers to pegs in contrast with the rapid defoliation associated with leaf spots (Subrahmanyam *et al.*, 1985). When the disease progresses, it may lead to production of small and shriveled seeds which subsequently have implication on yield (Subrahmanyam and McDonald, 1983). The disease symptoms describe the detrimental effects on groundnut growth and yield attributes hence this knowledge helps to decide the proper time for starting disease management in order to combat great losses.

#### *Common management options used to control groundnut rust disease*

##### *Cultural control and sanitary methods*

This employ practices that make the environment less attractive to pathogen and less favorable for their survival, dispersal, growth and reproduction while promoting the development of a desired plant.

It involves reducing inoculum, intercropping, crop rotation, plant quarantine regulations, mulching and soil amendments (Thurston, 1992; Sill, 1982). Some of effective practices reported against *P. arachidis* involves long break between successive groundnut crops aiming to disrupt viable urediniospores, removing volunteer groundnut plants and groundkeepers. Subrahmanyam and McDonald, 1983; suggested adjusting times of sowing as well as controlling weeds was effective as assisted in avoiding the environmental conditions conducive for disease build-up. Also intercropping pearl millet or sorghum with groundnut (1:3) has been reported to be useful in reducing the intensity of rust disease (Ghosh and Dayal, 1998). Other practices include strict plant quarantine regulations to avoid the movement and spread of rust on pods or seeds to disease free areas (Palti, 1981). Moreover application of high doses of phosphorous fertilizers (60-75kg phosphate/ha) has been reported to slow down the development of rust (Mayee, 1983). Although cultural control is effective in managing rust disease but those practices should be integrated with other management option to make it more effective.

##### *Biological control*

Biological control of fungal diseases of plants is eco-friendly and potential component of integrated disease management option which involves use of natural enemies including parasites, predators and competing organisms to low levels at which there is little economic damage (Pal and McSpadden, 2006). This method is highly specific to a particular pest and it provides long term solution to a pest problem and it reputed of being environmentally eco-friendly. According to Biswas *et al.*, 2010, 70% rust disease reduction was observed when treated with *Trichoderma viride* or *Trichoderma harzianum* as foliar spray. Similarly, Govindasamy and Balasubramanian, 1989; reported the significant reduction of number of uredosori/leaflet and uredospore/pustule by *Trichoderma harzianum* on detached groundnut leaves. Furthermore, according to Ghewande, 2006; maximum inhibition of *in vitro* germination of urediniospores and significant reduction in *in vivo* development of rust was

exhibited by *V. lecanii* and *P. islandicum*. Furthermore, a study by Meena *et al.*, 2010; proved efficacy of a talc-based powder formulation of *Pseudomonas fluorescens* strain Pfl in controlling groundnut late leaf spot and rust when applied as seed as well as foliar treatment.

The biological control option seems effective against groundnut rust disease nevertheless it involve research hence too expensive. It is only successful on confined places i.e. 'glasshouse pests' and pest problems located in small area/ islands moreover time and patience is needed. Basing on economic status of small holder farmers in Sub Saharan Africa, there is great need of searching of cheaper, effective and eco-friendly management option for management of groundnut rust disease.

#### *Use of resistance varieties*

Use of resistant variety is the simplest, practical, effective and economical option for rust management (Pande and Rao, 2001). It also saves time, energy and money spent on other control measures. According to (Subrahmanyam 1995; Subrahmanyam *et al.*, 1985) in recent years there have been concerted efforts by many countries to exploit genetic resistance to groundnut rust disease due to its importance. More than 13,000 accessions of groundnut germplasm were successfully screened at ICRISAT, India leading to the identification of over 160 genotypes with resistance to rust (Singh and Singh, 1997; Mehan *et al.*, 1996; Subrahmanyam and McDonald 1980). These resistant varieties are characterized by slow rusting type, increased incubation period, reduced infection frequency, reduced lesion diameter and low sporulation index (Mehan *et al.*, 1996; Subrahmanyam *et al.*, 1983).

Despite the fact that resistant varieties are effective in managing the rust disease, there are some challenges because in order to develop acceptable varieties it may take 10-15 years (Brown 1980) which is such a long time coming up with a single variety. Furthermore, the host-pathogen system resistance in groundnut rust is not long-lasting (Johnson, 1979). Similarly, sometimes the resistant varieties may have lower crop yield and poor quality hence not preferred

by farmers. Therefore, the limitations associated with use of resistant varieties open a door to explore other management options.

#### *Chemical control*

Numerous synthetic fungicides have been reported to be effective against groundnut rust disease. For instant several dust formulations of copper and the combinations of mancozeb and zinc and several sterol-inhibiting fungicides proved to be effective against groundnut rust (Subrahmanyam, 1997). Likewise, Chiteka *et al.* 1988; reported that when Tridemorph 2gm/lit of water sprayed 3-4 times at 15-20 days intervals gave good management of groundnut rust. Also according to (Culbreath *et al.*, 2002), spraying chlorothalonil 0.2%; or mancozeb 0.25% or Hexaconazole/propaconazole 30 days after germination till 15 days before harvesting at regular 10-15 day intervals reduced disease incidences. Likewise, Calixin at 0.5 ml/liter of water or Kavach at 2 g/liter of water controls groundnut rust (Culbreath *et al.*, 2002). The above mentioned synthetic fungicides respond quickly and effectively in managing the rust disease. However, due to poor socio economic conditions of small holder farmers in the developing world, they cannot afford or follow the recommendations for the application of the optimum amounts. In most cases they apply lower dosage which has implications in developing pathogen resistance to the chemicals (Monyo *et al.*, 2009). Subsequently, this calls for the affordable substitute to the chemical pesticides. The use of botanical pesticides previously called 'native pesticides' has gained reputation due to their less risks to environment, human, livestock and natural enemies.

#### *Pesticidal plant*

Currently, a lot of interest has been developed in the antimicrobial effects of pesticidal plants for plant disease management aiming to replace synthetic pesticides (Zida *et al.*, 2008). According to (Zida *et al.*, 2008; Davicino *et al.*, 2007), there are more than 250,000 higher plant species on earth that can be evaluated for their antimicrobial bioactive chemical compounds/phytochemicals. These phytochemicals occur naturally in plants and are classified into two

groups' i.e. primary metabolites and secondary metabolites. The primary metabolites (ethanol, lactic acid and amino acids) are mainly for growth, development and reproduction of the organism (Sofowora, 1993). While the secondary metabolites i.e. alkaloids, steroids, saponins, tannins and phenol compounds are synthesized and deposited in specific parts or in all parts of the plant serves as physical defense systems (Ndakidemi and Dakora, 2003; Joseph and Raj, 2010).

A number of pesticidal have been screened to be effective against foliar fungal disease of groundnut such as early leaf spots, late leaf spots and rust (Harsha *et al.*, 2011). According to Kishore and Pande, 2005; 38 plant extracts were screened against late leaf spot and rust disease of groundnut and found that aqueous leaf extracts (20%,w/v) of *Prosopis juliflora* and *Lycopersicon esculentum* completely inhibited the spore germination of *Phaeoisariopsis personata* and *Puccinia arachidis*. Similarly, chloroform extracts of *Hemionitis arifolia* was effective against *P. personata* and *P. arachidis* (Sahayaraj *et al.*, 2009). Another study by Eriyanto and Alfi (2016), reported the efficacy of crude extract

5% of *Ageratum conyzoides* against rust disease of groundnut. Kishore and Pande (2005) confirmed a positive effect of extracts of *Datura metel* (25g/L) and *Lawsonia inermis* (50g/L) against late leaf spot lesion and rust pustules by 65-74% compared with control. Also an aqueous extracts of *Eupatorium odoratum* L. showed antifungal activity against spore germination of *Puccinia arachidis* over control. Moreover, chloroform extracts of three fern plants were effective against *Phaeoisariopsis personata* and *Puccinia arachidis* (Sahayaraj *et al.*, 2009). These evidences proves the antifungal activity of plant extracts in managing rust-causing pathogen *Puccinia arachidis* and open a new prospect in plant disease management by attracting antimicrobial investigation of available plant species against groundnut rust disease. We are focusing on the potential phytochemical compounds possessed by the four selected pesticidal plant species (*Azadirachta indica*, *Moringa oleifera*, *Parthenium hysterophorus* and *Jatropha curcas*) since they have proved efficient against different fungal pathogens as described in Table 1. This call for further studies on how these selected plant species can effectively manage groundnut rust disease and improve crop production.

**Table 1.** Identified selected pesticidal plant species against fungal pathogens.

Plant	Plant part	Diseases/pathogen	References
<i>Azadirachta indica</i>	Leaf, stem bark, root, shoot,	Anthraco nose pepper <i>Fusarium solani</i> <i>Alternaria porri</i> Stemphylium blight diseases Protozoa, bacteria <i>Aspergillus flavus</i> , <i>Alternaria solani</i> <i>Colletotrichum capsici</i> , <i>Schizophyllum</i> <i>commune</i>	(Nduagu <i>et al.</i> , 2008) (Dharam and Sharma, 1985) (Atawodi and Atawodi, 2009) (Mohammad and Forough, 2007) (Shrivastava and Kshma, 2014; Muktarul <i>et al.</i> , 2009)
<i>Parthenium</i> <i>hysterophorus</i>	Shoot, leaf, root, stem	<i>Fusarium solani</i> <i>Alternaria alternata</i> <i>Aspergillus niger</i> , <i>Candidia albicans</i> <i>Fusarium oxysporum</i> <i>Aspegillus niger</i> <i>Aspegillus flavus</i> <i>Alternaria brassicae</i> , <i>Alternaria brassicola</i>	(Gupta <i>et al.</i> , 1996) (Pareek <i>et al.</i> , 2011) (Singh and Srivastava, 2013; Zunera <i>et al.</i> , 2012;) (Harsha <i>et al.</i> , 2011), (Ajay <i>et al.</i> , 2013; Barsagade and Wagh 2010) (Harsha <i>et al.</i> 2011; Rukhsana <i>et al.</i> , 2003)
<i>Moringa oleifera</i>	Leaves, seeds, stem, roots	<i>Rhizopus</i> <i>stolonifer</i> and <i>Microsporium gypseum</i> <i>Gibberella xylarioides</i> <i>G. candidum</i> <i>M. micheli</i> and <i>R. stolonifer</i>	(Ganie <i>et al.</i> , 2015) (Nkya <i>et al.</i> , 2014) (Jabeen <i>et al.</i> , 2008); (Oluduro, 2012); (Bamishaiye <i>et al.</i> , 2011)
<i>Jatropha curcas</i>	Leaves seeds, stem, roots	<i>Colletotrichum musae</i> <i>Colletotrichum gloeosporioides</i> <i>Fusarium oxysporum</i> , <i>Pythium aphanidermatum</i> , <i>Lasiodiplodia theobromae</i> , <i>Fusarium</i> <i>semitectum</i> , <i>Colletotrichum capsici</i> and <i>Colletotrichum gloeosporioides</i>	(Oskoueian <i>et al.</i> , 2011) (Saetae and Suntornsuk, 2010)

Phytochemical compounds found in four selected plant species (*Azadirachta indica*, *Jatropha curcas*,

*Parthenium hysterophorus* and *Moringa oleifera*) for pesticidal activity.

*Azadirachta indica*

*Azadirachta indica* also known as neem, it is native to India, Pakistan and Bangladesh, currently grown in tropical and semi-tropical regions. It is a large evergreen dense tree growing some 10 to 15 meter height and width of about 2-3 meter. Products made from neem trees have been used in India for over two millennia for their medicinal properties. Neem products possess good antifungal, antibacterial, antiviral, anthelmintic, antidiabetic, sedative and contraceptive since they possess most important phytochemicals steroids, phenolic compounds, saponins, alkaloids, glycosides, terpenoids, flavonoids and tannins (Ramadass and Subramanian, 2018; Shrivastava and Kshma 2014; Mohammad and Forough, 2007). Moreover Neem have other valuable active phytochemicals such as azadirachtin, nimbolinin, nimbin, nimbidin, nimbidol, salannin and quercetin, 6-desacetylnimbinene, nimbandiol, nimbolide, ascorbic acid, n-hexacosanol and amino acid, 7-desacetyl-7-benzoylazadiradione, 7-desacetyl-7-benzoylgedunin, 17-hydroxyazadiradione and nimbiol (Hossain *et al.*, 2011; Kokate *et al.*, 2010; Muktarul *et al.*, 2009).

These important ingredients plays important role in the inhibition of growth of numerous microbes such as viruses, bacteria and pathogenic fungi through cell wall breakdown (Hossain *et al.*, 2013); Mordue and Nisbet, 2000). Therefore, it is vital to explore the potential of neem extracts in management of groundnut rust disease.

*Moringa oleifera*

*Moringa oleifera* commonly known as Moringa, drumstick tree, ben oil tree, is the only genus in the family Moringaceae. It is native to northwestern India and widely cultivated in tropical and subtropical areas. Moringa tree is a small, graceful, deciduous tree with sparse foliage, often resembling a leguminous species. The tree grows to 8m height with smooth, dark grey; slash thin, yellowish (Fuglie, 2001).

Approximately all parts of the Moringa tree are edible and consumable by humans. It is used worldwide traditionally as medicine for various health conditions

since it has antifungal, antibacterial and anti-inflammatory properties (Bamishaiye *et al.*, 2011; Anyasor, *et al.*, 2011; Mahmood *et al.*, 2010; Fuglie, 2001; Sairam, 1999). The *Moringa oleifera* possess numerous bioactive components including vitamins, phenolic acids, flavonoids, saponins, alkaloids, isothiocyanates, tannins, glycosides, terpenoids and steroids (Oluduro, 2012; Jabeen *et al.*, 2008). *Moringa oleifera* plant has been proved to be effective against numerous fungal pathogens this encourages scientists for further studies on its efficacy against other diseases such rust disease of groundnut.

*Parthenium hysterophorus*

*Parthenium hysterophorus* L. (Asteraceae) is an annual herbaceous plant that can reach a height of 1.5m in its native range, but sometimes can grow up to 2.5m where it is an invasive species (McConnachie *et al.*, 2011). It is commonly known as carrot weed, white top, ragweed parthenium, congress grass and star weed. It is widely distributed in semi-arid, subtropical, tropical and warmer temperate regions with a preference for warmer temperate climates (Adkins and Shabbir, 2014; Shabbir and Bajwa, 2006; EPPO, 2018).

Researchers reported *P. hysterophorus* has antiviral, antifungal, antibacterial, anti-helmintic, anti-molluscal and anti-inflammatory properties (Sesha *et al.*, 2013; Sukanya *et al.* 2009). The major components of toxic being parthenin and other phenolic acids such as caffeic acid, vanillic acid, anisic acid, p-anisic acid, chlorogenic acid and parahydroxy benzoic acid are toxic to human beings and animals (Mahadevappa, 1997; Gupta *et al.*, 1996). The phytochemical investigation of *P. hysterophorus* revealed the presence of various chemical constituents viz., alkaloids, proteins, saponins, tannins, carbohydrate, glycosides, terpenoids, steroids, volatile oils, amino acids, amino sugars, lignans, phenolic compounds, flavonoids, metallic elements, organic acids and terpenoids (Lakshmanan *et al.*, 2013; Sankar and Dipak 2014; Pareek *et al.*, 2011; Das and Das, 1995; Bakhtiar *et al.*, 2012.) Possession of various phytochemicals enable its efficacy against different plant diseases hence the study on how *P. hysterophorus* can be effective in managing groundnut rust disease is important.



*Jatropha curcas*

*Jatropha curcas* L is a drought resistant perennial plant, belonging to genus *Jatropha* and is a member of the Euphorbiaceae family, producing oil containing seeds (Amit *et al.*, 2010). It originates from Mexico or neighboring parts of Central America, which is the only area where it has often been collected from undisturbed vegetation (Igbiosa and Igbiosa, 2009; Henning, 2007). *Jatropha* plants can easily be grown on marginal soils to help repopulate land, hence can be used to conserve the land (Igbiosa and Igbiosa, 2009).

All parts of *Jatropha* (seeds, leaves, bark, and roots) play a major role in the treatment of various disease pathogens, including bacteria and fungi (Prasad *et al.*, 2012; Duke, 1988). The phytochemical analysis of the methanol extract of the leaf, stem bark and root of *Jatropha curcas* revealed the presence of alkaloids, flavonoids, saponins, glycosides, tannins, terpenoids, resins and steroids in varying quantities (Muklesur *et al.* 2011; Oskoueian *et al.* 2011; Obasi *et al.*, 2011;) Saetae and Suntornsuk, 2010. The fungicidal properties of *J. curcas* against different fungal pathogens call for further research in the utilization of this plant in managing groundnut rust disease.

**Conclusion**

Based on researches and evidences from other research groups, the active ingredients found on the selected pesticidal plants (*Azadirachta indica*, *Moringa oleifera*, *Jatropha curcas* and *Parthenium hysterophorus*) are biodegradable in nature and possess less risks to the environment, human, livestock and natural enemies, Thus these plants may be used as possible substitute for synthetic fungicide in the management of groundnut rust disease. Moreover these plants are cheap and readily available to small holder farmers in Tanzania Africa, who do not have ready access to chemical fungicides.

**Acknowledgements**

We would like to acknowledge DAAD and Centre for Research, Agricultural Advancement, Teaching Excellence and Sustainability in Food and Nutrition Security (CREATES- FNS) for their financial support.

**References**

- Adkins S, Shabbir A.** 2014. Biology, ecology and management of the invasive *Parthenium* weed (*Parthenium hysterophorus* L.). Pest Management Science **70(7)**, 1023-1029.
- Agrios GN.** 2005. Plant Pathology. 5<sup>th</sup> Ed., Elsevier Academic Press, Amsterdam, New York, USA.
- Ajay K, Shailesh J, Tripti M.** 2013. Antimicrobial Potential of *Parthenium hysterophorus* Linn. Plant Extracts. Intern. Journal of Life science Biotechnology and Pharmacy Research **2(3)**, 232-236.
- Amit S, Sonal S, Rami U, Rajore S, Batra M.** 2010. Broad-spectrum antimicrobial properties of medicinally important plant, *Jatropha curcas* L. International Journal Pharmacology Science Review and Research **4(3)**, 11-13.
- Anyasor GN, Aina DA, Olushola MM, Aniyikaye AF.** 2011. Phytochemical constituent, proximate analysis, antioxidant, antibacterial and wound healing properties of leaf extracts of *Chromolaena odorata*. Annual Biology Research **2**, 441-451.
- Atawodi SE, Atawodi JC.** 2009. *Azadirachta indica* (Neem) a plant of microbial and pharmacological activity. Phytochemical Review **8**, 601-620.
- Aylor DE.** 1990. The role of intermittent wind in the dispersal of fungal pathogens, Annual Review. Phytopathological **28**, 73-92.
- Bakhtiar M, Rashid K, Yasir A.** 2012. Phytochemical and cytotoxic analysis of *Parthenium hysterophorus* selected from District Bannu, Pakistan. African Journal of Biotechnology **11(55)**, 11857-11860.
- Bamishaiye EI, Olayemi FF, Awagu EF, Bamshaiye OM.** 2011. Proximate and phytochemical composition of *Moringa oleifera* leaves at three stages of maturation. Advance Journal of Food Science Technology **3**, 233-237.
- Barsagade NB, Wagh GN.** 2010. Comparative screening of leaf extracts of common plants and weeds for their antibacterial and antifungal activities. Asiatic Journal of Biotechnology. Resources **3**, 227-232.

- Biswas S, Dasgupta N, Bratati D.** 2010. Antioxidant Activity of some Indian medicinal plants used for the treatment of diabetes mellitus. *Pharmacologyonline* **3**, 7-21
- Bromfield KR.** 1971. Peanut rust: a review of literature. *Journal of American Peanut Research Education Association* **3**, 111-112.
- Brown JF.** 1980. The genetics of resistance in plants to infection by pathogens. *Plant protection*, pp. 267-275.
- Chiteka ZA, Kucharek TA, Knauft DA, Gorbett DW, Shokes FM.** 1988. Components of resistance to late leaf spots in peanut. I. Levels and variability-implications for selection. *Peanut Science* **15**, 25-30.
- Culbreath AK, Stevenson KL, Brenneman TB.** 2002. Management of late leaf spot of peanut with benomyl and chlorothalonil: A study in preserving fungicide utility. *Plant Disease* **56**, 349-355.
- Das B, Das R.** 1995. Chemical investigation in *Parthenium hysterophorus* An-allelopathic plant. *Allelopathy Journal* **2(1)**, 99-104.
- Davicino R, Mattar MA, Casali YA, Graciela S, Margarita E, Micalizzi B.** 2007. Antifungal activity of plant extracts used in folk medicine in Argentina. *Revista Peruana de Biología* **14**, 247-251.
- Dharam V, Sharma RK.** 1985. Efficacy of fungicides XXIX studies on the fungicidal properties of neem oil. *Indian Journal of Plant Pathology* **3**, 241-242.
- Duke JA.** 1988. CRC. Handbook of Medicinal Herbs. CRC Press, Boca Raton, Florida, p. 253-254.
- EPPO.** 2018. PQR database. Paris: European and Mediterranean Plant Protection Organization.
- Eriyanto Y, Alfi I.** 2016. Methanolic extracts of three weeds as botanical fungicides to control peanut rust disease. *Nusant Ara Bioscience* **8(1)**, 117-122.
- Fuglie LJ.** 2001. The Miracle Tree: *Moringa Oleifera*: natural nutrition for the tropics, (Church World Service, Dakar, 1999). pp: 68. Revised in 2001 and published as The Miracle Tree: The Multiple Attributes of Moringa, p. 172
- Ganie SA, Mudasser Z, Surender SG, Surender SY, Ranjana S, Sujata G.** 2015. Antifungal Efficacy of *Moringa oleifera* Lam. *American Journal of Phytomedicine and Clinical Therapeutics* **3(1)**, 28-33.
- Ghewande MP.** 2009. Rust of groundnut an overview. In: Upadhyay RK, Mukherji KG, Chamola BP, Dubey OP (Eds) Integrated pest and disease management. APH Publishing Corp., New Delhi, pp 326-334, 110012, India.
- Ghosh PK, Dayal D.** 1998. Effect of varying levels of nitrogen in three groundnut-based intercropping systems. In: Proceeding of the International Conference on Food Security and Crop Science, CCS Haryana Agricultural University, Hissar, India, p. 133.
- Govindasamy V, Balasubramanian R.** 1989. Biological control of groundnut rust, *Puccinia arachidis*, by *Trichoderma harzianum*. *Journal of Plant Diseases and Protection* **96**, 337-341.
- Gupta N, Martin BM, Metcalfe DD, Rao PV.** 1996. Identification of a novel hydroxyproline-rich glycoprotein as the major allergen in *Parthenium* pollen. *The Journal of Allergy and Clinical Immunology* **98**, 903-912.
- Harsha M, Santosh G, Sharda S.** 2011. Antimicrobial and spermicidal activities of *Parthenium hysterophorus* Linn. and *Alstonia scholaris* Linn. *Indian Journal of Natural Products and Research* **2(4)**, 458-463.
- Hennen JF, Subhramanyam P, Figueiredo MB.** 1987. The taxonomy, life history & evolution of *Puccinia arachidis* Speg., Proceedings of a discussion group meeting on Groundnut Rust Disease held at ICRISAT Centre, Pattencheru, A.P. India, 24-28 Sep. 1984.
- Henning RK.** 2007. *Jatropha curcas* L. in H. A. M. van der Vossen and G. S. Mkamilo, (Eds.). Prota 11(1): Medicinal plants/Plantes médicinales. Plant Resources of Tropical Africa (PROTA), Wageningen, Netherlands.

- Hossain MA, Al-Toubi WAS, Weli AM, Al-Riyami QA, Al-Sabahi JN.** 2013. Identification and characterization of chemical compounds in different crude extracts from leaves of Omani neem. *Journal of Taibah University for Science* **7(4)**, 181-188.
- Hossain MA, Shah MD, Sakari M.** 2011. Gas chromatography-mass spectrometry analysis of various organic extracts of *Merremia borneensis* from Sabah. *Asian Pacific Journal of Tropical Medicine* **4(8)**, 637-641.
- Igbinosa OO, Igbinosa EOA.** 2009. Antimicrobial activity and phytochemical screening of stem bark extract of *Jatropha curcas* Linin. *Journal of Pharmacy and Pharmacology*, **3(2)**, 055-062
- Ivan S.** 2000. Short-distance dispersal of wheat rust spores by wind and rain. *Agronomie* **20**, 757-767
- Izge AU, Mohammed ZH, Goni A.** 2007. Levels of variability in groundnut (*Arachis hypogaea* L.) to Cercospora leaf spot disease-Implication for selection. *African Journal of Agriculture Research* **2**, 182-186.
- Jabeen R, Shahid MA, Jamil A, Ashra M.** 2008. Microscopic evaluation of the antimicrobial activity of seed extracts of *Moringa oleifera*. *Pakistan Journal Botany* **40**, 1349-1358.
- Johnson R.** 1979. The concept of durable resistance. *Phytopathology* **69**, 198-199.
- Joseph B, Raj SJ.** 2010. Pharmacognostic and phytochemical properties of *Aloe vera* Linn: an overview. *International Journal of Pharmaceutical Science* **4(2)**, 106-110.
- Kishore GK, Pande S.** 2005. Integrated management of late leaf spot and rust diseases of groundnut (*Arachis hypogaea* L.) with *Prosopis juliflora* leaf extracts and Chlorothalonil. *International Journal of Pest Management*. **51(4)**, 325-332.
- Kokate C, Purohit AP, Gokhale SB.** 2010. Pharmacognosy. Maharashtra, India: Nirali Prakashan; *Alternative Medicine* **13(120)**, 1-9.
- MAFC.** Agricultural Statistics 2010. Agricultural Basic Data.
- Mahmood KT, Tahira M, Ikram UH.** 2010. *Moringa oleifera*: a natural gift. *Journal of Pharmaceutical Science and Research* **2(11)**, 775-781.
- Mallaiah KV, Rao AS.** 1979. Groundnut rust: factors influencing disease development, sporulation and germination of uredospores. *Indian Journal of Phytopathology* **32**, 382-388.
- Mansoor H.** 2012. Grain Legumes Strategy for Tanzania. Presented at N2Africa workshop Tanzania, 6 and 7 November 2012.
- Mayee CD.** 1983. Epidemiology and forecasting of groundnut rust in Marathwada region: final report. Parbhani, Maharashtra, India: Marathwada Agricultural University, pp.76
- McConnachie AJ, Strathie LW, Mersie W, Gebrehiwot L, Zewdie K.** 2011. Current and potential geographical distribution of the invasive plant *Parthenium hysterophorus* (Asteraceae) in eastern and southern Africa. *Weed Research* **51(1)**, 71-84.
- Meena B, Radhajejalakshmi R, Marimuthu T, Vidhyasekaran P, Velazhahan R.** 2010. Biological control of Groundnut Late Leaf Spot and rust by seed treatment and foliar applications of a Powder Formulation of *Pseudomonas fluorescens*. *Biocontrol Science and Technology* **12**, 2.
- Mehan VK, Reddy PM, Subramanyam P, McDonald D, Singh AK.** 1996. Identification of new sources of resistance to rust and late leaf spot in peanut. *International Journal of Pest Management* **42**, 267-271.
- Mohammad MS, Forough M.** 2007. Investigation of Compounds from *Azadirachta indica* (Neem). *Asian Journal of Plant Science* **6**, 444-445.
- Monyo ES, Osiru MO, Kadyampakeni D, Mponda O, Chinyamunyamu B.** 2009. Improving Food Security and Nutrition in Malawi and Tanzania through Research on Edible Legumes. Proceedings of Stakeholder Workshops on Groundnut Production in Malawi and Tanzania held 1-2 March and 13 April 2007, Lilongwe (Malawi) and Mtwara (Tanzania).

- Mordue AJ, Nisbet AJ.** 2000. Azadirachtin from the neem tree *Azadirachta indica*: Its action against insects, *Anais da Sociedade Entomol´ogica do Brasil* **29(4)**, 615-632.
- Mponda OK, Kafiriti EM.** 2008. Jinsi ya kuzalisha Mbegu Bora za Karanga (Production of Improved Groundnut Variety Seeds). Naliendele Agricultural Research Institute, Mtwara, pp.17.
- Muklesur R, Siti HA, Mahmud TMM, Mohamad ZAR.** 2011. Extraction of *Jatropha curcas* fruits for antifungal activity against anthracnose (*Colletotrichum gloeosporioides*) of papaya. *African Journal of Biotechnology* **10(48)**, 9796-9799.
- Ndavidemi PA, Dakora FD.** 2003. Legume seed flavonoids and nitrogenous metabolites as signals and protectants in early seedling development. *Functional Plant Biology* **30**, 729-745.
- Nduagu C, Ekefan EJ, Nwankiti AO.** 2008. Effect of some crude plant extracts on growth of *Colletotrichum capsici* (Synd) Butler & Bis by, causal agent of pepper anthracnose. *Journal of Applied Bioscience* **6**, 184-190.
- Nkya JW, Paul E, Deusdedit K, Musa C.** 2014. In vitro evaluation of antifungal activity of *Moringa oleifera* Lam extracts against coffee wilt pathogen, *Gibberella xylarioides* Heim and Saccas. *American Journal of Research Communication* **2(12)**, 53-62.
- Obasi LN, Madus P, Ejikeme C, Egbuonu AC.** 2011. Antimicrobial and phytochemical activity of methanolic extract and its fractions of *Jatropha curcas* Linn stem bark. *African Journal of Pure and Applied Chemistry* **5(5)**, 91-96.
- Oluduro AO.** 2012. Evaluation of antimicrobial properties and nutritional potentials of *Moringa oleifera* Lam. leaf in South-Western Nigeria. *Malaysian Journal of Microbiology* **8**, 59-67.
- Oskoueian E, Abdullah N, Ahmad SWZ, Saad WZ, AR, Omar AR, Ho YW.** 2011. Bioactive compounds and biological activities of *Jatropha curcas* L. kernel meal extract. *International Journal of Molecular Science* **12(9)**, 5955-70.
- Pal KK, McSpadden GB.** 2006. Biological Control of Plant Pathogens. The Plant Health Instructor DOI: 10.1094/PHI-A-2006-1117-02.
- Palti J.** 1981. Cultural practices and Jectiaus crop diseases. Springer, Verlag,
- Pande S, Bandyopadhyay R, Blümmel M, Narayana RJ, Thomas D, Navi SS.** 2003. Disease management factors influencing yield and quality of sorghum and groundnut crop residues. *Field Crops Research* **84(1-2)**, 89-103.
- Pande S, Rao NJ.** 2001. Resistance of wild *Arachis* species to late leaf spot and rust in greenhouse trials. *Plant Disease* **85**, 851-855.
- Pandey MK, Monyo E, Ozias-Akins P, Liang X, Guimarães P, Nigam SN.** 2012. Advances in *Arachis* genomics for peanut improvement Patancheru, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics **96**, 978-92.
- Papavizas GC, Lewis JA.** 1988. The use of fungi in integrated control of plant diseases. In: Burge MN, editor. Fungi in biological control systems. Manchester, UK: Manchester University Press, pp. 235-253.
- Papavizas GC.** 1985. *Trichoderma* and *Glilotadium*: Biology, ecology and potential for biocontrol. *Annual Review of Phytopathology* **23**, 23-54.
- Pareek A, Suthar M, Rathore GS, Bansal V.** 2011. Feverfew (*Tanacetum parthenium* L.): A systematic review. *Pharmacognosy Reviews* **5**, 103-110.
- Patel VA, Vaishnav MV.** 1987. Assessment of losses in groundnut due to rust and tikka leaf spots in Gujrat., Gujarat Agricultural University, Sardar Krushinagar. *Research Journal* **12**, 52-53.
- Philipo M, Nchimbi-Msolla S.** 2019. Effect of Watering Regimes on Yield and Agronomic Traits of Exotic Groundnut Genotypes in Tanzania. *Journal of Advances in Biology and Biotechnology* **21(2)**, 1-6.

- Prasad DMR, Amirah I, Md. Maksudur RK.** 2012. *Jatropha curcas*: Plant of medical benefits. Journal of Medicinal Plants Research **6(14)**, 2691-2699.
- Ramadass N, Subramanian N.** 2018. Study of phytochemical screening of neem (*Azadirachta indica*). International Journal of Zoology Studies **3(1)**, 209-212.
- Rao PVS, Subrahmanyam P, McDonald D.** 1994. Effect of Temperature on Urediniospore Production and Germinability in *Puccinia arachidis*. Peanut Science **21**, 79-81.
- Rukhsana B, Afia K, Tabinda SC.** 2003. Antifungal Activity of Allelopathic Plant Extracts III. Growth Response of Some Pathogenic Fungi to Aqueous Extract of *Parthenium hysterophorus*. Pakistan Journal of plant Pathology **2(3)**, 145-156.
- Saetae D, Suntornsuk W.** 2010. Antifungal activities of ethanolic extract from *Jatropha curcas* seed cake. Journal of Microbiology and Biotechnology **20(2)**, 319-324.
- Sahayaraj K, Borgio JAF, Raju G.** 2009. Antifungal activity of three fern extracts on causative agents of groundnut early leaf spot and rust diseases. Journal of Plant Protection Research **49(2)**, 141-144
- Sairam TV.** 1999. Home remedies, Volume II: A Handbook of Herbal Cures for Commons Ailments. New Delhi, India: Penguin.
- Sankar NS, Dipak P.** 2014. Antioxidant potentials of *Parthenium hysterophorus* L. leaf extracts. Scientific Research Journal of India **3(2)**, 80-86.
- Savary S, Janeau JL,** 1986. Rain-induced dispersal in *Puccinia arachidis*, studied by means of a rainfall simulator. Netherland Journal of Plant Pathology **92**, 163-174.
- Savary S.** 1985. Effets du niveau de contamination et de la temperature sur quelques e'tapes du developpement de *Puccinia arachidis* Sveg. Aeronomie **5**, 479-485.
- Sesha SDM, Satyavathi K, Bhojaraju P.** 2013. Cytotoxic and Anthelmintic Activities of Methanolic Extract of *Parthenium Hysterophorus* L. Pharmacophore **4(6)**, 275-279.
- Shabbir A, Bajwa R.** 2006. Distribution of parthenium weed (*Parthenium hysterophorus* L.), an alien invasive weed species threatening the biodiversity of Islamabad. Weed Biology and Management **6(2)**, 89-95.
- Shrivastava DK, Kshma S.** 2014. Antifungal Activity of leaf extract of Neem (*Azadirachta Indica* Linn). International Journal of Current Microbiology Applied Science **3(5)**, 305-308.
- Sill WH.** 1982. Plant protection-an integrated interdisciplinary approach. State University Press, Ames, Iowa.
- Singh M, Singh RP.** 1997. Potato virus Y detection: Sensitivity of RT-PCR depends on the size of fragment amplified. Canadian Journal of Plant Pathology-Revue Canadienne De Phytopathologie **19**, 149-155.
- Singh P, Srivastava D.** 2013. Phytochemical screening and *in vitro* antifungal investigation of *Parthenium hysterophorus* extracts against *Alternaria alternata*. International Journal of Pharmacology **4(7)**, 190-193.
- Sofowora A.** 1993. Medicinal Plants and Traditional Medicine in Africa. Spectrum Books Ltd., Ibadan, Nigeria, pp. 191-289.
- Subrahmanyam P, Hammons RO, Nigam SN, McDonald D, Gibbons RW, Fan MY, and Yeh WL.** 1983a. International cooperative screening for resistance of peanut to rust and late leaf spot. Plant Diseases **67**, 1108-1111.
- Subrahmanyam P, McDonald D, Subba Rao PV.** 1983. Influence of host genotype on uredospore production and germinability in *Puccinia arachidis*. Phytopathology **73**, 726-729.

- Subrahmanyam P, McDonald D.** 1980. Groundnut rust-Its survival and carry-over in India. *Proceeding India Academic Science* **91**, 93-100.
- Subrahmanyam P, McDonald D.** 1983. Rust Disease of Groundnut. in: Information bulletin no. 13. International Crop Research Institute for the Semi-Arid Tropics (ICRISAT). Patancheru, Anhra Pradesh, India.
- Subrahmanyam P, McDonald D.** 1987. Groundnut rust disease: Epidemiology and control. In: *Groundnut Rust Disease: Proc Discuss Group Meet. 24-28 September, 1984, ICRISAT, Patancheru, AP 502324, India.* p 27-39.
- Subrahmanyam P, Reddy LJ, Gibbons RW, McDonald D.** 1985. Peanut rust: A major threat to peanut production in semiarid tropics. *Plant Disease* **69**, 813-819.
- Subrahmanyam P.** 1995. Screening methods and sources of resistance to rust and late leaf spot of groundnut. Report No. 9290663197. Pages 1-14. International Crops Research Institute for the Semi-Arid Tropics, Andhra Pradesh, India.
- Subrahmanyam P.** 1997. Rust. Pages 31-33 in: *Compendium of Peanut Diseases.* N. Kokalis-Burelle, D. M. Porter, R. Rodriguez-Kabana, D. H. Smith and P. Subrahmanyam, Eds. American Phytopathological Society, St. Paul, MN.
- Sukanya SL, Sudisha J, Hariprasad P.** 2009. Antimicrobial activity of leaf extracts of Indian medicinal plants against clinical and phytopathogenic bacteria. *African Journal of Biotechnology* **8(23)**, 6677-6682.
- Sunkad G, Kulkarni S.** 2007. Studies on perpetuation and carryover of groundnut rust (*Puccinia arachidis* Speg.) in Northern Karnataka. *Karnataka Journal of Agriculture Science* **20**, 297-300.
- Thurston HD.** 1992. Sustainable practices for plant disease management in farming systems. Westview Press, Boulder, Colorado.
- Ulbrich CW.** 1983. Natural variations in the analytical form of the raindrop size distribution. *Journal Climate Applied Meteorology* **22**, 1764-1765.
- USDA.** 2003. Report on the status of *Arachis germplasm* in the United States. United States Department of Agriculture Agricultural Research Service.
- Zida PE, Sérémé LP, Sankara V, Somda PI, Néya A.** 2008. Importance of seed-borne fungi of sorghum and pearl millet in Burkina Faso and their control using plant extracts. *Pakistan Journal of Biological Science* **11**, 321-331.
- Zunera Z, Shazia S, Sobiya S, Tahir M.** 2012. Antifungal potential of *Parthenium hysterophorus* L. plant extracts against *Fusarium solani*. *Scientific Research and Essays* **7(22)**, AA 2049-2054.