

2017-12

Evaluation of maize accessions for diversity, yield and tolerance to Maize Lethal Necrosis Disease in Tanzania

Kabululu, Mujuni

NM-AIST

<https://dspace.nm-aist.ac.tz/handle/20.500.12479/2490>

Provided with love from The Nelson Mandela African Institution of Science and Technology

**EVALUATION OF MAIZE ACCESSIONS FOR DIVERSITY, YIELD AND
TOLERANCE TO MAIZE LETHAL NECROSIS DISEASE IN TANZANIA**

Mujuni Sospeter Kabululu

**A Dissertation Submitted in Partial Fulfillment of the Requirements for the Degree of
PhD in Life Science of the Nelson Mandela African Institution of
Science and Technology**

Arusha, Tanzania

December, 2017

ABSTRACT

Maize (*Zea mays* L.) is among the most important crops in Tanzania with still low average yield of 1.2 metric tonnes per hectare as compared with potential yields of 4 to 5 metric tonnes per hectare. Low yield is due to a number of factors including pests and diseases. Recently, East Africa has been hard hit by a new deadly disease of maize called Maize Lethal Necrosis Disease (MLND). The disease started in Kenya in 2011 and later spread to other countries including Tanzania and it continues to spread fast in other countries. It is caused by a combination of two viruses i.e., *Maize chlorotic mottle virus* (MCMV) and any other Potyvirus, with *Sugarcane mosaic virus* (SCMV) reported for East Africa. The aim of this study was to evaluate genetic diversities of maize accessions (focusing more on landraces) with respect to their MLND resistance and optimum yield in Tanzania. Fifty one maize landraces from National Plant Genetic Resources Centre (NPGRC) in Tanzania, thirty four commercial varieties as checks for yield and thirteen elite lines from CIMMYT Kenya as checks for resistance against MLND were used. Three field experiments were conducted at Tengeru, Mlangarini and Selian in Tanzania for genetic diversity study and MLND evaluation, one field experiment was conducted at Naivasha Kenya for MLND evaluation. Molecular study was conducted at Nelson Mandela African Institution of Science and Technology in Tanzania. Significant ($p < 0.05$) variations were observed among the tested accessions and an OPV Situka 1 and a hybrid DH 04 were the best yielding varieties across Tengeru, Mlangarini and Selian with 116.01g and 115.90g per plant respectively. Landraces TZA 2793 and TZA 5170 were among the highest yielding accessions with 100.46g and 99.80g per plant respectively. The allele distribution and frequency associated with quantitative trait loci for SCMV resistance were detected with landraces and TZA 2793 as well as TZA 3544 expressed low MLND progression across Mlangarini and Naivasha. The results of significant genetic diversity and response against MLND of maize landraces tested in this study calls for further investigation to ascertain their utilization in breeding and crop improvement.


DECLARATION

I, **Mujuni Sospeter Kabululu** do hereby declare to the Senate of Nelson Mandela African Institution of Science and Technology that this dissertation is my own original work and that it has neither been submitted nor being concurrently submitted for degree award in any other institution.

Mujuni Sospeter Kabululu

Date

The above declaration is confirmed by



Dr. Tileye Feyissa

18/12/2017
Date

Prof. Patrick Alois Ndakidemi

Date

COPYRIGHT

This dissertation is copyright material protected under the Berne Convention, the Copyright Act of 1999 and other international and national enactments, in that behalf, on intellectual property. It must not be reproduced by any means, in full or in part, except for short extracts in fair dealing; for researcher private study, critical scholarly review or discourse with an acknowledgement, without a written permission of the Deputy Vice Chancellor for Academic, Research and Innovation, on behalf of both the author and the Nelson Mandela African Institution of Science and Technology.

CERTIFICATION

The undersigned certify that they have read the dissertation titled **Evaluation of Maize accessions for diversity, yield and tolerance to Maize Lethal Necrosis Disease in Tanzania** and recommend for examination in fulfillment of the requirements for the degree of PhD in Life Science and Bioengineering of the Nelson Mandela African Institution of Science and Technology.



Dr. Tileye Feyissa



Date

Prof. Patrick Alois Ndakidemi

Date

ACKNOWLEDGEMENT

First and foremost, I thank God for his grace and strength throughout my PhD study. I then acknowledge the Government of Tanzania through Nelson Mandela African Institution of Science and Technology (NM-AIST) for financial support. I recognize the support granted by the School of Life Sciences and Bioengineering and the NM-AIST management for allowing me to carry out laboratory activities as part of my research at NM-AIST laboratories. The permission to pursue this PhD study by my employer, Tropical Pesticides Research Institute (TPRI) is highly appreciated.

My sincere gratitude are extended to my esteemed supervisors, Dr. Tileye Feyissa and Prof. Patrick Ndakidemi for their scientific support and unlimited assistance right from the beginning of developing a proposal up to this point. I do thank Dr. Joseph Ndunguru for his professional contribution during proposal development.

I am also grateful to Dr. Beatus Lymo and Dr. Pavithravani B.V. for their assistance on laboratory work at NM-AIST during the DNA extraction and PCR operations. Mr. Juma Mpinga, Daniel Loishiye, Ali Mpakata and Wilson Mjenga are as well recognized for their assistance on field work at Tengeru, Selian, Mlangarini in Tanzania and Naivasha in Kenya, respectively.

I recognize the International Maize and Wheat Improvement Centre (CIMMYT) and National Plant Genetic Resources Center of Tanzania (NPGRC) for providing germplasm seeds utilized in this study; also CIMMYT together with Kenya Agriculture and Livestock Research Organization (KALRO) granted a space at the MLN screening facility in Naivasha to conduct a field experiment through artificial inoculation. I do appreciate the scientific support granted to me by Dr. Wilson Marandu, Dr. Bright Jumbo, Dr. George Mahuku and Dr. Dan Makumbi. Last but not least, I would not have reached this milestone without my parent's investment in my educational endeavor since my childhood. The steadfast love and support of my parents, wife, children and friends around me were the potential assets towards any of my achievement.

DEDICATION

I dedicate this dissertation to the Almighty God and also to my wife Grace D. John and my two little children Godlove Kabululu and Hannah Kabululu

TABLE OF CONTENTS

| | |
|---|-------------|
| ABSTRACT | ii |
| DECLARATION | iii |
| COPYRIGHT | iv |
| CERTIFICATION | v |
| ACKNOWLEDGEMENT | vi |
| DEDICATION | vii |
| TABLE OF CONTENTS | viii |
| LIST OF TABLES | xii |
| LIST OF FIGURES | xiv |
| LIST OF APPENDICES | xv |
| LIST OF ABBREVIATIONS AND SYMBOLS | xvi |
| CHAPTER ONE | 1 |
| General introduction | 1 |
| 1.0 Introduction..... | 1 |
| 1.1 Background information | 1 |
| 1.2 Research problem and justification..... | 2 |
| 1.3 Objectives | 3 |
| 1.3.1 General objective | 3 |
| 1.3.2 Specific objectives | 3 |
| 1.4 Hypothesis..... | 3 |
| CHAPTER TWO | 4 |
| Literature review | 4 |
| 2.1 Introduction..... | 5 |
| 2.2 Maize Lethal Necrosis Disease (MLND)..... | 5 |
| 2.3 Genetic diversity of maize accessions | 7 |
| 2.3.1 Morphological diversity..... | 7 |
| 2.3.2 Diversity at molecular level | 10 |
| 2.4 Importance of landraces | 12 |

| | |
|--|-----------|
| 2.5 Conclusion | 13 |
| CHAPTER THREE..... | 15 |
| Evaluation of agronomic performance of local and improved maize varieties in Tanzania..... | 15 |
| 3.1 Introduction..... | 16 |
| 3.2 Materials and methods | 16 |
| 3.3 Results and discussion | 18 |
| 3.3.1 Analysis of variance of agronomic traits | 18 |
| 3.3.2 Correlation between parameters..... | 21 |
| 3.3.3 GGE biplot analysis | 21 |
| 3.4 Conclusion | 25 |
| CHAPTER FOUR..... | 26 |
| Genetic diversity of maize landraces from Tanzania as compared with commercial improved varieties and elite lines through morphological characterization..... | 26 |
| 4.1 Introduction..... | 27 |
| 4.2 Materials and methods | 28 |
| 4.2.1 Seed materials | 28 |
| 4.2.2 Field location and experimental design..... | 30 |
| 4.2.3 Data analysis | 30 |
| 4.3 Results..... | 32 |
| 4.3.1 Variability in quantitative characters | 32 |
| 4.3.2 Qualitative characters..... | 34 |
| 4.3.3 Principal components analysis (PCA)..... | 37 |
| 4.3.4 Cluster analysis | 38 |
| 4.4 Discussion..... | 40 |
| 4.4.1 Quantitative morphological traits..... | 40 |
| 4.4.2 Qualitative morphological traits..... | 41 |
| 4.4.3 Cluster analysis | 41 |

| | |
|---|-----------|
| 4.5 Conclusion | 43 |
| CHAPTER FIVE | 44 |
| Evaluating the response of maize accessions against maize lethal necrosis disease (MLND) in east Africa..... | 44 |
| 5.1 Introduction..... | 45 |
| 5.2 Materials and methods | 47 |
| 5.2.1 Natural MLND field infestation experiment (Mlangarini, Tanzania)..... | 47 |
| 5.2.2 Artificial inoculation experiment (Naivasha, Kenya) | 49 |
| 5.3 Results..... | 51 |
| 5.3.1 Area under disease progress curve..... | 51 |
| 5.3.2 Means, variances and heritability..... | 52 |
| 5.3.3 Cluster analysis | 53 |
| 5.3.4 Correlation matrix | 53 |
| 5.4 Discussion..... | 56 |
| 5.5 Conclusion | 59 |
| CHAPTER SIX | 60 |
| Analyses of genetic diversity of maize landraces in Tanzania using SSR markers for maize lethal necrosis disease resistance..... | 60 |
| 6.1 Introduction..... | 61 |
| 6.2 Materials and methods | 63 |
| 6.2.1 Plant materials..... | 63 |
| 6.2.2 Extraction of DNA..... | 64 |
| 6.2.3 Simple Sequence Repeats markers | 64 |
| 6.2.4 Polymerase Chain Reaction and gel electrophoresis | 64 |
| 6.2.5 Data analysis | 67 |
| 6.3 Results..... | 67 |
| 6.3.1 Simple Sequence Repeats polymorphism..... | 67 |

| | |
|--|------------|
| 6.3.2 Analysis of molecular variance (AMOVA) | 68 |
| 6.3.3 Genetic relationship between groups of accessions | 69 |
| 6.3.4 Genetic variation and diversity | 70 |
| 6.3.5 Cluster and principal coordinate analyses | 70 |
| 3.3.6 Genetic diversity associated with MLND resistance | 71 |
| 6.4 Discussion | 74 |
| 6.5 Conclusion | 77 |
| CHAPTER SEVEN..... | 79 |
| General discussion, conclusion and recommendations..... | 79 |
| 7.1 General discussion | 79 |
| 7.2 Conclusion | 85 |
| 7.3 Recommendations | 86 |
| REFERENCES..... | 87 |
| APPENDICES..... | 106 |

LIST OF TABLES

| | |
|---|----|
| Table 1: Some descriptors used to evaluate genetic diversity through morphological characterization (IBPGR, 1991)..... | 9 |
| Table 2: Maize genotypes used in the evaluation of agronomic performances in Tanzania during April to September 2015..... | 17 |
| Table 3: Performance of maize local cultivars (TZA lines), OPV, improved varieties and elite lines from CYMMIT on grain yield and yield components..... | 20 |
| Table 4: The Pearson correlation coefficient of eight parameters obtained from yield evaluation experiment of 68 maize accessions at three locations..... | 21 |
| Table 5: List of maize accessions and their source as used in the genetic diversity evaluation study in Tanzania..... | 29 |
| Table 6: Some descriptors used to evaluate genetic diversity through morphological characterization (IBPGR, 1991)..... | 32 |
| Table 7: Mean values with standard errors, minimum and maximum (range), and coefficient of variation of the 19 quantitative morphological descriptors generated from the performance of the three groups of accessions..... | 33 |
| Table 8: The 12 qualitative morphological descriptors with their percentage (%) frequency accessions distribution as generated from the performance of three different groups within 68 maize accessions used in this study..... | 35 |
| Table 9: Proportions of variability contributions given by the 25 morphological traits in different principle components..... | 38 |
| Table 10: List of maize accessions and their sources as used in the screening experiments for resistance against maize lethal necrosis disease (MLND) in Kenya and Tanzania.... | 48 |
| Table 11: The selected 20 accessions at each location of experiment with the lowest relative area under disease progress curve (rAUDPC)..... | 52 |
| Table 12: Response of maize accessions to MLND under artificial inoculation at Naivasha in Kenya and natural infestation at Mlangarini in Tanzania both on a 1–5 scale..... | 53 |
| Table 13: The Pearson correlation coefficient of 29 parameters in relation to the maize lethal necrosis diseases (MLND) scores obtained from a disease screening experiment of 68 maize accessions, probability significance at $p < 0.05$ | 55 |

| | |
|---|----|
| Table 14: List of maize accessions and their sources as used in the evaluation of genetic diversity for maize lethal necrosis disease resistance. | 65 |
| Table 15: The 30 primers used in this study with their names, repeat units, bin location and sequence details. | 66 |
| Table 16: The selected 20 primers with their number of alleles, bin location, repeat units and polymorphism information content (PIC)..... | 68 |
| Table 17: Analysis of molecular variance (AMOVA) for the 96 accessions used in this study. | 69 |
| Table 18: Pairwise genetic differentiation among the three populations, Nei's genetic identity (above diagonal) and genetic distance (below diagonal)..... | 70 |
| Table 19: Genetic diversity indices for the three groups of accessions. | 70 |
| Table 20: Allelic distribution and frequency of the three groups of accessions along with the two loci associated with SCMV resistance QTL's. | 71 |

LIST OF FIGURES

- Figure 1:** The “which win where” feature of the GGE biplot for maize grain yield of 68 genotypes in three locations, where g stands for genotypes name. 22
- Figure 2:** The average environment coordination (AEC) graphic that rank genotypes relative to an ideal genotype (the center of the concentric circles) through a GGE biplot for maize grain yield of 68 genotypes at three locations. 23
- Figure 3:** The GGE biplot displaying the relation among experimental locations, their discriminating ability and representativeness through evaluation of 68 genotypes. . 24
- Figure 4:** Map of Tanzania showing the collection sites of the 50 maize landraces 31
- Figure 5:** Box plots displaying variability in morphological descriptors given by the three different groups of CIMMYT inbred lines, commercial/Improved varieties and landraces from Tanzania. Small empty boxes inside big boxes represent median, big vertical boxes show a range of values falling between 25% and 75% and the vertical lines cover the range of non-outliers. 34
- Figure 6:** Principal component analysis distributing the 68 accessions into the first two components as performed through 25 morphological traits. 36
- Figure 7:** Dendrogram for cluster analyses based on Euclidean genetic distance with average linkage of the 25 morphological characters generated from the performance of 68 maize accessions. Words in brackets show the source of seeds/collection site. 39
- Figure 8:** Dendrogram for cluster analyses based on Euclidean genetic distance with group average of the 57 accessions generated from their performance against MLND in two experiments at Naivasha, Kenya and Mlangarini, Tanzania. 54
- Figure 9:** Pie chart showing the distribution of genetic variation generated by 96 accessions divided into three groups viz landraces, commercial varieties and CIMMYT inbred lines. 69
- Figure 10:** Relationships among the 96 accessions displayed by principal coordinate analysis using SSR data. 72
- Figure 11:** Dendrogram for cluster analyses based on Ward's algorithm method for genetic distance among the 96 accessions analyzed using SSR. 73

LIST OF APPENDICES

| | |
|--|-----|
| Appendix 1: Mean values on grain yield and yield components of all maize accessions used on yield evaluation study..... | 106 |
|--|-----|

LIST OF ABBREVIATIONS AND SYMBOLS

| | |
|---------|---|
| ' | Minutes |
| " | Seconds |
| ° | Degrees |
| % | Percent |
| µl | Microlitre |
| °C | Degrees centigrade |
| AEC | Average Environment Coordination |
| AFLP | Amplified Fragment Length Polymorphisms |
| AMOVA | Analysis of Molecular Variance |
| ANOVA | Analysis of Variance |
| ASI | Anthesis Silking Interval |
| AUDPC | Area Under Disease Progress Curve |
| BLUP | Best Linear Unbiased Prediction |
| CGIAR | Consultative Group on International Agricultural Research |
| CIMMYT | International Maize and Wheat Improvement Center |
| cm | Centimeter |
| CML | CIMMYT Maize Lines |
| COSTAT | Statistical Software |
| CTAB | Cetyl trimethylammonium bromide |
| CV | Coefficient of Variation |
| DNA | Deoxyribonucleic Acid |
| dNTPs | deoxyribonucleoside triphosphate |
| E | East |
| EDTA | Ethylenediaminetetraacetic acid |
| ELISA | Enzyme-Linked Immunosorbent Assay |
| g | Grams |
| Genstat | General Statistics Software |
| GGE | Genotype plus Genotype by Environment interaction |
| h | Gene diversity |

| | |
|-----------|--|
| ha | Hectare |
| HCl | Hydrochloric Acid |
| I | Shannon – Weaver diversity index |
| IBPGR | International Board for Plant Genetic Resources |
| KALRO | Kenya Agricultural and Livestock Research Organization |
| kg | Kilograms |
| L. | Linnaeus (famous botanist) |
| LSD | Least Significant Difference |
| M | Molar mass |
| m.a.s.l | Meters above sea level |
| MCMV | Maize Chlorotic Mottle Virus |
| MDMV | Maize Dwarf Mosaic Virus |
| min | Minutes |
| ml | Millilitre |
| MLN | Maize Lethal Necrosis |
| MLND | Maize Lethal Necrosis Disease |
| mM | Millimolar |
| MT | Metric tones |
| N | Nitrogen |
| <i>nm</i> | Nanometre |
| NM-AIST | Nelson Mandela African Institution of Science and Technology |
| NPGRC | National Plant Genetic Resource Centre |
| NPK | Nitrogen, Phosphorus, Potassium |
| NTSYS pc | Numerical Taxonomy and Multivariate Analysis System |
| OPV | Open Pollinated Variety |
| p | Probability |
| PAST | Statistical Software |
| PC | Principal Component |
| PCA | Principal Component Analysis |
| PCR | Polymerase chain reaction |

| | |
|------------|---|
| pH | Potential of Hydrogen to specify the acidity or alkalinity of solutions |
| PIC | Polymorphism Information Content |
| QTL | Quantitative Trait Loci |
| RAPD | Random Amplified Polymorphic DNA |
| rAUDPC | Relative Area Under Disease Progress Curve |
| RCBD | Randomized Complete Block Design |
| RFLP | Restriction Fragment Length Polymorphism |
| S | South |
| SCMV | Sugarcane Mosaic Virus |
| SSR | Simple Sequence Repeat |
| STATISTICA | Statistical Software |
| TBE | Tris/Borate/EDTA buffer |
| TE | Tris-EDTA buffer |
| TZA | The code identifying accessions collected in Tanzania |
| U | Unit, measure for Taq polymerase |
| UPGMA | Unweighted Pair Group Method with Arithmetic Mean |
| UV | Ultra Violet |
| WSMV | Wheat Streak Mosaic Virus |

CHAPTER ONE

General introduction

1.0 Introduction

1.1 Background information

Maize (*Zea mays* L.) is one of the most important crops in the world, because it is one of the main sources of human food, animal feed, and raw materials for industrial processes (Romay *et al.*, 2013). In Tanzania, maize is the major food and cash crop as it provides 60% of dietary calories and more than 50% of utilizable protein (Mwakalinga and Massawe, 2007). It is cultivated in all the agro-ecological zones of Tanzania with an average of 2 million hectares, which is about 45% of the cultivated area in the country (Katinila *et al.*, 1998, Otunge *et al.*, 2010). The economic importance of maize in Tanzania cannot be over emphasized since its short supply is normally equated to the national food insecurity (Katinila *et al.*, 1998). However, despite the importance of maize, the general average yields are still very low with 1.2 MT per hectare as compared with the estimated potential yields of 4 to 5 MT per hectare (Moshi *et al.*, 1990; Otunge *et al.*, 2010). This has been attributed to many factors such as lack of quality agricultural inputs (seeds, fertilizer), drought, pests and diseases to mention a few. Plant diseases are a potential threat to global food security which has always been calling upon extensive research using a wide range of methodologies. Plant diseases can cause losses of 30% to 50% or even more for major crops (Ali and Yan, 2012). Maize for instance, is affected by more than 100 pathogens, where some of them cause diseases with severe impacts in different locations depending on various factors (Ali and Yan, 2012). Recently, East Africa has been hit hard by the deadly disease known as Maize Lethal Necrosis (MLN) which was not there before while potyviruses that form one of the causing pathogens used to exist. The disease is caused by a combination of *Maize Chlorotic Mottle Virus* (MCMV) and other Potyviruses such as *Sugar Cane Mosaic Virus* (SCMV), *Maize Dwarf Mosaic Virus* and or *Wheat Streak Mosaic Virus*. Since the disease is now in few areas within the region and the spread is very fast, there is a need to screen a wide range of accessions for MLN disease resistance. Maize landraces are usually genetically heterogeneous populations which are adapted to the local environments and linked to the local farming systems as well as resistance to biotic and abiotic stresses (Tokatlidis and Vlachostergios, 2016). Characterization of genetically heterogeneous populations using

conventional and molecular tools has been the most potential and efficient ways of establishing diversities for different important traits including disease resistance (Prasanna, 2012). This study involved evaluation of Tanzanian maize landraces' diversities and compared with commercial varieties (checks for yield and yield related parameters) as well as elite lines from CIMMYT (as checks for MLND resistance) and hence screen all those materials for yield and MLN disease resistance.

1.2 Research problem and justification

Maize lethal necrosis disease (MLND) presents an immediate threat to food security in Tanzania with concomitant long-term consequences. The disease infection rate can reach 100% with yields being severely affected up to a complete loss of a maize crop (Adams *et al.*, 2013; Wangai *et al.*, 2012b). The occurrence of Maize Lethal Necrosis Disease (MLND) in Eastern Africa has recently posed a serious challenge to maize production with potential negative impact on trade and food security (Mahuku *et al.*, 2015). The control of MLND is complicated because it is caused by a combination of more than one virus (*Maize chlorotic mottle virus* - MCMV and *Sugarcane mosaic virus* - SCMV) which are difficult to separately identify them based on visual symptoms (Ali and Yan, 2012). The outcome following the synergistic effect through the combinational viral infection is a serious damage that eventually kills infected plants (Wangai *et al.*, 2012a). In addition, maize plants appear to be susceptible to MLND at all growth stages from seedling to maturity (CGIAR, 2012). Stakeholders in all circles and capacities agree that the foremost priority is to identify MLND resistant maize varieties (CGIAR, 2012) as well as those with other relevant traits such as yield. Valuable natural resources of maize in terms of genetic diversity are considered to play a key role in breeding programs (Reif *et al.*, 2006). It presents a wider range of traits that can be exploited to generate new novel varieties against maize production challenges such as MLND and low yield. A good understanding of genetic diversity within and among maize landraces ensures effective utilization in plant breeding program (Rao and Hodgkin 2002). However, only 2% of the world maize germplasm collection is being absorbed in breeding programs (Dowswell *et al.*, 1996) and an important part of maize germplasm is still in the hands of small farmers (Carvalho *et al.*, 2004). Landraces and local cultivars in maize present a population which is genetically diverse and have been under farmer's selection for many years in terms of adaptation, plant characteristics, yield, biotic and abiotic

stress tolerance or resistance (Wasala *et al.*, 2013). They have so far not efficiently being utilized because of unreliable information on agronomic as well as genetic potentials (Nass *et al.*, 1993). This study aimed at evaluating the genetic diversity levels among maize landraces with other maize accessions for the purpose of screening these maize materials against MLND and low yield for potential use in breeding programs. The information obtained on maize accessions' diversity, resistance levels against MLND and yield are expected to contribute towards speedy deployment of MLND resistant materials and higher yielding varieties to generate improved maize productivity and food security.

1.3 Objectives

1.3.1 General objective

Evaluating the diversity of maize accessions for MLND resistance and optimum yield in Tanzania.

1.3.2 Specific objectives

- 1) To evaluate agronomic performance of local and improved maize varieties in Tanzania
- 2) To assess genetic diversity of maize landraces from Tanzania as compared with commercial varieties and elite lines through morphological characterization
- 3) To evaluate the response of maize accessions against maize lethal necrosis disease (MLND)
- 4) To assess genetic diversity of maize landraces in Tanzania using SSR markers for maize lethal necrosis disease resistance

1.4 Hypothesis

The following hypotheses were tested for drawing up inferences;

Ho. The selected Maize accessions have no agronomic potentials worth for breeding programs

Ho. The selected Maize accessions possess low morphological diversity for crop improvement

Ho. The selected Maize accessions possess no varied range of responses against MLND

Ho. The selected Maize accessions possess low molecular genetic diversity for crop improvement

CHAPTER TWO

Literature review

Genetic diversity of maize accessions for maize lethal necrosis disease resistance¹

Mujuni Sospeter Kabululu¹ Joseph Ndunguru^{1,2} Patrick Ndakidemi¹ * and Tileye Feyissa^{1,3}

¹Department of Sustainable Agriculture, Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Arusha, Tanzania.

²Mikocheni Agricultural Research Institute, P.O. Box 6226, Dar-Es-Salaam, Tanzania.

³Institute of Biotechnology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia.

***Corresponding author:** patrick.ndakidemi@nm-aist.ac.tz

Abstract

Maize is among the most preferred crop in Tanzania and other parts of the world. However, its production has been facing a number of challenges including Maize Lethal Necrosis Disease (MLND). The control of MLND in Eastern Africa is complicated as it is caused by a combination of more than one virus that is *Maize Chlorotic Mottle Virus* (MCMV) and *Sugarcane Mosaic Virus* (SCMV). Stakeholders agree that the priority is to identify MLND resistant maize varieties. Therefore, genetic diversity provides the source of traits required to breed against maize production challenges such as MLND. The study of genetic diversity in maize accessions often involves characterizing morphological plant characteristics as well as molecular marker techniques to study variation at DNA level. This review explores different literatures that address the importance of genetic diversity and the possibility of generating information towards obtaining potential materials against maize production challenges in general and MLND in particular.

Key words: Genetic diversity, Landraces, Maize, MLND, Molecular markers, Morphological characterization.

¹ Article published in Indian Journal of Agricultural Research (*Vol. 51, Issue 1, pp 17 - 24*)

2.1 Introduction

Maize (*Zea mays* L.) is one of the most important crops in the world, because it is one of the main sources of human food, animal feed, and raw materials for industrial processes (Romay *et al.*, 2013). In Tanzania, maize is the major food and cash crop where its short supply is normally equated to the national food insecurity (Katinila *et al.*, 1998). However, despite its importance, the general average yields are still very low with 1.2 MT per hectare as compared with the estimated potential yields of 4 to 5 MT per hectare (Moshi *et al.*, 1990; Otunge *et al.*, 2010). Cultivation of maize is hampered by several diseases which cause serious grain loss (Anjichi, 2005; Pechanova and Pechan, 2015). Recently, East Africa has been hard hit by a deadly disease known as Maize Lethal Necrosis (MLN) which was not there before while potyviruses that form one of the causing pathogens used to exist (Wangai *et al.*, 2012a; Wangai *et al.*, 2012b; Adams *et al.*, 2014; Gowda *et al.*, 2015; Mahuku *et al.*, 2015). The disease is caused by a combination of *Maize Chlorotic Mottle Virus* (MCMV) and other Potyviruses such as *Sugar cane Mosaic Virus* (SCMV), *Maize Dwarf Mosaic Virus* or *Wheat Streak Mosaic Virus* (Scheets, 1998; De Groote *et al.*, 2016; Isabirye and Rwomushana, 2016). There is a need to screen a wide range of accessions for MLN disease tolerant or resistant materials. Maize landraces (accessions) are usually genetically heterogeneous populations (Ignjatovic *et al.*, 2013) which are typically selected by farmers for better adaptation to specific environments as well as resistance to biotic and abiotic stresses (Aci *et al.*, 2013). Characterization of those genetically heterogeneous populations using conventional and molecular tools is the most efficient way of establishing diversities for different important traits including disease resistance (Anumalla *et al.*, 2015; Prasanna, 2012). This means, developing improved varieties with required traits through plant breeding would very much depend on the availability of a wide and reliable crop genetic diversity (Abraha *et al.*, 2014; Sharma *et al.*, 2015).

2.2 Maize Lethal Necrosis Disease (MLND)

MLND was first observed in areas of South Rift Valley region of Kenya in 2011 and spread to several other places of Kenya (Wangai *et al.*, 2012b). Later, the disease was identified as MLN after serological and molecular tests were carried out on infected maize plants from Bomet County and Nakuru County in 2012 (Wangai *et al.*, 2012b). The same year, the disease was also

reported to spread to neighbouring countries of Tanzania and Uganda (Makumbi and Wangai, 2013) and later it was found in Rwanda (Adams *et al.*, 2014). The control of the disease has been reported to be difficult due to the combination of more than one virus that do not separately result into any significant symptoms (DSMZ, 2014; Xia *et al.*, 2016). MLND occurs as a result of a positive interaction between *Maize Chlorotic Mottle Virus* (MCMV) and any of the cereal viruses in the family, Potyviridae, such as *Sugarcane Mosaic Virus* (SCMV), *Maize Dwarf Mosaic Virus* (MDMV), or *Wheat Streak Mosaic Virus* (WSMV) (Adams *et al.*, 2014; Makone *et al.*, 2014; Liu *et al.*, 2016). In Eastern Africa, the disease has been reported to be caused by a combination of MCMV and SCMV infection (Gowda *et al.*, 2015; Mezzalama *et al.*, 2015; Kiruwa *et al.*, 2016). The two viruses together inflict serious damage or even completely kill infected plants (Scheets, 1998; CGIAR, 2012) and farmers in the affected areas have been reported to experience extensive to total crop loss (Wangai *et al.*, 2012b). The disease causing viruses are mainly transmitted by insects (Makone *et al.*, 2014) from plant to plant or by wind from field to field over long distances (CGIAR, 2012; Mezzalama *et al.*, 2015). MCMV is normally transmitted by thrips and beetles while SCMV is transmitted by aphids (CGIAR, 2012; Kiruwa *et al.*, 2016). The viruses can as well be transmitted from one generation to another through seed which are infected thus enhancing the possibility for wide spread of MLND (Zhang *et al.*, 2011). Infected plants show mild to severe mottling on the leaves (Gowda *et al.*, 2015; De Groote *et al.*, 2016; Kagoda *et al.*, 2016), usually starting from the base of young leaves in the whorl and extending upward towards the leaf tips (Wangai *et al.*, 2012b). Other symptoms include stunting and premature aging of the plants, dying (necrosis) of the leaf margins that progresses to the mid-rib and eventually the entire leaf (Wangai *et al.*, 2012b; Gowda *et al.*, 2015). Necrosis of young leaves in the whorl before expansion, leads to a symptom known as “dead heart” and eventually plants death (Kagoda *et al.*, 2016). In addition, infected plants often bear barren ears which are small and deformed with little or no seed set (CGIAR, 2012; Gowda *et al.*, 2015; Kagoda *et al.*, 2016).

2.3 Genetic diversity of maize accessions

Genetic diversity refers to the heritable genetic variation that occurs within populations of particular organisms (Rao and Hodgkin, 2002). The diversity in plants provides an opportunity for developing new varieties and improved cultivars with desirable characteristics (Govindaraj *et al.*, 2015; Saleh *et al.*, 2016). A number of methodologies exist for the assessment of genetic diversity in maize, those are (i) morphological characterization (Ristic *et al.*, 2014) (ii) biochemical characterization that uses electrophoresis to detect allelic variants of enzymes at gene level (Govindaraj *et al.*, 2015) (iii) pedigree that employ the extraction of genealogical information (Drinic *et al.*, 2012) and (iv) DNA molecular analyses (Sao *et al.*, 2015) such as Simple Sequence Repeats (SSR), Restriction Fragment Length Polymorphism (RFLP) (Mondini *et al.*, 2009) etc.

2.3.1 Morphological diversity

The morphological characteristics (phenotype) express the genetic constitution (genotype) of a given organism, while in other words genetic constitution give rise to what we see (Liao *et al.*, 2010; Uphoff *et al.*, 2015). However, the expression of phenotype (morphological characteristics) is always affected by the impact of environment (Anumalla *et al.*, 2015), the so called genotype x environment interaction. On the other hand, the impact by environmental interaction coupled with the expression of genetic constitution presents the advantage that can only be obtained with morphological markers (Durga *et al.*, 2015). The following equation shows the interaction;

$$P = f(G + E + (G \times E)),$$

where P stands for phenotype, G for genotype and E for environmental influences, and the interaction term $G \times E$ refers to their joint effects (Uphoff *et al.*, 2015). Genetic variation (diversity) can be evaluated using morphological characterization (Mondini *et al.*, 2009). Morphological characterization of maize is conducted through assessing plant characteristics that are given as the list of descriptors provided by the International Board for Plant Genetic Resources (IBPGR, 1991). The morphological traits that are used to evaluate maize genetic

diversity include (i). Vegetative data. (ii). Ear data. and (iii). Kernel data (Table 1). Collected data from the parameters (descriptors) as shown in Table 1 are further subjected to statistical analysis which, generate informative results that explain the behaviour of each crop accession as well as how accessions relate to each other. The results also help to identify promising accessions through potential traits evaluation. Therefore, analysis of variance (ANOVA) can be performed for all measured traits in order to test the significance of variation among accessions (Beyene *et al.*, 2005). The standardized traits mean values can be used to perform principal component (PC) and cluster analyses (to calculate similarities or dissimilarities between accessions) using softwares such as NCSS 2000 (Jerry, 2000) or NTSYS pc 2.1 (Rahman *et al.*, 2008). Further cluster analysis can be conducted on the Euclidean distance matrix with the unweighted pair group method based on arithmetic averages (UPGMA) (Beyene *et al.*, 2005).

Table 1: Some descriptors used to evaluate genetic diversity through morphological characterization (IBPGR, 1991).

| DESCRIPTO R NO. | ITEM | DESCRIPTION |
|-------------------------------|---|--|
| <u>VEGETATIVE DATA</u> | | |
| 4.1.1 | Days to tasseling | Number of days from sowing to when 50% of the plants have shed pollen |
| 4.1.2 | Days to silking | Number of days from sowing to when silks have emerged on 50% of the plants |
| 4.1.4 | Plant height [cm] | From ground level to the base of the tassel. After milk stage |
| 4.1.5 | Ear height [cm] | From ground level to the node bearing the uppermost ear. After milk stage |
| 4.1.6 | Foliage | Rating of total leaf surface. After milk stage. Observed on at least 20 representative plants |
| 4.1.7 | Number of leaves above the uppermost ear including ear leaf | Counted on at least 20 representative plants. After milk stage |
| 4.1.9 | Stem colour | Indicate up to three stem colours in the order of frequency, Observed between the two topmost ears. At flowering |
| 4.1.12 | Sheath pubescence | At flowering |
| 6.1.2 | Leaf length | From ligule to apex. Measure the leaf which subtends the uppermost ear. After flowering |
| 6.1.3 | Leaf width [cm] | Mid-way along its length. Measured on the same leaf as 6.1.2 |
| 6.1.5 | Leaf orientation | After flowering |
| 6.1.6 | Presence of leaf ligule | After flowering |
| <u>EAR DATA</u> | | |
| 6.2.2 | Ear length [cm] | Length |
| 6.2.4 | Ear diameter [cm] | Measured at the central part of the uppermost ear |
| 4.2.3 | Kernel row arrangement | Use the uppermost ear |
| 4.2.4 | Number of kernel rows | Count number of kernel rows in the central part of the uppermost |
| 6.2.5 | Cob diameter [cm] | Diameter at the center part of the uppermost ear cob |
| 6.2.6 | Rachis diameter [cm] | Diameter at the center of the inner part of the cob |
| 6.2.8 | Number of kernels per row | Count kernels of any row |
| 6.2.9 | Cob colour | Observe the colour of cob |
| 6.2.10 | Shape of uppermost ear | Observe shape |
| <u>KERNEL DATA</u> | | |
| 4.3.1 | Kernel type | Indicate up to three kernel types in the order of frequency |
| 4.3.2 | Kernel colour | Indicate up to three kernel colours in the order of frequency |
| 4.3.3 | 1000 kernel weight [g] | Adjusted to 10% moisture content |
| 6.3.1 | Kernel length [mm] | Average of 10 consecutive kernels from one row in the middle of the uppermost ear, measured with calliper |
| 6.3.2 | Kernel width [mm] | Measured on the same 10 kernels as 6.3.1 |
| 6.3.3 | Kernel thickness [mm] | Measured on the same 10 kernels as 6.3.1 |
| 6.3.4 | Shape of upper surface of kernel | Observe shape |
| 6.3.5 | Pericarp colour | Observe colour |
| 6.3.6 | Aleurone colour | Observe colour |
| 6.3.7 | Endosperm colour | Observe colour |

2.3.2 Diversity at molecular level

In the past few decades, analyzing genetic diversity based on phenotypic traits (morphological characterization) has been enhanced with the use of molecular (DNA) markers (Ristic *et al.*, 2014). This is due to the fact that discrepancies encountered with morphological markers are checked by the use of molecular markers for the purpose of obtaining variation at the DNA level (Dubreuil *et al.*, 2006). Evaluating genetic diversity of genetic resources at molecular level is essential due to the fact that morphological differences themselves are usually determined by a small number of genes and may not be representative of genetic divergence in the entire genome (Brown-Guedira *et al.*, 2000). However, the importance of these genetic resources and their potentials for selection has been constrained due to limited amount of important traits being characterized at molecular level (Rao and Hodgkin, 2002; Drinic *et al.*, 2012; Prasanna, 2012; Sood *et al.*, 2014). A number of reports on maize populations have been showing a considerable amount of variability on morphological and agronomic traits (Ihsan *et al.*, 2005) as well as at molecular level (Legesse *et al.*, 2006). This call for the efforts of ensuring that potential useful traits found in a wide range of plant genetic resources are made available to plant breeders for the purpose of crop improvement (Frese *et al.*, 2012; Maxted *et al.*, 2013).

i. Molecular marker techniques

Molecular marker techniques present the ability to detect variation at the DNA level through breeding programs and plant biotechnology (Anumalla *et al.*, 2015). PCR based molecular marker techniques have made it possible for breeders and other scientists to estimate genetic diversity through the use of different molecular markers (Arif *et al.*, 2010; Poczai *et al.*, 2013). Genetic diversity results from the genetic variation among individuals and is expressed in DNA sequences (Bindroo and Moorthy, 2014; Osawaru *et al.*, 2015). It can be categorized in terms of the number of different alleles existing in different populations, distribution of those alleles in the chromosomes, the impact these alleles have on performance and the general variability among different populations under various environmental conditions (Rao and Hodgkin, 2002; Mondini *et al.*, 2009; Bindroo and Moorthy, 2014). Some of the techniques that have been applied in molecular studies include RFLPs (Mondini *et al.*, 2009), RAPDs (Brown-Guedira *et al.*, 2000; Mondini *et al.*, 2009), AFLPs (Mondini *et al.*, 2009), and SSRs (Kumari *et al.*, 2005;

Beyene *et al.*, 2005; Mondini *et al.*, 2009; Aci *et al.*, 2013; Abraha *et al.*, 2014). However, these different marker techniques emphasize on different features (Abdel-Mawgood, 2012) and different aspects of genetic diversity (Matsuoka *et al.*, 2002; Mondini *et al.*, 2009). Therefore, different marker techniques may lead to different results and the range of variation produced can be different (Hodgkin *et al.*, 2001). Among those several marker techniques, microsatellites (SSR) have been exploited in many ways (Ignjatovic-Micic *et al.*, 2013) and specifically for characterizing genetic diversity in Maize (Reif *et al.*, 2006).

ii. Simple Sequence Repeats markers

These are microsatellites which are abundant and occur frequently and randomly (Ristic *et al.*, 2014) in eukaryotic genomes that are examined (Matsuoka *et al.*, 2002; Wan *et al.*, 2004). Irrespective of the fact that microsatellites are time consuming and costly, they actually are advantageous in terms of ease of use, high levels of inherited variation, co-dominant, reliable and highly reproducible (Rao and Hodgkin, 2002; Mondini *et al.*, 2009; Ristic *et al.*, 2014). The following constitutes major steps towards executing molecular characterization procedures:

Extraction of DNA

Extraction of DNA from sample to be analyzed is the first step for molecular marker diversity analysis (Semagn *et al.*, 2006). DNA is extracted from the leaf samples taken from young seedlings (Legesse *et al.*, 2006; Semagn *et al.*, 2006) using the CTAB procedure as explained by Saghai-Marroof *et al.* (1984). Sometimes modifications can be made depending on circumstances. Primers of a specific marker type such as SSR are selected on the basis of their genomic locations (Kumari *et al.*, 2005).

Polymerase Chain reaction (PCR)

The first step with PCR is denaturation process or melting step which separate the two DNA strands (template DNA), this step requires very high temp 94 - 98 °C (Kumari *et al.*, 2005; Legesse *et al.*, 2006). Annealing step follows that allow primers to bind to the complementary sequences on the template DNA and the temperature here ranges from 40 - 60 °C (Matsuoka *et*

al., 2002; Kumari *et al.*, 2005). The next step is elongation after the primers are bound, this requires a temperature of 72 °C (Matsuoka *et al.*, 2002; Kumari *et al.*, 2005).

Gel electrophoresis

Finally, the DNA sample is loaded into wells of agarose gel and then the gel is ran and scanned under UV light on transilluminator for interpreting the results (Yılmaz *et al.*, 2012). From the gel electrophoresis, polymorphism (variation) is expected to be determined (Senior *et al.*, 1998). Also similarity matrix is analysed by NTSYS-pc analytical package to generate hierarchical classification by the use of Unweighted Pair Group Method using Arithmetic Averages (UPGMA) (Kumari *et al.*, 2005).

2.4 Importance of landraces

Landraces are plant populations which are cultivated by local farmers that have historic origin, unique identity with no any formal crop improvement (Prasanna, 2012; Hagenblad *et al.*, 2016). They are often endowed with diverse genetic inheritance (Zeven, 1998) as well as local adaption and strong connection to traditional farming systems (Camacho *et al.*, 2005). They are typically selected by farmers for better adaptation to specific environment (Ristic *et al.*, 2014), yield potential, nutritional qualities and resistance/tolerance to biotic and abiotic stresses (Prasanna, 2012). However, domestication and selection of maize are said to cause reduced genetic diversity in the maize genome as compared with its progenitor population (Pineda-Hidalgo *et al.*, 2013). The reduction in the genetic diversity of crops represents an increase in vulnerability to new pests and diseases (De Jaramillo, 2009; Ogwu *et al.*, 2014). It has been reported that about 80% of African farmers grow only landraces since they are able to reuse the seeds in many seasons while on the other hand only 20% grow improved varieties together with landraces (Anjichi *et al.*, 2005). In Mexico, landraces occupy more than 80% of the area under maize production (Mercer and Wainwright, 2007). In Tanzania, landraces and traditional cultivars attract more attention than commercial improved cultivars (Mitawa and Marandu, 1996). In addition, between 1974 to 2000, maize researchers under maize research program in Tanzania have managed to utilize potentials from maize germplasm sourced from within and outside the country to release a number of varieties (Kirway *et al.*, 2000). Some of those were able to be adopted because they

possessed some traits preferred by the local communities besides being connected to the existing farming practices (Westengen *et al.*, 2014). The traits of those released varieties extend from plant characteristics, yield performance, disease resistance or tolerance to mention but a few. This gives an indication that maize germplasm presents an opportunity for making genetic improvement against maize production challenges. Landraces in Mexico attract great attention to both farmers and researchers (Rodriguez *et al.*, 1998). Tuxpeño maize for instance, is a Mexican landrace that was domesticated in the Oaxaca Chiapas region (Rodriguez *et al.*, 1998; Prasanna, 2010) and it is a cultivar which is very productive in the fertile lowland and has been used in the breeding programmes (Rodriguez *et al.*, 1998; Prasanna, 2010). Another example is Tuxpeño crema, a Mexican cultivar, which is characterized as a late maturing cultivar and resistant to tropical foliar diseases with white kernels, short and strong stalk (Rodriguez *et al.*, 1998). Therefore, landraces are expected to be a very important source of new and unique alleles which have not yet been well exploited (Mercer and Wainwright, 2007; Olson *et al.*, 2012). However, limited characterization data of landraces have caused difficulties to use, manage and conserve them (Rao and Hodgkin, 2002; Drinic *et al.*, 2012; Prasanna, 2012; Sood *et al.*, 2014). Another challenge with maize landraces is that many plant breeders limit their breeding research by using germplasm that contains a narrow genetic base (Prasanna, 2012). Thus, there is an urgent need to establish efficient and well organized characterization of maize germplasm for creating comprehensive information that would be useful to generate strong maize varieties against production challenges (Saad and Rao, 2001; Drinic *et al.*, 2012; Prasanna, 2012) including MLND in particular.

2.5 Conclusion

Maize continues to be the most preferred staple food and cash crop in Tanzania as well as in other parts of the world. However, maize is affected by many pathogens and some of them cause big negative impacts to its productivity. The outbreak of a new disease in East Africa called Maize Lethal Necrosis Disease (MLND) presents immediate threat to food security as well as uncertain long-term consequences. MLND infection rate reaches 100% and yields severely affected up to a complete loss of the crop. The International Maize and Wheat Improvement Center (CIMMYT) and Kenya Agriculture and Livestock Research Organization (KALRO) have

conducted joint studies from 2012 to 2014 with about 25,000 germplasm of elite inbred lines, Double Haploid lines, and hybrids against MLND under artificial inoculation and more than 95 percent of the tested materials were found to be completely susceptible. Stakeholders in maize research and development admit that expanding the range of maize germplasm for MLND screening is still needed to increase the possibility of acquiring materials with resistance or tolerance against the disease. Landraces and local cultivars in maize present a population which is genetically diverse and have been under farmer's selection for many years in terms of adaptation, plant characteristics, yield, biotic and abiotic stress tolerance or resistance. However, only 2% of the world maize germplasm collection is being absorbed in breeding programs and an important part of maize germplasm (landraces and local materials) are still in the hands of small farmers. Hence, good understanding of the performance on yield and resistance against MLND through screening a wide range of maize accessions ensure effective utilization of the genetic resources available for the fight against the current major challenges in maize production.

CHAPTER THREE

Evaluation of agronomic performance of local and improved maize varieties in Tanzania²

Mujuni Sospeter Kabululu^{1*}, Tileye Feyissa^{1,2}, Patrick Alois Ndakidemi¹

1. Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Arusha, Tanzania

2. Institute of Biotechnology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

* **For correspondence:** kabululum@nm-aist.ac.tz

Abstract

Maize (*Zea mays* L.) is one of the most important crops and yet it realizes very low average yield in Tanzania due to a number of production challenges including the use of poor varieties by most of the farmers. Availability of an efficient and well organized agronomic, morphological and genetic knowledge of maize germplasm make it easy to select the best recombinants and plan well for crosses in future breeding programs. The objective of this study was to evaluate the agronomic performance of 50 Tanzanian maize local varieties compared with 7 commercial varieties and 11 elite lines from CIMMYT in Nairobi. The genotypes were laid out in a randomized complete block design (RCBD) in three sites located in Arusha region of Tanzania in 2015. The data obtained was subjected to statistical analysis using COSTAT and STATISTICA analytical packages. These software identified significant variations among genotypes and their interactions with the environments. The GGE biplot analyses identified the best genotypes with respect to mean yield and stability. An open pollinated variety (OPV), Situka 1 and a hybrid DH 04 had generally the best performance in terms of grain yield and stability across all the 3 locations. A local cultivar TZA 2793 emerged to be the most promising landrace with overall appealing yield and stability performance. The results of this study may be a good source of new allelic diversity that could be used for developing different important elite maize materials.

Key words: *Maize, Tanzania, agronomic performance, OPV, hybrid, local cultivars, GGE biplot*

² Article published in Indian Journal of Agricultural Research (Vol. 51, Issue 3, pp 233 -238)

3.1 Introduction

Maize (*Zea mays* L.) receives much of attention worldwide because of its importance as food and feed (Guruprasad *et al.*, 2016; Dogan *et al.*, 2015). In Tanzania, maize is important as it is normally equated to the national food security (Katinila *et al.*, 1998). However, the general average yield of maize is still very low with 1.2 MT per hectare (Makurira *et al.*, 2007) as compared with the potential estimated yield of 4 to 5 MT per hectare (Otunge *et al.*, 2010). The low yield is attributed to the use of poor varieties, abiotic and biotic constraints, low soil fertility, poor agronomic management and uncertain input supply (Lisuma *et al.*, 2006). There is a need to identify adapted maize breeding materials that possess traits with high potentials to be used by maize breeders to develop superior cultivars (Sharma *et al.*, 2015). Successful plant breeding requires a more careful choice of genotypes that would be used to develop desirable recombinants (Toshimenla *et al.*, 2016). Only two percent of the world maize germplasm collection is being absorbed in breeding programs (Dowswell *et al.*, 1996) and an important part of maize germplasm is still in the hands of small-scale farmers (Carvalho *et al.*, 2004). Landraces and local cultivars in maize present a population which is genetically diverse and have been under farmer's selection for many years. Selection criteria include adaptation to local climatic conditions, plant characteristics, yield, biotic and abiotic stress tolerance or resistance (Wasala *et al.*, 2013). However, landraces are so far not efficiently being utilized because of unreliable information on agronomic as well as genetic potentials (Nass *et al.*, 1993). The objective of this study was to evaluate the agronomic performance of local cultivars in comparison to improved maize varieties in Tanzania as well as elite lines from CIMMYT in order to identify potential breeding materials.

3.2 Materials and methods

In this study, 68 accessions (Table 2) were laid out in a randomized complete block design (RCBD) with three replications. The study was conducted in three sites namely Mlangarini (S 03° 26' 12", E 036 ° 47' 13.4" at 1128 m.a.s.l.); Tengeru (S 03° 22' 30.2", E 036 ° 48' 30.2" at 1237 m.a.s.l.) and Selian (S 03° 21' 31.4", E 036 ° 37' 51.9" at 1415 m.a.s.l.).

Table 2: Maize genotypes used in the evaluation of agronomic performances in Tanzania during April to September 2015.

| SNo. | Accession name | Status | Source | SNo. | Accession name | Status | Source |
|------|--------------------|-----------------------------|--------------------|------|----------------|----------------|--------------------|
| 1 | CKDHL0500 | Elite line | CIMMYT, Nairobi | 35 | TZA3167 | Local cultivar | Genebank, Tanzania |
| 2 | CKDHL120552 | Elite line | CIMMYT, Nairobi | 36 | TZA3171 | Local cultivar | Genebank, Tanzania |
| 3 | CKSBL10205 | Elite line | CIMMYT, Nairobi | 37 | TZA3181 | Local cultivar | Genebank, Tanzania |
| 4 | CLRCY034-B | Elite line | CIMMYT, Nairobi | 38 | TZA3206 | Local cultivar | Genebank, Tanzania |
| 5 | CLRCY039 | Elite line | CIMMYT, Nairobi | 39 | TZA3310 | Local cultivar | Genebank, Tanzania |
| 6 | CLYN261 | Elite line | CIMMYT, Nairobi | 40 | TZA3536 | Local cultivar | Genebank, Tanzania |
| 7 | CML440 | Elite line | CIMMYT, Nairobi | 41 | TZA3544 | Local cultivar | Genebank, Tanzania |
| 8 | CML442 | Elite line | CIMMYT, Nairobi | 42 | TZA3585 | Local cultivar | Genebank, Tanzania |
| 9 | CML443 | Elite line | CIMMYT, Nairobi | 43 | TZA3614 | Local cultivar | Genebank, Tanzania |
| 10 | CML544 | Elite line | CIMMYT, Nairobi | 44 | TZA3837 | Local cultivar | Genebank, Tanzania |
| 11 | TZMI730 | Elite line | CIMMYT, Nairobi | 45 | TZA3914 | Local cultivar | Genebank, Tanzania |
| 12 | DEKALB (DK 8031) | Hybrid | Agro Input shop | 46 | TZA3926 | Local cultivar | Genebank, Tanzania |
| 13 | DH04 | Hybrid | Agro Input shop | 47 | TZA3958 | Local cultivar | Genebank, Tanzania |
| 14 | PIONEER (Phb 3253) | Hybrid | Agro Input shop | 48 | TZA3971 | Local cultivar | Genebank, Tanzania |
| 15 | SC403 | Hybrid | Agro Input shop | 49 | TZA4020 | Local cultivar | Genebank, Tanzania |
| 16 | SITUKA1 | OPV | Agro Input shop | 50 | TZA4164 | Local cultivar | Genebank, Tanzania |
| 17 | SITUKAM1 | OPV | Agro Input shop | 51 | TZA4203 | Local cultivar | Genebank, Tanzania |
| 18 | TMV-1 | OPV | Agro Input shop | 52 | TZA4320 | Local cultivar | Genebank, Tanzania |
| 19 | TZA163 | Local cultivar ^b | Genebank, Tanzania | 53 | TZA4351 | Local cultivar | Genebank, Tanzania |
| 20 | TZA1723 | Local cultivar | Genebank, Tanzania | 54 | TZA4574 | Local cultivar | Genebank, Tanzania |
| 21 | TZA1724 | Local cultivar | Genebank, Tanzania | 55 | TZA4667 | Local cultivar | Genebank, Tanzania |
| 22 | TZA1745 | Local cultivar | Genebank, Tanzania | 56 | TZA5102 | Local cultivar | Genebank, Tanzania |
| 23 | TZA1753 | Local cultivar | Genebank, Tanzania | 57 | TZA5129 | Local cultivar | Genebank, Tanzania |
| 24 | TZA1757 | Local cultivar | Genebank, Tanzania | 58 | TZA5138 | Local cultivar | Genebank, Tanzania |
| 25 | TZA212 | Local cultivar | Genebank, Tanzania | 59 | TZA5162 | Local cultivar | Genebank, Tanzania |
| 26 | TZA2263 | Local cultivar | Genebank, Tanzania | 60 | TZA5169 | Local cultivar | Genebank, Tanzania |
| 27 | TZA2264 | Local cultivar | Genebank, Tanzania | 61 | TZA5170 | Local cultivar | Genebank, Tanzania |
| 28 | TZA2330 | Local cultivar | Genebank, Tanzania | 62 | TZA5200 | Local cultivar | Genebank, Tanzania |
| 29 | TZA2338 | Local cultivar | Genebank, Tanzania | 63 | TZA5205 | Local cultivar | Genebank, Tanzania |
| 30 | TZA2731 | Local cultivar | Genebank, Tanzania | 64 | TZA5618 | Local cultivar | Genebank, Tanzania |
| 31 | TZA2793 | Local cultivar | Genebank, Tanzania | 65 | TZA5619 | Local cultivar | Genebank, Tanzania |
| 32 | TZA2813 | Local cultivar | Genebank, Tanzania | 66 | TZA599 | Local cultivar | Genebank, Tanzania |
| 33 | TZA2843 | Local cultivar | Genebank, Tanzania | 67 | TZA608 | Local cultivar | Genebank, Tanzania |
| 34 | TZA2904 | Local cultivar | Genebank, Tanzania | 68 | TZA93 | Local cultivar | Genebank, Tanzania |

^a OPV = Open Pollinated Varieties

^b Local cultivar = Local cultivated variety

Sowing was done in 4 rows at a spacing of 75 cm between rows and 30 cm within rows of 4 m long each. Two seeds were planted followed by thinning to one plant per hill in two weeks after sowing. At the time of sowing, the recommended dose of fertilizer (40:20:20 kg /ha NPK) was applied in the form of urea, single super phosphate and muriate of potash. Full dose of phosphorus, potash and 50% nitrogen was applied as basal dose and 50% nitrogen top dressed 30 days after sowing. The experiments were conducted during rainy season and irrigation was applied whenever necessary. Sowing at Mlangarini, Selian and Tengeru were both in April, 2015 while harvesting was done in September, 2015 for Mlangarini and October, 2015 for both Selian and Tengeru. Data were recorded on the two inner rows for days to 50% silking and anthesis (tasseling), ear diameter, ear length, number of kernel rows per ear, number of kernels per row, 1000 seed weight and grain yield per plant. The yield per plant data were collected by harvesting grains in all plants of the two inner rows and divided by the total number of plants harvested. Collected data were subjected to COSTAT and STATISTICA for analysis of variance and the correlation between parameters. Also, the analysis using GGE biplot was used to determine Genotype main effect and Genotype by Environment interaction where the variability factor was expected to be maize accessions (Genotype) and location (Environment). The Genotype plus Genotype by environment interaction (GGE biplot) analysis is a multivariate analytical technique that uses principal component analysis (PCA) to explore the relation between genotypes, environments and their interaction. It partition genotypes and genotype by environment interaction (G + GE) into principal components (Yan, 2001). Yield data were subjected to evaluation among the three groups, that is, landraces, improved varieties popularly grown in Tanzania and elite materials from CIMMYT, Nairobi, Kenya to see how they differ and perform in terms of yield.

3.3 Results and discussion

3.3.1 Analysis of variance of agronomic traits

In general, high significant ($p < 0.05$) difference among genotypes, locations as well as their interactions (Location x Genotype) were observed except for the interaction on ear length with environment which, was not significantly different (Table 3). OPV Situka 1 and hybrid DH04 emerged to be the best genotypes with respect to mean grain yield per plant with 116.01 g and

115.90 g, respectively across the three locations (Table 3). The least mean grain yield was obtained from a local cultivar TZA 2813 with 11.52 g per plant. The detailed table showing the performance of all the 68 accessions across all three locations is given in Appendix 1. The OPV maize cultivars are reported to be able to produce grain yield which is the same or a little more than hybrids (Omondi *et al.*, 2014) because they are in most cases domesticated and developed under localized marginal areas (Gudu *et al.*, 2005).

The local cultivar TZA 2813 took the longest time (100.67 days) to 50% silking while another local cultivar TZA 1724 was late maturing among all genotypes with 90.67 days to 50% anthesis (tasseling) (Table 3). On the other hand, a hybrid SC 403 was the earliest with both 63.67 days to 50% anthesis (tasseling) as well as 69.78 days to 50% silking. However, OPV's Situka 1 and Situka M1 were as well among the genotypes that took shorter time to 50% anthesis (tasseling) and silking. The number of days to anthesis or tasseling or even to silking signify maturity differences (Olaoye, 2009). The genotypes which took the shortest time to reach maturity had also high grain yield as compared with those that were late to reach maturity (Table 3). This is due to the fact that early maturing cultivars possess an opportunity to escape early occurring stresses and adapt to stresses that would occur at the end of the season (Salami *et al.*, 2007). Silking delay causes maize to become barren and it is associated with poor development of an ear during flowering (Edmeades *et al.*, 1993). This is also evidenced by the frequent occurrence of negative association between days to 50% flowering and grain yield (Bolafios and Edmeades, 1996). The current study showed that hybrid Pioneer (Phb 3253) with respect to yield related-parameters had the highest ear length, ear diameter and number of kernels per row while hybrid SC 403 had the highest 1000 seed weight (Table 3). The trend of improved varieties (hybrids) to perform better than landraces in terms of yield and other yield-related parameters had also been found by Wasala *et al.* (2013).

Table 3: Performance of maize local cultivars (TZA lines), OPV, improved varieties and elite lines from CYMMIT on grain yield and yield components.

| Genotype name | Status | Yield per plant (g) | 1000 Kernel weight (g) | Days to 50% Tasseling | Days to 50% Silking | Ear Length (cm) | Ear diameter (cm) | Number of Kernels per row | Number of Kernel rows |
|-------------------------------------|-----------------------------|---------------------|------------------------|-----------------------|---------------------|--------------------|-------------------|---------------------------|-----------------------|
| The first ten genotypes | | | | | | | | | |
| SITUKA1 | OPV ^a | 116.01±12.28 | 275.51±19.47 | 64.33±2.25 | 70.89±2.67 | 16.02±0.89 | 4.28±0.17 | 31.17±2.08 | 13.05±0.33 |
| DH04 | Hybrid | 115.90±14.48 | 273.74±15.33 | 72.11±2.05 | 75.44±2.01 | 15.58±0.41 | 4.51±0.07 | 32.65±1.75 | 12.78±0.13 |
| PIONEER | Hybrid | 112.73±17.19 | 258.10±25.60 | 69.89±1.76 | 75.89±2.04 | 17.66±0.36 | 4.64±0.05 | 36.07±0.96 | 13.83±0.18 |
| TMV-1 | OPV | 108.14±10.84 | 247.27±13.52 | 73.44±1.82 | 77.78±1.88 | 16.50±0.52 | 4.36±0.07 | 31.92±1.71 | 13.92±0.50 |
| DEKALB | Hybrid | 104.74±9.11 | 302.73±19.86 | 66.44±2.29 | 69.89±2.39 | 16.63±0.51 | 4.58±0.09 | 33.09±1.17 | 12.42±0.13 |
| TZA2793 | Local cultivar ^b | 100.46±11.22 | 260.68±10.45 | 78.33±2.38 | 84.00±2.74 | 15.86±0.39 | 4.24±0.08 | 29.36±1.47 | 12.45±0.20 |
| TZA5170 | Local cultivar | 99.80±14.55 | 285.36±26.12 | 77.44±2.46 | 83.00±2.71 | 15.14±0.54 | 4.38±0.09 | 26.77±1.55 | 11.78±0.26 |
| SITUKAMI | OPV | 99.41±14.94 | 301.99±13.68 | 64.11±2.19 | 71.56±1.89 | 15.95±0.33 | 4.43±0.08 | 30.30±1.16 | 13.19±0.22 |
| TZA2263 | Local cultivar | 97.54±17.01 | 260.73±20.96 | 80.33±2.69 | 86.11±3.29 | 15.35±0.69 | 4.37±0.12 | 28.86±1.73 | 12.77±0.14 |
| SC403 | Hybrid | 95.14±14.76 | 313.55±18.15 | 63.67±2.05 | 69.78±1.96 | 16.77±0.59 | 4.52±0.11 | 33.10±1.88 | 12.67±0.16 |
| Ten genotypes in the middle | | | | | | | | | |
| TZA3544 | Local cultivar | 63.34±7.75 | 272.89±15.38 | 78.11±2.75 | 85.67±2.82 | 14.54±0.70 | 4.21±0.06 | 23.41±1.78 | 12.47±0.18 |
| TZA3914 | Local cultivar | 61.16±7.01 | 271.82±15.96 | 79.33±2.95 | 82.89±3.28 | 14.83±0.78 | 4.26±0.11 | 22.73±1.59 | 12.25±0.40 |
| TZA4574 | Local cultivar | 60.84±8.43 | 221.23±23.26 | 81.44±2.92 | 89.44±3.50 | 13.59±0.81 | 3.67±0.17 | 25.07±2.53 | 10.90±0.36 |
| CLYN261 | CIMMYT Line | 60.46±10.82 | 232.81±24.03 | 84.56±2.75 | 85.89±2.78 | 13.01±0.55 | 3.65±0.13 | 18.96±1.82 | 13.01±0.21 |
| TZA1723 | Local cultivar | 60.36±5.02 | 265.78±15.32 | 81.89±2.84 | 89.33±2.87 | 15.11±0.49 | 3.73±0.06 | 23.90±0.96 | 11.10±0.17 |
| TZA5138 | Local cultivar | 60.13±11.55 | 213.58±11.39 | 76.78±2.02 | 83.33±2.35 | 14.18±0.48 | 4.14±0.08 | 25.37±1.66 | 13.52±0.41 |
| TZA5162 | Local cultivar | 59.20±13.58 | 219.55±25.49 | 86.56±3.31 | 93.78±3.34 | 14.34±0.39 | 3.80±0.10 | 24.17±1.75 | 11.43±0.30 |
| TZA2264 | Local cultivar | 58.28±11.09 | 250.20±14.40 | 78.44±2.26 | 84.11±2.73 | 14.03±0.43 | 4.37±0.11 | 26.81±1.28 | 13.05±0.27 |
| TZA3167 | Local cultivar | 57.66±6.25 | 272.20±14.34 | 77.33±2.49 | 83.89±2.66 | 14.72±0.63 | 3.98±0.08 | 24.62±1.08 | 11.12±0.12 |
| TZA3310 | Local cultivar | 57.39±7.29 | 238.47±13.35 | 82.72±2.52 | 90.72±2.39 | 13.83±0.36 | 3.87±0.08 | 24.57±1.95 | 11.09±0.33 |
| The last ten genotypes | | | | | | | | | |
| TZA3181 | Local cultivar | 41.26±6.98 | 269.77±27.96 | 81.22±3.78 | 89.67±4.32 | 14.25±0.89 | 3.8±0.14 | 23.15±1.47 | 10.90±0.82 |
| TZA93 | Local cultivar | 40.45±2.91 | 271.26±18.52 | 77.89±2.20 | 87.33±1.98 | 16.46±0.48 | 4.05±0.09 | 25.27±0.99 | 11.37±0.18 |
| TZA5618 | Local cultivar | 36.69±5.03 | 219.87±24.64 | 85.78±2.99 | 94.44±3.31 | 15.41±0.67 | 4.01±0.08 | 21.50±1.33 | 11.99±0.26 |
| CKSBL10205 | CIMMYT Line | 35.95±11.31 | 185.63±15.87 | 79.56±2.22 | 81.22±2.17 | 10.80±0.55 | 3.28±0.18 | 17.72±1.91 | 12.53±0.96 |
| TZA1757 | Local cultivar | 35.72±8.24 | 284.61±24.16 | 85.11±3.46 | 93.00±3.73 | 14.59±0.64 | 4.23±0.17 | 23.35±1.31 | 11.61±0.40 |
| CML544 | CIMMYT Line | 29.20±5.18 | 192.49±19.20 | 78.00±2.81 | 80.56±2.84 | 12.72±0.50 | 3.69±0.09 | 18.84±0.73 | 12.94±0.37 |
| CML440 | CIMMYT Line | 29.03±4.80 | 174.27±10.31 | 72.11±1.98 | 76.22±2.52 | 11.48±0.30 | 3.36±0.11 | 17.88±0.75 | 12.82±0.37 |
| TZA1724 | Local cultivar | 24.47±3.27 | 248.36±25.28 | 90.67±2.98 | 101.33±2.97 | 16.00±0.52 | 3.72±0.13 | 20.18±2.43 | 10.12±0.40 |
| TZMI730 | CIMMYT Line | 21.14±3.85 | 204.93±21.38 | 87.78±1.88 | 91.00±2.10 | 10.32±0.73 | 3.34±0.19 | 15.44±2.45 | 12.01±0.63 |
| TZA2813 | Local cultivar | 11.52±2.16 | 174.11±15.90 | 88.22±3.10 | 100.67±3.41 | 11.39±0.80 | 3.37±0.10 | 13.18±1.81 | 11.19±0.62 |
| Grand mean | | 62.94±1.48 | 249.59±2.59 | 78.98±0.40 | 85.70±0.45 | 14.52±0.10 | 4.01±0.02 | 24.59±0.26 | 12.21±0.06 |
| 2 WAY ANOVA (F - Statistics) | | | | | | | | | |
| Location (L) | | 49.31*** | 155.64*** | 1342.61*** | 1321.48*** | 131.61*** | 129.77*** | 113.69*** | 26.93*** |
| Genotypes (G) | | 6.59*** | 5.48*** | 30.65*** | 36.79*** | 6.92*** | 15.40*** | 10.86*** | 9.43*** |
| Interaction (L x G) | | 1.45** | 1.59*** | 1.41** | 1.64*** | 1.12 ^{ns} | 1.50** | 1.60*** | 1.98*** |

^a OPV = Open Pollinated Varieties; ^b Local cultivar = Local cultivated variety; Means in a column followed by the same alphabets are not significantly different; a letter followed by a dash and then a letter means to reduce a number of letters where that indicates a range of letters from the first to the last after the dash in alphabetical order. *** signify the statistical significance difference at a probability level of $p < 0.05$, ^{ns} stands for non significant difference

Note: Means of all parameters are ranked following the performance in terms of yield per plant.

3.3.2 Correlation between parameters

The Pearson's correlation analysis expressed significant ($p < 0.05$) relationships among eight parameters (traits) except for the relationship between number of kernel rows per an ear and 1000 kernel weight. Similar results of non-significant relationship was found by Zarei *et al.* (2012) thus supporting this study. The analysis showed a significant negative correlation between days to 50% anthesis and days to 50% silking with the rest of the parameters (Table 4). The number of days to anthesis and silking are very important because when the duration is long, it eventually causes yield and other related parameters to be low (Bolafios and Edmeades, 1996). The rest of the yield-related parameters which include ear length, ear diameter, number of kernels per row, number of kernel rows per ear, 1000 kernel weight and grain yield per plant were positively correlated to each other (Table 4). Correlation between traits and identification of fundamentally important traits which contribute to each other as well as to the ultimate goal is useful in selection during breeding programs.

Table 4: The Pearson correlation coefficient of eight parameters obtained from yield evaluation experiment of 68 maize accessions at three locations.

| | Days to 50% tasseling | Days to 50% silking | Ear length (cm) | Ear diameter (cm) | Number of kernels per row | Number of kernel rows per ear | 1000 kernel weight (g) | Yield per plant (g) |
|-------------------------------|-----------------------|---------------------|-----------------|-------------------|---------------------------|-------------------------------|------------------------|---------------------|
| Days to 50% tasseling | 1 | | | | | | | |
| Days to 50% silking | 0.944** | 1 | | | | | | |
| Ear length (cm) | -0.485** | -0.440** | 1 | | | | | |
| Ear diameter (cm) | -0.430** | -0.404** | 0.656** | 1 | | | | |
| Number of kernels per row | -0.551** | -0.527** | 0.721** | 0.740** | 1 | | | |
| Number of kernel rows per ear | -0.262** | -0.320** | 0.273** | 0.524** | 0.417** | 1 | | |
| 1000 kernel weight (g) | -0.285** | -0.241** | 0.469** | 0.532** | 0.395** | 0.033 ^{ns} | 1 | |
| Yield per plant (g) | -0.457** | -0.464** | 0.524** | 0.536** | 0.622** | 0.310** | 0.401** | 1 |

** signify the statistical significance difference at a probability level of $p < 0.05$, ^{ns} stands for non significant different

3.3.3 GGE biplot analysis

The first two principal components (PC1 and PC2) explained 91.2% of the total variance given by the 68 genotypes across the three locations (Fig. 1). PC1 accounted for 69.5% and PC2 21.7%, where PC1 explained genotype productivity and PC2 expressed genotype stability as explained by Yan *et al.* (2000). That means Genotypes having PC1 reading less than 0 are likely

to be high yielding and the vice versa is also true. On the other hand, genotypes with PC2 values close to zero are more stable (Hugos and Abay, 2013).

i. Which win where?

Fig. 1 displays a polygon which is obtained through joining points where genotypes are located furthest from origin of the biplot while other remaining genotypes are contained within the polygon (Yan and Tinker, 2006). The lines extending from the biplot origin toward points where perpendicular lines are formed gives out divisions of what is called sectors. The winning genotypes appear on the corners (vertex) of respective sectors. Thus, Pioneer (Phb 3253) emerged to be the best on yield at Selian experimental site while Situka 1 gave the best yield at both Tengeru and Mlangarini sites. In addition, TZA 2263, TZA 5162, TZA 2813, TZA 1724 and TZA 4320 were genotypes that expressed general adaptation with no specific location (Fig. 1).




Figure 1: The “which win where” feature of the GGE biplot for maize grain yield of 68 genotypes in three locations, where g stands for genotypes name.

ii. Genotypes performance against an ideal genotype

In Fig. 2, the 68 maize genotypes were ranked in relation to mean yield as well as stability with reference to an ideal genotype. The ideal genotype is found at the center of the inner circle as a point along the single arrow Average Environment Coordination (AEC) x - axis (Fig. 2) (Yan and Tinker, 2006). The point (an ideal genotype) possesses the longest vector with zero value across the double arrow axis. Thus, any genotype found near to the point of an ideal genotype can be considered promising than others. Therefore, Situka 1 and hybrid DH 04 were found to be the most desirable genotypes than others because they had the highest yield and still were more stable. Genotypes TMV 1, TZA 2793 and SC 403 could still be considered desirable while hybrid Pioneer (Phb 3253) was among the higher yielding genotypes but less stable across environments. On the other hand, TZA 2813 gave the least overall mean yield and TZA 4320 was the most unstable genotype (Fig. 2).




Figure 2: The average environment coordination (AEC) graphic that rank genotypes relative to an ideal genotype (the center of the concentric circles) through a GGE biplot for maize grain yield of 68 genotypes at three locations.

iii. Test environment evaluation

Fig. 3 presents the test-environment evaluation of the three locations where the line that joins biplot origin and any location, corresponds to the standard deviation derived from the mean of the genotypes in a certain environment (Yan and Tinker, 2006). It shows Tengeru to be more discriminative than the others. This means that Tengeru was able to provide much information about differences among genotypes than other locations. In line with explanation by Mehari *et al.* (2015), Tengeru was more strongly correlated to Mlangarini as the angle between them was small, while the angle between both Tengeru and Mlangarini with Selian was bigger and hence not closely related to them. Therefore, the biplot identified Tengeru to be the test environment that is close to an ideal environment which is eventually defined to be both discriminating and representative (Yan *et al.*, 2007).




Figure 3: The GGE biplot displaying the relation among experimental locations, their discriminating ability and representativeness through evaluation of 68 genotypes.

3.4 Conclusion

The significant differences expressed among the 68 genotypes and the interactions among those genotypes and environments suggest that genotypes were different (variable) from each other and they responded differently in different environments. This variability is desirable for use in breeding programs to develop improved maize varieties required by the farmers for specific traits and even for specific location. Successful plant breeding then requires careful selection of genotypes that possess desirable traits for best combinations. The correlations between traits are also another important knowledge that helps on predicting the required performance in terms of a certain trait. For example, the negative correlation between yield and flowering parameters help to have an indirect selection of higher yielding genotypes through flowering behaviour. The determination of best genotypes also requires knowing the location with high discriminative power, meaning the location which is able to provide much information about differences among genotypes. This study showed that an OPV Situka 1 and a hybrid DH 04 were generally the best performing genotypes in terms of grain yield and stability as well as for other yield-related parameters across all the three locations. However, TZA 2793 was a local cultivar that expressed promising performance for yield. The maize landraces have always been considered less productive than other improved varieties, but they present an important source of genetic variability that can be exploited during a search for genes against biotic and abiotic stresses. Therefore, identifying potential traits as well as genotypes possessing those important traits make it possible to plan for crosses and obtain required characteristics for future breeding programs.

CHAPTER FOUR

Genetic diversity of maize landraces from Tanzania as compared with commercial improved varieties and elite lines through morphological characterization³

Mujuni Sospeter Kabululu¹, Tileye Feyissa^{1,2}, Patrick Alois Ndakidemi^{1*}

¹Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania

²Institute of Biotechnology, Addis Ababa University, Addis Ababa, Ethiopia

***Corresponding email:** patrick.ndakidemi@nm-aist.ac.tz

Abstract

Maize production challenges require well-known genetic diversity that are well adapted to geographic sites to ensure effective improvement. This study aimed at conducting morphological evaluation for 50 maize landraces from Tanzania compared with 7 commercial varieties and 11 elite lines from CIMMYT in Kenya. The experiments were laid out in randomized complete block design (RCBD) at three locations in Arusha region, Tanzania. Data were collected on 19 quantitative and 12 qualitative traits that were subjected to analysis of variance, descriptive and multivariate statistics. Significant variations ($p < 0.05$) were observed for all traits while higher contribution for accessions variability were found with yield, a thousand kernel weight, flowering traits, kernel, ear and vegetative plant characteristics. Commercial varieties were characterized by significant yield (107.4g per plant) and yield related parameters namely a thousand seed weight, number of rows per an ear, ear diameter and ear length. Also early days to tasseling and silking of 67.7 and 73 days, respectively. CIMMYT elite lines were characterized by significant low plant and ear height of 138.9cm and 50.6cm, respectively as well as flint kernel type. Landraces were more diverse in every trait evaluated with significant long anthesis-silking interval of 7.5 days and large ear height of 95.9cm. Some landraces such as TZA 2793 and TZA 5170 expressed significant traits that would be tapped for further crop improvement. Other landraces clustered themselves irregularly in terms of their collection sites within their

³ Article published in International Journal of Biosciences (*Vol. 10, Issue 3, pp 309 - 322*)

major group due to selection and exchange of seeds. Thus, farmers as custodians of landraces are supposed to be involved in a systematic selection and breeding.

Key words: Genetic diversity, Maize, Landraces, Morphological characterization, Improved varieties

4.1 Introduction

Maize (*Zea mays* L.) sustains a huge population in the world (Romay *et al.*, 2013) and is even equated to the national food security in Tanzania (Katinila *et al.*, 1998). However, its average yield is still very low with 1.2 MT per hectare in Tanzania as compared with the estimated potential yields of 4 to 5 MT per hectare (Moshi *et al.*, 1990; Otunge *et al.*, 2010). Low yield has been attributed to factors such as lack of quality inputs (eg seeds, fertilizer), drought, pests and diseases. Nevertheless, maize is a crop which is potentially diverse in terms of phenotypic and genetic characters (Whitt *et al.*, 2002). The genetic variation of maize constitute a very important package for breeding (Prasanna, 2010; Yao *et al.*, 2007) which requires the availability of desirable characters for maize crop improvements (Ristic *et al.*, 2013). However, for the past decades, breeding in maize have been concentrated in short breeding programs that uses inbred lines, elite lines and breeder materials (Cömertpay, 2012; Yao *et al.*, 2007). These materials are in most cases uniform such that for a long time they have caused the existence of a narrow based genetic background (Shiri *et al.*, 2014; Yao *et al.*, 2007). The narrow based genetic background has always been coupled with genetic erosion and habitat alteration that resulted in an increased sensitivity to new pathogenic races as well as decreased resistance and tolerance to environmental extremes (Prasanna, 2010). Germplasm are heterogeneous in nature and are open pollinated with a wide range of adaptability to an extensive range of environmental variability (Rahman *et al.*, 2008). Maize landraces are reported to be genetically heterogeneous populations which have been selected by farmers for environmental adaptability (Aci *et al.*, 2013; Ignjatovic *et al.*, 2013). They can also be used to explore for resistance and tolerance against biotic and abiotic environmental stress factors (Molin *et al.*, 2013). Salami *et al.* (2015) found significant morphological variation with Benin local and improved maize varieties on all traits with distinctive potential highlight on early maturity and sensitivity to maize streak virus. Traits such as plant growth, tassel characteristics and yield had a significant contribution to phenotypic

variation between maize landraces that were assessed by Ristic *et al.* (2014). Significant amount of variability was observed by Rahman *et al.* (2008) from the morphological traits evaluated in maize population from Pakistan. Italian maize landraces showed significant morphological variation on earliness, plant architecture traits, tassel, ear and kernel characteristics (Hartings *et al.*, 2008). Asare *et al.* (2016) suggested that maize landraces presents a significant genetic diversity reserve of important morphological characteristics on phenology, plant growth, grain yield, and leaf photosynthesis that reflects farmer preferences and worth for maize crop improvement.

In Tanzania, Nestory and Reuben (2016) evaluated maize landraces from northern part of the country and obtained high traits variability that serve as an opportunity for enhancing genetic improvement to maize germplasm required by the community. On the other hand, Bucheyeki (2012) evaluated maize landraces from Tanzania and identified potential sources of northern leaf blight disease resistance. However, Tanzania still holds a vast majority of germplasm that remain marginally exploited and are of great importance for food, adaptability, resistance to pests and diseases as well as for quality attributes (Ngwediagi *et al.*, 2009).

The general decreasing trend in maize production and yield in Tanzania is caused by recurrent abiotic and biotic stresses (Bucheyeki, 2012) and recently Tanzania and East Africa in general has been hit hard by another new deadly disease called Maize Lethal Necrosis disease (Kabululu *et al.*, 2017; Kiruwa *et al.*, 2016; Wangai *et al.*, 2012b). Thus detailed characterization of landraces and other germplasm is required to establish a gene pool for crop improvement (Drinic *et al.*, 2012; Obeng-Antwi, 2012; Saad and Rao, 2001). The objective of this study was therefore to evaluate the genetic diversity of maize accessions from Tanzania through morphological characterization in order to establish the existing genetic diversity worth for maize crop improvement.

4.2 Materials and methods

4.2.1 Seed materials

The 68 accessions used in this study included 50 landraces collected in Tanzania, seven improved commercial varieties in Tanzania and eleven elite lines from CIMMYT in Nairobi,

Kenya (Table 5). Seeds of the 50 landraces of maize were obtained from the National Plant Genetic Resources Centre (NPGRC) in Arusha, Tanzania and were collected from different parts of the country (Fig. 4). The sampling of the collected maize accessions was done considering a wide distribution over the country.

Table 5: List of maize accessions and their source as used in the genetic diversity evaluation study in Tanzania.

| SNo. | Accession | Source | SNo. | Accession | Source |
|-------------|--------------------|--------------------|-------------|------------------|---------------|
| 1 | CKDHL0500 | CIMMYT | 35 | TZA3171 | Kigoma |
| 2 | CKDHL120552 | CIMMYT | 36 | TZA3181 | Kigoma |
| 3 | CKSBL10205 | CIMMYT | 37 | TZA3206 | Tabora |
| 4 | CLRCY034-B | CIMMYT | 38 | TZA3310 | Tabora |
| 5 | CLRCY039 | CIMMYT | 39 | TZA3536 | Morogoro |
| 6 | CLYN261 | CIMMYT | 40 | TZA3544 | Morogoro |
| 7 | CML440 | CIMMYT | 41 | TZA3585 | Mtwara |
| 8 | CML442 | CIMMYT | 42 | TZA3614 | Mtwara |
| 9 | CML443 | CIMMYT | 43 | TZA3837 | Mtwara |
| 10 | CML544 | CIMMYT | 44 | TZA3914 | Mara |
| 11 | DEKALB (DK8031) | Commercial variety | 45 | TZA3926 | Mara |
| 12 | DH04 | Commercial variety | 46 | TZA3958 | Mara |
| 13 | PIONEER (Phb 3253) | Commercial variety | 47 | TZA3971 | Mara |
| 14 | SC403 | Commercial variety | 48 | TZA4020 | Mwanza |
| 15 | SITUKA1 | Commercial variety | 49 | TZA4164 | Kagera |
| 16 | SITUKAM1 | Commercial variety | 50 | TZA4203 | Mwanza |
| 17 | TMV-1 | Commercial variety | 51 | TZA4320 | Kagera |
| 18 | TZA163 | Mtwara | 52 | TZA4351 | Kagera |
| 19 | TZA1723 | Njombe | 53 | TZA4574 | Mwanza |
| 20 | TZA1724 | Njombe | 54 | TZA4667 | Mwanza |
| 21 | TZA1745 | Njombe | 55 | TZA5102 | Tanga |
| 22 | TZA1753 | Mbeya | 56 | TZA5129 | Tanga |
| 23 | TZA1757 | Mbeya | 57 | TZA5138 | Tanga |
| 24 | TZA212 | Mbeya | 58 | TZA5162 | Tanga |
| 25 | TZA2263 | Lindi | 59 | TZA5169 | Tanga |
| 26 | TZA2264 | Lindi | 60 | TZA5170 | Tanga |
| 27 | TZA2330 | Lindi | 61 | TZA5200 | Tanga |
| 28 | TZA2338 | Mtwara | 62 | TZA5205 | Tanga |
| 29 | TZA2731 | Morogoro | 63 | TZA5618 | Manyara |
| 30 | TZA2793 | Morogoro | 64 | TZA5619 | Manyara |
| 31 | TZA2813 | Tanga | 65 | TZA599 | Singida |
| 32 | TZA2843 | Tanga | 66 | TZA608 | Singida |
| 33 | TZA2904 | Ruvuma | 67 | TZA93 | Rukwa |
| 34 | TZA3167 | Kigoma | 68 | TZMI730 | CIMMYT |

Also seven improved commercial maize varieties were obtained from the agro-input shops in Arusha and CIMMYT inbred lines were sourced from CIMMYT in Nairobi, Kenya. Both improved commercial maize varieties and CIMMYT inbred lines were used in this study as checks.

4.2.2 Field location and experimental design

Morphological characterization experiment was laid out in a randomized complete block design with three replications. The experiment was conducted in three locations in Arusha region as follows; Mlangarini at S 03° 26' 12", E 036° 47' 13.4" with elevation of 1128 m.a.s.l.; Tengeru at S 03° 22' 30.2", E 036° 48' 30.2" with elevation of 1237 m.a.s.l and Selian at S 03° 21' 31.4", E 036° 37' 51.9" with elevation of 1415 m.a.s.l. Maize accessions were planted in rows per each plot at spacing of 75 cm between rows and 30 cm within rows. Each row was 4 meters long and with an approximation of 44,444 plants population per hectare. The seeds were planted during the rainy season in 2015 where irrigation was applied whenever necessary to provide an optimum growing condition as possible. Two seeds were planted per hill followed by thinning to one plant per hill in two weeks after sowing. At planting, NPK (40:20:20) was applied at a rate of 100 kg N per hectare and top dressing with Urea (46%) were applied later at a rate of 100 kg N per hectare. Data collection generally included: vegetative, ear and kernel characteristics (Table 6). The data were collected according to the descriptor list by IBPGR (1991). Yield per plant data were collected by harvesting grains on the two inner rows and divided by the total number of plants harvested.

4.2.3 Data analysis

Descriptive statistics on means, minimum, maximum, standard errors and coefficient of variation using STATISTICA 8.0 were obtained for quantitative traits while one-way analysis of variance (ANOVA) was applied for significance test on morphological differences among maize accessions. Box and whisker plot was also used to show variability within quantitative characters. Cluster analysis was performed based on average linkage method through Genstat discovery edition 4 to generate similarity and dissimilarity among accessions and eventually comparing between groups of accessions clustered together. Principle component analysis was

done to identify traits with significant contribution to the overall variation within each principle component.




Figure 4: Map of Tanzania showing the collection sites of the 50 maize landraces

Table 6: Some descriptors used to evaluate genetic diversity through morphological characterization (IBPGR, 1991).

| ITEM | DATA COLLECTION PROCEDURE | |
|--|---------------------------|---|
| <u>Vegetative data</u> | | |
| Days to tasseling | D50T | Number of days from sowing to when 50% plants shed pollen |
| Days to silking | D50S | Number of days from sowing to 50% plants with silks |
| Anthesis-Silking Interval | ASI | Difference between days to silking and anthesis stages |
| Plant height (cm) | PH | Ground level to the base of the tassel. After milk stage |
| Ear height (cm) | EH | Ground to the node at the uppermost ear. After milk stage |
| Foliage | FG | Rating of total leaf surface. After milk stage, on 20 plants |
| Number of leaves above the uppermost ear | NLAUME | Counted on at least 20 plants. After milk stage |
| Stem colour | SC | Observed between the two topmost ears. At flowering |
| Sheath pubescence | SP | The hairy condition of leaf base encasing the stem of a plant |
| Leaf length | LL | From ligule to apex of the uppermost ear leaf . After flowering |
| Leaf width (cm) | LW | Mid-way along its length. Measured on the same leaf |
| <u>Ear data</u> | | |
| Ear length (cm) | EL | Measured from the base to the tip of the uppermost ear |
| Ear diameter (cm) | ED | Measured at the central part of the uppermost ear |
| Kernel row arrangement | KRA | Pattern and arrangement of rows of the uppermost ear |
| Number of kernel rows | NKR | Counting kernel rows in the central part of the uppermost ear |
| Cob diameter (cm) | CD | Mid-way of cob length |
| Rachis diameter (cm) | RD | Diameter of the inner part of the cob |
| Number of kernels per row | NK_R | Count number of kernel in a single row of the uppermost ear |
| Cob colour | CC | Rating colours of the cobs |
| Shape of uppermost ear | SOU ME | Determining the shape through observation |
| <u>Kernel data</u> | | |
| Kernel type | KT | Indicate up to three kernel types in the order of frequency |
| Kernel colour | KC | Indicate up to three kernel colours in the order of frequency |
| 1000 kernel weight (g) | 1000KW | Adjusted to 10% moisture content |
| Kernel length (mm) | KL | Average of 10 kernels from the row in the middle |
| Kernel width (mm) | KW | Measured on the same 10 kernels |
| Kernel thickness (mm) | KTH | Measured on the same 10 kernels |
| Shape of upper surface of kernel | SOUSOK | Determining the shape through observation |
| Endosperm colour | EC | Colour of the tissue inside the seeds |
| Yield per plant (g) | Y_P | Grain yield per plant in grams |

4.3 Results

4.3.1 Variability in quantitative characters

The descriptive statistics and analysis of variance of the 19 quantitative traits revealed a significant ($p < 0.05$) variation of all the traits among landraces (Table 7). Significant variation was observed in commercial varieties except for leaf width, cob diameter, rachis diameter, kernel width, kernel thickness and yield per plant. The CIMMYT elite lines had also significant traits variability except for only ear diameter and cob diameter. Significant coefficient of variation was observed with anthesis-silking interval as well as with yield per plant. The box and Whisker plot farther displayed the performance and variability among the groups of maize accessions, where the overall grain yield per plant was higher with commercial varieties, but high variability is seen

with landraces portrayed by the range of non outliers values (Fig. 5A). This was also observed with the weight of a thousand seed weight (Fig. 5B) where commercial varieties generated heavier seeds than other groups, but also the variability of this trait in landraces is more than other groups. Landraces showed high variation in number of kernel rows per ear while commercial varieties exhibited higher variability for number of kernels per row than other groups (Fig. 5D and E). For flowering behaviour, commercial varieties flowered earlier as compared with other groups that had almost the same median value. Landraces showed high variation in flowering as well as for plant height when compared with other groups (Fig. 5C and F).

Table 7: Mean values with standard errors, minimum and maximum (range), and coefficient of variation of the 19 quantitative morphological descriptors generated from the performance of the three groups of accessions.

| ^aDescriptor | Landraces | | | Commercial varieties | | | CIMMYT lines | | |
|-------------------------------|--------------------------|--------------|-----------|-----------------------------|--------------|-----------|--------------------------|--------------|-----------|
| | Mean ± SE | Range | CV | Mean ± SE | Range | CV | Mean ± SE | Range | CV |
| D50T | 80.2±0.4 ^{***} | 61.0-94.7 | 6.6 | 67.7±0.9 ^{***} | 62.3-75.0 | 6.0 | 80.6±0.9 ^{***} | 70.0-91.7 | 6.5 |
| D50S | 87.7±0.5 ^{***} | 74.0-108.3 | 7.4 | 73±0.7 ^{***} | 69.0-79.7 | 4.5 | 84.6±0.9 ^{***} | 73.7-94.3 | 6.2 |
| ASI | 7.5±0.2 ^{***} | 2.0-15.7 | 32.7 | 5.3±0.4 ^{***} | 2.0-7.7 | 32.5 | 4.0±0.4 ^{***} | 0.3-8.0 | 52.7 |
| NLAUME | 6.5±0.0 ^{***} | 5.6-7.7 | 5.4 | 6.7±0.1 ^{***} | 6.1-7.1 | 4.5 | 6.6±0.1 ^{***} | 5.2-7.6 | 10.3 |
| PH (cm) | 205.4±1.8 ^{***} | 162.1-275.0 | 11.0 | 211.8±3.4 ^{***} | 187.3-239.1 | 7.3 | 138.9±3.6 ^{***} | 96.6-172.9 | 15.0 |
| EH (cm) | 95.9±1.4 ^{***} | 59.6-156.8 | 18.1 | 78.7±2.4 ^{***} | 61.9-100.2 | 14.1 | 50.6±1.6 ^{***} | 29.8-65.6 | 18.7 |
| LL (cm) | 80.1±0.4 ^{***} | 69.2-95.7 | 6.2 | 83.2±0.8 ^{***} | 74.2-90.2 | 4.7 | 70.7±1.3 ^{***} | 59.1-85.5 | 10.8 |
| LW (cm) | 9.1±0.1 ^{***} | 7.5-12.0 | 7.5 | 9.7±0.1 ^{ns} | 9.1-11.1 | 5.3 | 8.7±0.2 ^{***} | 7.0-11.0 | 12.4 |
| EL (cm) | 14.6±0.1 ^{***} | 10.9-18.4 | 8.8 | 16.4±0.2 ^{***} | 15.1-18.2 | 4.5 | 12.7±0.3 ^{***} | 9.4-20.0 | 15.4 |
| ED (cm) | 4.8±0.0 ^{***} | 3.6-6.2 | 9.1 | 5.2±0.1 ^{***} | 4.3-6.1 | 7.4 | 4.4±0.1 ^{ns} | 3.3-5.4 | 12.9 |
| CD (cm) | 3.0±0.0 ^{***} | 2.3-4.0 | 10.6 | 3.1±0.1 ^{ns} | 2.8-3.8 | 8.1 | 2.8±0.1 ^{ns} | 2.3-3.6 | 10.6 |
| RD (cm) | 1.6±0.0 ^{***} | 1.2-3.6 | 14.5 | 1.6±0.0 ^{ns} | 1.5-1.8 | 4.4 | 1.4±0.0 ^{***} | 1.1-1.7 | 11.8 |
| NK_R | 24.7±0.3 ^{***} | 12.6-33.3 | 15.1 | 32.6±0.4 ^{***} | 29.6-37.0 | 6.3 | 19.1±0.5 ^{***} | 14.2-26.5 | 14.6 |
| NKR | 12.0±0.1 ^{***} | 9.1-14.7 | 9.0 | 13.1±0.1 ^{***} | 12.3-14.8 | 5.2 | 12.6±0.2 ^{***} | 9.0-14.8 | 9.0 |
| KL (cm) | 1.0±0.0 ^{***} | 0.8-1.2 | 6.7 | 1.1±0.0 ^{***} | 1.0-1.3 | 6.3 | 0.9±0.0 ^{***} | 0.7-1.1 | 11.3 |
| KW (cm) | 1.0±0.0 ^{***} | 0.8-1.4 | 7.4 | 1.0±0.0 ^{ns} | 0.9-1.2 | 6.6 | 0.8±0.0 ^{***} | 0.7-0.9 | 5.5 |
| KTH (cm) | 0.5±0.0 ^{***} | 0.4-0.9 | 12.8 | 0.5±0.0 ^{ns} | 0.4-0.5 | 4.8 | 0.6±0.0 ^{***} | 0.5-0.8 | 11.1 |
| 1000KW (g) | 256.2±2.7 ^{***} | 160.0-340.0 | 12.9 | 281.8±6.5 ^{***} | 237.0-334.4 | 10.5 | 199.1±4.4 ^{***} | 148.2-257.3 | 12.6 |
| Y/P (g) | 60.6±1.7 ^{***} | 9.8-133.7 | 34.3 | 107.4±4.8 ^{ns} | 50.1-136.6 | 20.4 | 45.2±3.0 ^{***} | 11.5-85.9 | 38.4 |

^a abbreviation of the descriptors are defined and explained in Table 2

^{***} stands for significant difference at P <0.05

^{ns} not significant different




Figure 5: Box plots displaying variability in morphological descriptors given by the three different groups of CIMMYT inbred lines, commercial/Improved varieties and landraces from Tanzania. Small empty boxes inside big boxes represent median, big vertical boxes show a range of values falling between 25% and 75% and the vertical lines cover the range of non-outliers.

4.3.2 Qualitative characters

The qualitative data of the 68 accessions were categorized into three groups and they showed landraces to be more diverse as compared to the commercial varieties and CIMMYT lines (Table 8). This means the landraces were distributed in classes of all the 12 qualitative traits within the accessions while for the other groups the accessions distribution was not in all classes of each trait. The percentage distribution of accessions at each trait differentiated the three groups of accessions in terms of foliage (rating of total leaf surface), where landraces were mostly characterized as intermediate (44.9%), commercial varieties were large (63.5%) while CIMMYT

Table 8: The 12 qualitative morphological descriptors with their percentage (%) frequency accessions distribution as generated from the performance of three different groups within 68 maize accessions used in this study.

| Morphological descriptor | Descriptor | Landraces | Commercial Varieties | CIMMYT Lines |
|----------------------------------|---------------------|------------------|-----------------------------|---------------------|
| Stem colour | Green | 95.11 | 100.00 | 100.00 |
| | Purple | 4.89 | 0.00 | 0.00 |
| Sheath pubescence | Sparse | 1.11 | 4.76 | 3.03 |
| | Intermediate | 94.22 | 85.71 | 82.83 |
| Foliage | Dense | 4.67 | 9.52 | 14.14 |
| | Small | 18.67 | 12.70 | 70.71 |
| | Intermediate | 44.89 | 23.81 | 26.26 |
| Tassel type | Large | 36.44 | 63.49 | 3.03 |
| | Primary | 3.56 | 0.00 | 23.23 |
| Tassel size | Primary-secondary | 96.44 | 100.00 | 76.77 |
| | Small | 4.44 | 6.35 | 70.71 |
| Cob colour | Medium | 48.44 | 49.21 | 25.25 |
| | Large | 47.11 | 44.44 | 4.04 |
| | White | 96.00 | 100.00 | 100.00 |
| Shape of uppermost ear | Red | 3.33 | 0.00 | 0.00 |
| | Purple | 0.67 | 0.00 | 0.00 |
| | Cylindrical | 0.67 | 0.00 | 3.03 |
| Shape of upper surface of Kernel | Cylindrical-conical | 6.22 | 0.00 | 17.17 |
| | Conical | 93.11 | 100.00 | 79.80 |
| | Shrunken | 3.33 | 41.27 | 9.09 |
| Kernel row arrangement | Indented | 23.11 | 33.33 | 12.12 |
| | Level | 1.33 | 25.40 | 26.26 |
| | Rounded | 71.78 | 77.78 | 3.03 |
| | Pointed | 0.44 | 0.00 | 0.00 |
| | Regular | 94.67 | 71.43 | 89.90 |
| Kernel type | Irregular | 4.00 | 0.00 | 10.10 |
| | Straight | 1.33 | 28.57 | 0.00 |
| | Semi-floury | 5.78 | 20.63 | 7.07 |
| | Dent | 55.78 | 53.97 | 28.28 |
| | Semi-dent | 32.89 | 14.29 | 19.19 |
| Kernel colour | Semi-flint | 5.56 | 11.11 | 41.41 |
| | Flint | 0.00 | 0.00 | 4.04 |
| | White | 84.44 | 100.00 | 67.68 |
| | Yellow | 9.33 | 0.00 | 28.28 |
| | Purple | 0.44 | 0.00 | 0.00 |
| Endosperm colour | Variegated | 2.67 | 0.00 | 2.02 |
| | White cap | 1.33 | 0.00 | 2.02 |
| | Red | 1.78 | 0.00 | 0.00 |
| | White | 94.44 | 100.00 | 78.79 |
| | Pale yellow | 0.44 | 0.00 | 0.00 |
| | Yellow | 4.67 | 0.00 | 21.21 |
| | White cap | 0.44 | 0.00 | 0.00 |

lines were small (70.7%). With regard to tassel size, landraces and commercial varieties were medium with 48.4% and 49.2%, respectively, while CIMMYT lines were small with 70.7%. On

4.3.3 Principal components analysis (PCA)

The analyses of principal components for the 25 morphological traits are shown in Table 9. The first six components expressed 78.36% of the total variation and each had an eigenvalue of more than one. The first principal component (PC1) in particular accounted for 33.36% of the total morphological variation of the studied traits. Morphological traits that highly contributed to the PC1 include leaf length, ear length, ear diameter, number of kernels per row, kernel length, 1000 kernel weight, and yield per plant. Principal component two (PC2) accounted for 17.91% of the total variation and was highly influenced by days to 50% tasseling and silking, anthesis silking interval, plant height, ear height, kernel width and thickness. Morphological traits that had high contributions to Principal component three (PC3), which accounted for 8.51% variations, were number of leaves above uppermost ear, cob diameter, rachis diameter and number of kernel rows. The fourth component (PC4) was influenced by shape of uppermost ear, kernel colour, leaf width and contributed 7.94% of the total variation. The fifth component had variability contribution of 5.76% as caused by kernel row arrangement and endosperm colour. The sixth component contributed 4.89% variation given by shape of upper surface of kernel and kernel type. The PCA plot further characterized the three groups of accessions differently with specific traits discriminating them on a plotted plane (Fig. 6). The commercial varieties were grouped on the upper left hand side quadrant, CIMMYT lines grouped themselves on the upper right hand side quadrant. Landraces were mostly scattered along the origin of the plane and to all the quadrants.

Table 9: Proportions of variability contributions given by the 25 morphological traits in different principle components.

| Trait | PC 1 | PC 2 | PC 3 | PC 4 | PC 5 | PC 6 |
|--------------------------------------|---------------|---------------|--------------|--------------|--------------|--------------|
| Days to 50% tasseling | 0.177 | -0.299 | 0.282 | 0.020 | 0.159 | -0.162 |
| Days to 50% silking | 0.149 | -0.369 | 0.220 | -0.007 | 0.113 | -0.161 |
| Anthesis Silking Interval | -0.003 | -0.374 | -0.059 | -0.079 | -0.072 | -0.079 |
| Shape of uppermost ear | -0.042 | -0.092 | -0.107 | 0.478 | -0.421 | -0.040 |
| Shape of upper surface of kernel | 0.180 | -0.169 | -0.057 | -0.124 | -0.166 | 0.325 |
| Kernel row arrangement | -0.068 | 0.116 | -0.071 | -0.287 | 0.478 | -0.202 |
| Kernel type | 0.177 | 0.054 | 0.201 | 0.091 | -0.067 | 0.508 |
| Kernel colour | 0.029 | 0.067 | -0.212 | 0.278 | 0.220 | 0.172 |
| Endosperm colour | 0.088 | 0.062 | 0.064 | 0.359 | 0.407 | 0.378 |
| Number of leaves above uppermost ear | -0.003 | 0.062 | 0.401 | 0.309 | -0.315 | -0.194 |
| Plant height (cm) | -0.247 | -0.286 | 0.033 | 0.110 | 0.045 | -0.027 |
| Ear height (cm) | -0.156 | -0.378 | 0.059 | 0.020 | 0.074 | -0.092 |
| Leaf length (cm) | -0.231 | -0.177 | 0.173 | 0.193 | 0.215 | -0.076 |
| Leaf width (cm) | -0.184 | 0.003 | 0.117 | 0.393 | 0.259 | -0.047 |
| Ear length (cm) | -0.270 | -0.095 | -0.185 | 0.033 | 0.113 | 0.189 |
| Ear diameter (cm) | -0.324 | -0.006 | 0.149 | -0.094 | -0.049 | 0.050 |
| Cob diameter (cm) | -0.258 | -0.027 | 0.312 | -0.186 | -0.077 | 0.184 |
| Rachis diameter (cm) | -0.179 | -0.028 | 0.370 | -0.247 | -0.082 | 0.324 |
| Number of kernels per row | -0.307 | 0.104 | -0.100 | 0.057 | -0.037 | 0.007 |
| Number of kernel rows | -0.099 | 0.281 | 0.416 | -0.070 | 0.093 | 0.000 |
| Kernel length (cm) | -0.320 | 0.026 | -0.100 | -0.043 | -0.064 | -0.117 |
| Kernel width (cm) | -0.198 | -0.277 | -0.183 | -0.033 | -0.094 | 0.193 |
| Kernel thickness (cm) | 0.164 | -0.313 | -0.001 | 0.009 | 0.171 | 0.197 |
| 1000 Kernel weight (g) | -0.262 | -0.100 | -0.180 | -0.115 | -0.054 | 0.201 |
| Yield per plant (g) | -0.279 | 0.166 | -0.014 | 0.152 | 0.055 | 0.005 |
| Eigenvalues | 8.34 | 4.48 | 2.13 | 1.99 | 1.44 | 1.22 |
| Total variance (%) | 33.36 | 17.91 | 8.51 | 7.94 | 5.76 | 4.89 |
| Cumulative total variance (%) | 33.36 | 51.27 | 59.77 | 67.71 | 73.47 | 78.36 |

4.3.4 Cluster analysis

The dendrogram of the 68 maize accessions evaluated based on average linkage analysis is presented in Fig. 7. The combined analysis was generated from the 19 quantitative and 12 qualitative traits. The dendrogram at a distance of 0.945, clustered the accessions into four different groups following their similarity and dissimilarity distances (Fig. 7). Cluster I was

comprised of all the seven commercial varieties and two landraces (TZA 2793 and TZA 5170), cluster II had one CIMMYT line (CML 442) and two landraces (TZA 3206 and TZA 5169), cluster III grouped the rest of 46 landraces while 10 CIMMYT lines were grouped into cluster IV.




Figure 7: Dendrogram for cluster analyses based on Euclidean genetic distance with average linkage of the 25 morphological characters generated from the performance of 68 maize accessions. Words in brackets show the source of seeds/collection site.

4.4 Discussion

4.4.1 Quantitative morphological traits

Maize is reported to be among the crops with high genetic diversity in terms of morphological as well as genetic variability (Hartings *et al.*, 2008). The maize accessions used in this study expressed a huge amount of variability in terms of quantitative characteristics. Landraces were found to be more variable than commercial varieties and the CIMMYT lines. Significant coefficient of variation among all evaluated traits in this study includes anthesis-silking interval and yield per plant. This was also reported by Sharma *et al.* (2010) in that significant genetic variation among maize landrace populations were found through yield related traits and flowering characteristics. The flowering behaviour might define the maturity differences among accessions (Olaoye, 2009) and they can also be connected to the yielding ability that early maturing accessions could generate high grain yield while those which are late maturing produce low yield (Lafitte *et al.*, 1997). A wide range of variation in flowering behaviours could signify the potential variability within accessions that would help on developing genotypes adaptable to different areas with different characteristics (Cömertpay, 2012). The existence of wide variability among the 68 accessions evaluated were further strengthened by box plot and Whisker, where landraces had more variability as compared with other groups despite their general low yield and other related parameters. Moreover, the principal component analysis identified quantitative morphological traits in different components that highly contributed to the total variation expressed by the accessions under study. The traits include 1000 kernel weight, plant height, ear height, yield per plant and days to 50% silking. That means, traits with high values in principal components present the potential characteristics for discriminating and identifying important accessions. The traits could also be used to characterize several maize landrace populations and discover potential candidates as parents for generating elite materials. Moreover, the principal component analysis expressed the distinction of the three groups of accessions used in this study with the traits contributing to their discriminating behaviours. Commercial varieties were discriminately identified by high yield, a thousand seed weight, number of rows per ear, ear diameter, ear length and early days to tasseling and silking. CIMMYT elite lines were characterized by significant low plant and ear height. Landraces were scattered along all

quadrants in a PCA plane, which signify them to be more diverse than the rest of the accessions involved in this study.

4.4.2 Qualitative morphological traits

The qualitative traits observed in this study explained distribution of accessions within each trait which differ among landraces, commercial varieties and CIMMYT lines. Only landraces had accessions distributed in each trait and not for commercial varieties and CIMMYT lines. The frequency distribution in percentage of accessions within traits differentiated the three groups in terms of foliage, tassel size, shape of upper surface of kernel and kernel type. Other traits like stem colour, sheath pubescence, tassel type, cob colour, shape of upper most ear, kernel row arrangement, kernel colour and endosperm colour characterized the three groups similarly though with different percent distribution. The former traits were able to discriminate between and within the three groups while the later identified differences just within each group. Traits that had higher percentage distribution of accessions towards one class within a trait include stem colour (green), sheath pubescence (intermediate), tassel type (primary-secondary), cob colour (white), shape of uppermost ear (conical), shape of kernel upper surface (rounded), kernel row arrangement (regular), kernel colour (white) and endosperm colour (white). The defined trait classes with high percentage accessions distribution might reflect farmers' preferences through successive selection (Ntundu *et al.*, 2006; Louette and Smale, 2000). In addition to the influence of farmers in shaping the structure of maize population, other factors such as species biology, geographical positioning, climatic settings, agricultural systems, biodiversity and local traditions have also an impact on population structuring (Prasanna, 2010; Pressoir and Berthaud, 2004).

4.4.3 Cluster analysis

The clustering indicated that the groups of accessions (landraces, commercial varieties and CIMMYT lines) were quite different from each other, though admixtures were observed. The grouping of the accessions mostly reflected individual performance and type of accessions. The commercial varieties and the two landraces in Cluster I were characteristically isolated due to distinct performance in high yield per plant, 1000 kernel weight, number of kernels per row, leaf length, plant height and few number of days to 50% tasseling and silking (Table 7). On the other

hand, landraces were more diverse in performance especially those that were grouped in cluster III. They had no specific unique behaviour except for ear height and the lengthy to anthesis-silking interval which explain their variability in terms of flowering time. CIMMYT elite lines in cluster IV had unique characteristics of low yield per plant, small plant sizes, mostly semi flint kernel type and many days to tasseling (Table 7). The mixed cluster II with one CIMMYT line and two landraces occurred due to the very unique characteristics that isolated them from the specific groups they were supposed to be. For example, CML 442 had a very high grain yield per plant as compared with the rest of CIMMYT lines. It also had intermediate sheath pubescence as compared with sparse pubescence that the rest of CIMMYT lines had (data not shown). On the other hand, TZA 5169 had the red kernel and cob colour, while TZA 3206 had yellow kernel colour and purplish cob colour different from the rest of the landraces (data not shown). The big cluster III of 46 landraces lack consistent originality grouping in terms of collection sites (Fig. 7). This observation was similar to the finding by Sun *et al.* (2016) who observed geographically close populations of Chinese sweetgum in different clusters. This implies that the landraces involved are comprised of a heterogeneous group that would have occurred through repeated exchange and selection of germplasm executed by farmers. The results of exchange and selection create the occurrence of irregular pattern of clustering (Ntundu *et al.*, 2006). Other reports also relate heterogeneity groupings with socio-economic factors, cultural, biological (open pollination) and migration of maize germplasm from one region to another (Hartings *et al.*, 2008; N'Da *et al.*, 2015; Cömertpay *et al.*, 2012). The findings of Ashimogo and Rukulantile (2000) explained that 35.4% of farmers in three regions studied in Tanzania use maize seeds they acquired from their neighbours and 60.1% grow their own farm-saved seeds. Furthermore, the clustering displayed some unique placement of accessions collected from Tanga to different clusters. Accession TZA 5170 from Tanga together with accession TZA 2793 from Morogoro were found in cluster I along with commercial varieties, also accession TZA 5169 from Tanga was grouped with other unique accession of TZA 3206 from Tabora region and CIMMYT line CML 442 in cluster II. In a mixed cluster III accessions TZA 2813, TZA 5162, TZA 5102 and TZA 5200 from Tanga were isolated and exerts higher distances (dissimilarities) with other accessions. This suggests a source of high variation is found in Tanga region.

4.5 Conclusion

This study revealed significant range of genetic diversity in the 68 maize accessions evaluated. This might provide a source of variation required for breeding programs to hold back the genetic vulnerability as a result of the recurrent outbreak of new strains of pest and diseases. It also offers an opportunity to widen the genetic background of the available maize germplasm because the materials that are currently at disposal for several breeding programs are composed of narrow genetic base. The traits that expressed high contribution towards total variability across the three groups of maize accessions used in this study include quantitative characteristics such as a 1000 kernel weight, plant height, ear height and yield per plant. The qualitative traits that had a significant contribution include foliage, tassel size, shape of upper surface of kernel and kernel type. All these traits might serve the purposes of generally discriminating among several populations through morphological characterization. However, each group was characterized specifically from other groups with specific traits and might as well signify the potentials expressed by each group. Cluster analysis identified the potential of landraces towards contributing a wealth of genetic resource for future breeding. Two landraces, TZA 2793 and TZA 5170, were grouped together with commercial varieties. The rest of the landraces possessed a wide range of variability in different traits that form a significant gene pool. That means the accessions might have strong contributions for producing superior varieties when used for introgressing promising traits. The cluster analysis also disclosed the expression of landraces lacking regular pattern in clustering within their major group. This elucidate the fact that farmers select cultivars based on their preferences and also exchange seed crop materials with fellow farmers even from very distant regions. Farmers play a significant role in shaping the structure of landrace population existing in a certain area. This calls for systematic involvement of farmers in breeding and selection process through participatory breeding in order to have an organized process of population structuring.

CHAPTER FIVE

Evaluating the response of maize accessions against maize lethal necrosis disease (MLND) in east Africa⁴

Mujuni Sospeter Kabululu^{1*}, Tileye Feyissa^{1,2}, Patrick Alois Ndakidemi¹

1. Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Arusha, Tanzania

2. Institute of Biotechnology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

* **For correspondence:** kabululum@nm-aist.ac.tz

Abstract

Maize lethal necrosis (MLN) is a viral disease that has severely affected maize crop and has put the east African community in jeopardy since maize is a crop that supports millions of people. The disease started in Kenya and spread to Tanzania, Uganda, South Sudan, Rwanda, DR Congo, Ethiopia and it continues to spread fast in other countries as well. It is caused by a combination of two viruses of *Maize chlorotic mottle virus* (MCMV) and any of the Potyvirus, but *Sugarcane mosaic virus* (SCMV) in particular for east Africa. Selection and production of genetically resistant varieties is the main solution of choice to this problem. Therefore, the objective of this study was to screen and select resistant maize genotypes. Field experiment with artificial inoculation was done at Naivasha in Kenya whereas natural field infestation was done at Mlangarini in Tanzania to screen for resistant local cultivars from Tanzania and to compare with elite lines from CIMMYT, Kenya. Data collected were subjected to Meta-R. Version 5.0, Genstat discovery edition 4 and STASTICA softwares for analyses. There were significant difference on mean disease occurrence and relative area under disease progress curve (rAUDPC) at Naivasha and no significant difference were found for Mlangarini. Accessions TZA2793 and TZA3544 had the lowest rAUDPC in both locations and they had both no significant difference with resistant check varieties. In addition, the cluster analysis grouped a landrace TZA2793 together with moderately resistant and resistant check varieties and hence considered promising for further improvements. Heritability at Naivasha was much higher (0.86) as compared to Mlangarini (0.18) and across locations (0.30). Significant negative correlation was found on

⁴ Submitted to Virus Disease Journal - Springer

MLND scores with sheath pubescence, kernel type and endosperm colour while significant positive correlations were found with anthesis silking interval, tassel size, ear damage and kernel type.

Key words: *Maize, Tanzania, disease resistance, landrace, rAUDPC, MLND, east Africa, virus*

5.1 Introduction

Maize lethal necrosis (MLN) is a viral disease that adversely affects maize crops in East Africa and other parts of the world (Mahuku *et al.*, 2015). In East Africa, it started in Kenya in 2011 as unknown disease and later was identified as *maize lethal necrosis disease* (MLND) after serological and molecular tests were carried out on infected maize plants from Bomet County and Nakuru County in 2012 (Wangai *et al.*, 2012b). In 2012, the disease was also reported to have spread in other areas of Kenya, neighbouring countries of Tanzania and Uganda and later to South Sudan, Rwanda, DR Congo, Ethiopia and it continues to spread fast into other countries. The emergence of MLND in Eastern Africa has added more challenges to the maize farming communities and other maize based-stakeholders (CGIAR, 2012). The control of this disease is difficult due to the fact that it is caused by a combination of more than one virus (Mahuku *et al.*, 2015; DSMZ, 2014). MLND occurs as a result of a positive interaction between *Maize chlorotic mottle virus* (MCMV) and any of the cereal viruses in the family, Potyviridae, such as *Sugarcane mosaic virus* (SCMV), *Maize dwarf mosaic virus* (MDMV), or *Wheat streak mosaic virus* (WSMV) (CIMMYT, 2013, Adams *et al.*, 2014). Specifically in East Africa, MLND is caused by MCMV and SCMV. These two viruses together inflict serious damage or even completely kill infected plants. For instance, *Maize chlorotic mottle virus* (MCMV) alone has been reported to be able to cause significant losses (Mahuku *et al.*, 2015). Farmers in the affected areas have been reported to experience extensive to total crop loss (Adams *et al.*, 2013; Wangai *et al.*, 2012b). The disease causing viruses are mainly transmitted from plant to plant by some insect vectors and they may also be carried by wind from field to field over long distances while spreading the disease (CGIAR, 2012). MCMV is normally transmitted by thrips and beetles while SCMV is transmitted by aphids (Kiruwa *et al.*, 2016). Infected plants show mild to severe mottling on the leaves, usually starting from the base of young leaves in the whorl and extending

upward towards the leaf tips. Also, stunting and premature aging of the plants, dying (necrosis) of the leaf margins that progresses to the mid-rib and eventually the entire leaf. In addition, necrosis of young leaves in the whorl before expansion has been reported leading to a symptom known as “dead heart” and eventually plant death (Wangai *et al.*, 2012b).

The control of the MLN disease relies on several management packages. However, generating varieties that are genetically resistant is important and economically feasible (Saleh *et al.*, 2016; Mahuku *et al.*, 2015; Wisser *et al.*, 2006). Thus, there is an urgent need to find germplasm with resistance traits and eventually establish genomic regions (markers) or map linked to MLN disease resistance. The identified candidates with those markers would then be crossed through breeding with MLND susceptible varieties carrying other desirable traits such as yield to obtain potential varieties against MLND (Semagn *et al.*, 2015; Ragimekula *et al.*, 2013; Ali and Yan, 2012). So far, there have been initiatives by several local and international institutions to evaluate as much number of germplasm against MLND as possible, but most of the materials ended up being susceptible. For instance, International Maize and Wheat Improvement Center (CIMMYT) and Kenya Agriculture and Livestock Research Organization (KALRO) have conducted joint studies from 2012 to 2014 with about 25,000 germplasm of elite inbred lines, Double Haploid lines, and hybrids against MLND under artificial inoculation and more than 95 percent of the tested materials were found to be completely susceptible (Gowda *et al.*, 2015). In this regard, Gowda *et al.* (2015) found 14 promising lines from 615 inbred lines that expressed MLND resistance and could be used as potential candidates for further improvements. Also, some few promising lines were identified with improved response to MCMV alone and they require further characterization to determine whether they have some level of resistance or tolerance to MLND (Mahuku *et al.*, 2015). Generally there is a believe that there are so far no maize materials which are completely resistant to the disease and hence further screening of more germplasm is justified (Mahuku *et al.*, 2015). Thus, a need arises to widen the source of germplasm with more diverse nature so as to increase the chance of finding more resistant sources against these 2 disease causing viruses and all kinds of strains associated with the viruses. High diversity in maize germplasm presents the opportunity to have a wide range of genetic variations that would enable it to produce sources of traits for survival against extreme

conditions like MLND. Maize landraces provide the most significant diverse sources of traits that are tolerant and / or resistant against pest and diseases (George *et al.*, 2004). Furthermore, maize genes within its genome are reported to be distant to an extent that any significant correlation in germplasm gives an indication of relationship between genetic variation and phenotypic trait variation (Wisser *et al.*, 2011). Therefore, phenotypic variation is an important aspect in plant breeding as it presents a significant component in MLND screening because it expresses the combined effects of all genes (Ragimekula *et al.*, 2013). In addition, Benson *et al.* (2015) pointed out that some genes conferring disease resistance are linked to phenotypic variations and other growth process. The aim of this study, therefore was to evaluate the phenotypic response of maize landraces, commercial varieties and CIMMYT lines against *maize lethal necrosis disease* (MLND).

5.2 Materials and methods

5.2.1 Natural MLND field infestation experiment (Mlangarini, Tanzania)

i. Materials and field layout

The experiment had 50 landraces, 7 commercial varieties from Tanzania and 11 inbred lines from CIMMYT Kenya (Table 10). A MLND hot spot site at Mlangarini located at S 03° 26' 12", E 036° 47' 13.4" and an elevation of 1128 meters above sea level was selected for establishing a natural field MLND infestation in Tanzania. The experiment was laid out in a randomized complete block design (RCBD) with three replications. Maize accessions were planted in rows of 4 meters long at spacing of 75 cm between rows and 30 cm within rows. The experiment was planted during rainy season and irrigated whenever it was necessary as well as other agronomic management practices to make sure that the plants faced no stress for a complete expression of their response to the disease. Two seeds were planted per hill followed by thinning to one plant two weeks after sowing. At planting, NPK (40:20:20) were applied at a rate of 100 kg N per hectare and top dressing was applied with Urea (46%) at a rate of 100 kg N per hectare.

Table 10: List of maize accessions and their sources as used in the screening experiments for resistance against maize lethal necrosis disease (MLND) in Kenya and Tanzania.

| NAIVASHA, KENYA | | | | | | MLANGARINI, TANZANIA | | | | | |
|-----------------|---------------------|-------------------|------|-----------------------|--------|----------------------|---------------------------|--------|------|----------|--------|
| SNo. | Acc name | Source | SNo. | Acc name ^c | Source | SNo. | Acc name | Source | SNo. | Acc name | Source |
| 1 | CML395 ^S | CML ^a | 35 | TZA4203 | GBTZ | 1 | CKDHL0500 ^{MR} | CML | 35 | TZA3167 | GBTZ |
| 2 | CML494 ^R | CML | 36 | TZA4320 | GBTZ | 2 | CKDHL120552 ^{MR} | CML | 36 | TZA3171 | GBTZ |
| 3 | TZA163 | GBTZ ^b | 37 | TZA4351 | GBTZ | 3 | CKSBL10205 ^{MR} | CML | 37 | TZA3181 | GBTZ |
| 4 | TZA1723 | GBTZ | 38 | TZA4574 | GBTZ | 4 | CLRCY034B ^R | CML | 38 | TZA3206 | GBTZ |
| 5 | TZA1724 | GBTZ | 39 | TZA4667 | GBTZ | 5 | CLRCY039 ^R | CML | 39 | TZA3310 | GBTZ |
| 6 | TZA1745 | GBTZ | 40 | TZA5102 | GBTZ | 6 | CLYN261 ^R | CML | 40 | TZA3536 | GBTZ |
| 7 | TZA1753 | GBTZ | 41 | TZA5129 | GBTZ | 7 | CML440 ^{MR} | CML | 41 | TZA3544 | GBTZ |
| 8 | TZA1757 | GBTZ | 42 | TZA5138 | GBTZ | 8 | CML442 ^S | CML | 42 | TZA3585 | GBTZ |
| 9 | TZA212 | GBTZ | 43 | TZA5162 | GBTZ | 9 | CML443 ^{MR} | CML | 43 | TZA3614 | GBTZ |
| 10 | TZA2263 | GBTZ | 44 | TZA5169 | GBTZ | 10 | CML544 ^{MR} | CML | 44 | TZA3837 | GBTZ |
| 11 | TZA2264 | GBTZ | 45 | TZA5170 | GBTZ | 11 | DEKALB | CMMV | 45 | TZA3914 | GBTZ |
| 12 | TZA2330 | GBTZ | 46 | TZA5200 | GBTZ | 12 | DH04 | CMMV | 46 | TZA3926 | GBTZ |
| 13 | TZA2338 | GBTZ | 47 | TZA5201 | GBTZ | 13 | PIONEER | CMMV | 47 | TZA3958 | GBTZ |
| 14 | TZA2731 | GBTZ | 48 | TZA5618 | GBTZ | 14 | SC403 | CMMV | 48 | TZA3971 | GBTZ |
| 15 | TZA2793 | GBTZ | 49 | TZA5619 | GBTZ | 15 | SITUKA M1 | CMMV | 49 | TZA4020 | GBTZ |
| 16 | TZA2813 | GBTZ | 50 | TZA599 | GBTZ | 16 | SITUKA1 | CMMV | 50 | TZA4164 | GBTZ |
| 17 | TZA2843 | GBTZ | 51 | TZA608 | GBTZ | 17 | TZMI730 ^{MR} | CML | 51 | TZA4203 | GBTZ |
| 18 | TZA2904 | GBTZ | 52 | TZA93 | GBTZ | 18 | TMV1 | CMMV | 52 | TZA4320 | GBTZ |
| 19 | TZA3167 | GBTZ | | | | 19 | TZA163 | GBTZ | 53 | TZA4351 | GBTZ |
| 20 | TZA3171 | GBTZ | | | | 20 | TZA1723 | GBTZ | 54 | TZA4574 | GBTZ |
| 21 | TZA3181 | GBTZ | | | | 21 | TZA1724 | GBTZ | 55 | TZA4667 | GBTZ |
| 22 | TZA3206 | GBTZ | | | | 22 | TZA1745 | GBTZ | 56 | TZA5102 | GBTZ |
| 23 | TZA3310 | GBTZ | | | | 23 | TZA1753 | GBTZ | 57 | TZA5129 | GBTZ |
| 24 | TZA3536 | GBTZ | | | | 24 | TZA1757 | GBTZ | 58 | TZA5138 | GBTZ |
| 25 | TZA3544 | GBTZ | | | | 25 | TZA212 | GBTZ | 59 | TZA5162 | GBTZ |
| 26 | TZA3585 | GBTZ | | | | 26 | TZA2263 | GBTZ | 60 | TZA5169 | GBTZ |
| 27 | TZA3614 | GBTZ | | | | 27 | TZA2264 | GBTZ | 61 | TZA5170 | GBTZ |
| 28 | TZA3837 | GBTZ | | | | 28 | TZA2330 | GBTZ | 62 | TZA5200 | GBTZ |
| 29 | TZA3914 | GBTZ | | | | 29 | TZA2338 | GBTZ | 63 | TZA5205 | GBTZ |
| 30 | TZA3926 | GBTZ | | | | 30 | TZA2731 | GBTZ | 64 | TZA5618 | GBTZ |
| 31 | TZA3958 | GBTZ | | | | 31 | TZA2793 | GBTZ | 65 | TZA5619 | GBTZ |
| 32 | TZA3971 | GBTZ | | | | 32 | TZA2813 | GBTZ | 66 | TZA599 | GBTZ |
| 33 | TZA4020 | GBTZ | | | | 33 | TZA2843 | GBTZ | 67 | TZA608 | GBTZ |
| 34 | TZA4164 | GBTZ | | | | 34 | TZA2904 | GBTZ | 68 | TZA93 | GBTZ |

^R stands for resistant check variety; ^{MR} stands for moderate resistant check variety; ^S stands for susceptible check variety; ^a CML stands for CIMMYT Lines; ^b GBTZ stands for Genebank Tanzania; ^c Acc name stands for Accession name.

ii. Data collection

Standardized screening protocol for MLND that was developed under the collaboration between USDA-ARS/Ohio State University, the Monsanto Company and KALRO was used to assess different resistance levels shown by the maize materials used in this study. The disease occurrence at each plot for every maize accession was scored using a 1 to 5 scale. The following is the description of each scoring scale;

1 = no MLN symptoms

2 = fine chlorotic streaks on lower leaves

3 = chlorotic mottling throughout plant

4 = excessive chlorotic mottling and dead heart

5 = complete plant necrosis

Along with MLND score data, the following characterization data were recorded; days to 50% tasseling, days to 50% silking, anthesis-silking interval, tassel size, foliage, plant height, ear height, leaf length, leaf width, number of leaves above uppermost ear, leaf pubescence, sheath pubescence, ear damage, ear length, ear diameter, cob colour, shape of uppermost ear, number of kernel rows, number of kernels per row, kernel length, kernel width, kernel thickness, kernel colour, endosperm colour, shape of upper surface of kernel, kernel row arrangement, 1000 kernel weight, kernel type, yield per plant.

5.2.2 Artificial inoculation experiment (Naivasha, Kenya)

i. Field layout

The experiment was laid out in an alpha lattice design with two replications. Two maize seeds per hill were sown in a 3 meters row and later thinned to one seed per hill. Naivasha is located at latitude 0° 43' S, longitude 36° 26' E, 1896 meters above sea level in Kenya. All other required agronomic managements were applied.

ii. Materials and artificial inoculation

The experiment involved 50 landraces from Tanzania and 2 inbred lines from CIMMYT Kenya (Table 10). The optimum viral ratio of combination between MCMV and SCMV was (1:4) for uniform MLND stress on plants. Inoculum production was prepared in two separate screen houses. Each viral inoculum was prepared by taking 1.0 g of infected leaf samples (viral presence confirmed by ELISA) and mixed with 20 ml of cold 0.1M Potassium phosphate buffer at pH 7.0. The mixture was homogenized using a blender and then sieved with cheese cloth to remove debris. Then 1.0 g carborundum was added to 1 litre of inoculum extract. The first plant inoculation in the field was done at 4 - 6 leaf stage using motorized mist blower (Solo 423). The second inoculation was done one week later after the first in order to make sure that maximum MLND pressure is inflicted on plants.

iii. Data collection

Data scoring on disease severity were taken using a 1 - 5 scale as described above. The MLND ratings on maize accession's response were recorded at 14, 28, 42 and 56 days after inoculation.

iv. Data analysis

The means estimated from the collected data of the response of maize accessions against MLND were predicted and adjusted towards the true population means using the statistical software Meta-R. Version 5.0. The analysis produced what is called best linear unbiased prediction (BLUP) for each accession and also coefficient of variations, genotypic variations, residual variations and heritability.

The response of maize accessions to MLND together with other morphological characters was related to each other through Pearson's correlation coefficient using STATISTICA 8.0 analytical software. Cluster analysis was performed based on group average method through Genstat discovery edition 4 to obtain clusters based on similarity and dissimilarity of accessions due to their response performance on MLND resistance. The disease severity was also measured using area under the disease progress curve (AUDPC) values which were obtained through an equation by Forbes *et al.* (2014) as follows;

$$\text{AUDPC} = \sum_{i=1}^{n-1} [(y_i + y_{i+1})/2] * [(t_{i+1} - t_i)]$$

where; Y_i is the percentage (or disease assessment score) of affected foliage at the i^{th} reading, t_i is the time (days) of each reading at the i^{th} observation and n is the total number of readings or observation. Moreover, "t" stands for days after planting, emergence or inoculation. The AUDPC produce an estimation of the amount of disease occurring on crop plants across the growing season. The relative area under the disease progress curve (rAUDPC) was calculated to take care of the interaction between genotypes and environments using the following formula;

$$\text{rAUDPC} = \text{Specific AUDPC} / ((\text{Days to the last reading} - \text{Days to the first reading}) * 100)$$

Both AUDPC and rAUDPC was calculated using Microsoft Excel spreadsheet program.

5.3 Results

5.3.1 Area under disease progress curve

The relative area under disease progress curve (rAUDPC) extended from 0.20 to 0.36 for Mlangarini experiment while at Naivasha rAUDPC was from 0.66 to 0.90 (Table 11). There was a significant ($p < 0.05$) variation with the disease progress curve at Naivasha while no significant difference was observed on plant response to MLND at Mlangarini. The average rAUDPC for Mlangarini was 0.25 and Naivasha was 0.76. Accessions DH04, TZA2263, CML442, DEKALB, TZA3544, TZA2904, TZA2793, SITUKA M1, and SITUKA1 expressed resistance at Mlangarini because they had the least rAUDPC and were not significantly different to the resistant check varieties (Table 11). On the other hand, accessions TZA3914, TZA4203, TZA3926, TZA1724, TZA3544, TZA5200, TZA2793, TZA2330, TZA2813, TZA3536, and TZA3167 expressed resistance at Naivasha as they had the least rAUDPC and were not significantly different from the resistant check variety (Table 11). Accessions CKDHL120552, SITUKA1, SITUKA M1, TZA2793 at Mlangarini and accession CML 494 at Naivasha had the least rAUDPC. The most susceptible accessions were TZMI730 and TZA2338 for Mlangarini closely followed by the susceptible check CML395 for Naivasha. Accessions TZA2793 and TZA3544 expressed the lowest rAUDPC at both locations.

Table 11: The selected 20 accessions at each location of experiment with the lowest relative area under disease progress curve (rAUDPC).

| MLANGARINI | | NAIVASHA | |
|---|---------------|-----------------------|----------------|
| Accession name | rAUDPC | Accession name | rAUDPC |
| Twelve accessions with the lowest rAUDPC | | | |
| CKDHL120552 | 0.20e | CML494 | 0.66j |
| SITUKA1 | 0.20e | TZA3167 | 0.68ij |
| SITUKA M1 | 0.20e | TZA3536 | 0.68ij |
| TZA2793 | 0.20e | TZA2813 | 0.68ij |
| TZA2904 | 0.21e | TZA2330 | 0.68hij |
| TZA3544 | 0.22de | TZA2793 | 0.68hij |
| DEKALB | 0.22de | TZA5200 | 0.68hij |
| CML442 | 0.22de | TZA3544 | 0.68hij |
| TZA2263 | 0.22de | TZA1724 | 0.69ghij |
| DH04 | 0.22de | TZA3926 | 0.69ghij |
| CLYN261 | 0.22cde | TZA4203 | 0.70fghij |
| CLRCY039 | 0.22cde | TZA3914 | 0.70fghij |
| Ten accessions with the highest rAUDPC | | | |
| TZMI730 | 0.36a | TZA2338 | 0.90a |
| TZA5200 | 0.33ab | CML395 | 0.88ab |
| TZA2813 | 0.32abc | TZA3614 | 0.87abc |
| TZA5138 | 0.31abcd | TZA2843 | 0.85abcd |
| TZA163 | 0.30abcde | TZA5102 | 0.85abcd |
| TZA3837 | 0.29abcde | TZA2904 | 0.85abcd |
| TZA2843 | 0.29abcde | TZA3181 | 0.84abcde |
| TZA599 | 0.29abcde | TZA5619 | 0.84abcde |
| TZA1723 | 0.29abcde | TZA2264 | 0.83abcdef |
| TZA2338 | 0.28abcde | TZA5138 | 0.83abcdefg |
| Grand mean | 0.25 | Grand mean | 0.76 |
| F-statistics | 1.29ns | F-statistics | 2.39 ** |
| CV (%) | 19.53 | CV (%) | 7.35 |
| LSD | 0.078 | LSD | 0.112 |

5.3.2 Means, variances and heritability

There was significant difference on maize accessions tested for MLND resistance at Naivasha, but no significant differences were observed at Mlangarini as well as across the two locations (Table 12). The mean MLND score at Naivasha was higher with a mean of 4.22 ranging from 2.31 to 4.89 while at Mlangarini the mean score was lower at 1.47 and across the two locations with 2.87. The genotypic variance recorded was high with 0.55 at Naivasha than at Mlangarini

and across environments. The residual variance was low (0.18) at Naivasha as compared with Mlangarini and across the locations. The heritability at Naivasha was much higher (0.86) as compared with Mlangarini (0.18) and across locations (0.30). The general coefficient of variation was higher at Mlangarini (29.91%) than at Naivasha and across the two environments.

Table 12: Response of maize accessions to MLND under artificial inoculation at Naivasha in Kenya and natural infestation at Mlangarini in Tanzania both on a 1–5 scale.

| Environment | Mean (range) | Genotypic variance | Residual variance | Heritability | LSD | CV (%) |
|---------------------|----------------------------------|--------------------|-------------------|--------------|------|--------|
| Naivasha | 4.22 (2.31 - 4.89)** | 0.55 | 0.18 | 0.86 | 0.85 | 10.03 |
| Mlangarini | 1.47 (1.39 - 1.71) ^{ns} | 0.02 | 0.19 | 0.18 | 0.88 | 29.91 |
| Across environments | 2.87 (2.47 - 3.07) ^{ns} | 0.07 | 0.19 | 0.30 | 0.61 | 10.65 |

5.3.3 Cluster analysis

The dendrogram which was generated from the 57 maize accessions analyzed based on group average is presented in Fig. 8. The analysis involved combining MLND scores collected from the two locations of Mlangarini in Tanzania and Naivasha in Kenya. The obtained dendrogram grouped the accessions into four different clusters based on their similarity and dissimilarity distances. Cluster I included two landraces, TZA 2813 and TZA 5200, cluster II grouped the 47 landraces, cluster III had just TZMI 730 and cluster IV contained 7 accessions of CKDHL0500, CKSBL10205, CKDHL120552, TZA2793, CLRCY039, CLYCN261, CLRCY034.

5.3.4 Correlation matrix

The Pearson correlation matrix presented the interrelationship between MLND scores and other morphological characters collected during this study (Table 13). There was a significant correlation between MLND scores and anthesis-silking interval, sheath pubescence, foliage, tassel size, ear damage, kernel type, endosperm colour, plant height, ear height, leaf length, ear length, ear diameter, number of kernels per row, kernel length, kernel width and 1000 kernel weight.




Figure 8: Dendrogram for cluster analyses based on Euclidean genetic distance with group average of the 57 accessions generated from their performance against MLND in two experiments at Naivasha, Kenya and Mlangarini, Tanzania.

Table 13: The Pearson correlation coefficient of 29 parameters in relation to the maize lethal necrosis diseases (MLND) scores obtained from a disease screening experiment of 68 maize accessions, probability significance at $p < 0.05$.

| 1 ^a | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | |
|----------------|--------|--------|--------|--------|--------|--------|--------|-------|--------|--------|--------|--------|--------|-------|--------|--------|-------|-------|-------|-------|-------|--------|--------|--------|--------|-------|--------|-------|------|------|
| 1 | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | 0.93* | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | 0.30* | 0.62* | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 4 | -0.18 | -0.18 | -0.09 | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | 0.07 | 0.16 | 0.28* | 0.14 | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | | |
| 6 | -0.11 | 0.00 | 0.25 | -0.07 | 0.13 | 1.00 | | | | | | | | | | | | | | | | | | | | | | | | |
| 7 | -0.08 | 0.12 | 0.49* | -0.04 | 0.24 | 0.65* | 1.00 | | | | | | | | | | | | | | | | | | | | | | | |
| 8 | -0.38* | -0.36* | -0.11 | 0.26 | 0.00 | 0.00 | 0.01 | 1.00 | | | | | | | | | | | | | | | | | | | | | | |
| 9 | 0.04 | 0.24 | 0.54* | -0.18 | 0.38* | 0.35* | 0.48* | -0.11 | 1.00 | | | | | | | | | | | | | | | | | | | | | |
| 10 | -0.32* | -0.37* | -0.28* | 0.01 | 0.00 | 0.16 | -0.01 | 0.04 | 0.03 | 1.00 | | | | | | | | | | | | | | | | | | | | |
| 11 | 0.14 | 0.24 | 0.32* | -0.03 | 0.2 | -0.26* | 0.02 | 0.13 | 0.05 | -0.16 | 1.00 | | | | | | | | | | | | | | | | | | | |
| 12 | 0.05 | 0.00 | -0.11 | 0.18 | 0.00 | -0.27* | -0.24 | -0.03 | -0.27* | -0.57* | 0.09 | 1.00 | | | | | | | | | | | | | | | | | | |
| 13 | 0.07 | -0.02 | -0.22 | 0.02 | -0.32* | -0.40* | -0.44* | -0.18 | -0.32* | 0.09 | 0.19 | 0.09 | 1.00 | | | | | | | | | | | | | | | | | |
| 14 | -0.15 | -0.15 | -0.06 | 0.13 | -0.12 | -0.03 | 0.01 | 0.70* | -0.09 | -0.15 | 0.09 | -0.03 | -0.07 | 1.00 | | | | | | | | | | | | | | | | |
| 15 | 0.03 | -0.09 | -0.30* | -0.08 | -0.35* | -0.09 | -0.29* | -0.06 | -0.12 | 0.07 | -0.13 | -0.04 | 0.36* | 0.19 | 1.00 | | | | | | | | | | | | | | | |
| 16 | 0.18 | 0.06 | -0.23 | -0.10 | -0.13 | -0.05 | -0.19 | -0.16 | -0.29* | 0.16 | -0.18 | -0.35* | 0.14 | -0.01 | -0.02 | 1.00 | | | | | | | | | | | | | | |
| 17 | 0.05 | 0.21 | 0.46* | -0.08 | 0.29* | 0.65* | 0.76* | -0.12 | 0.58* | 0.14 | -0.25 | -0.31* | -0.44* | -0.06 | -0.20 | -0.12 | 1.00 | | | | | | | | | | | | | |
| 18 | 0.25 | 0.41* | 0.56* | -0.11 | 0.39* | 0.61* | 0.71* | -0.12 | 0.58* | -0.03 | -0.13 | -0.19 | -0.49* | -0.07 | -0.28* | -0.18 | 0.92* | 1.00 | | | | | | | | | | | | |
| 19 | 0.13 | 0.18 | 0.19 | -0.24 | -0.06 | 0.58* | 0.62* | -0.08 | 0.28* | 0.07 | -0.36* | -0.21 | -0.35* | 0.05 | -0.03 | 0.10 | 0.75* | 0.68* | 1.00 | | | | | | | | | | | |
| 20 | -0.02 | -0.07 | -0.13 | 0.04 | -0.27* | 0.27* | 0.22 | 0.31* | 0.02 | 0.18 | -0.38* | -0.32* | -0.12 | 0.21 | 0.20 | 0.12 | 0.31* | 0.21 | 0.41* | 1.00 | | | | | | | | | | |
| 21 | -0.21 | -0.06 | 0.30* | -0.03 | 0.2 | 0.59* | 0.66* | 0.01 | 0.57* | 0.27* | -0.10 | -0.16 | -0.28* | 0.01 | -0.05 | -0.41* | 0.74* | 0.65* | 0.54* | 0.25 | 1.00 | | | | | | | | | |
| 22 | -0.30* | -0.24 | 0.04 | -0.05 | 0.24 | 0.49* | 0.53* | -0.06 | 0.38* | 0.24 | -0.32* | -0.04 | -0.48* | -0.21 | -0.26 | -0.07 | 0.57* | 0.46* | 0.50* | 0.27* | 0.59* | 1.00 | | | | | | | | |
| 23 | -0.55* | -0.47* | -0.05 | -0.01 | 0.13 | 0.59* | 0.49* | 0.16 | 0.30* | 0.39 | -0.30* | -0.10 | -0.36* | 0.04 | -0.16 | -0.16 | 0.51* | 0.38* | 0.39* | 0.24 | 0.66* | 0.67* | 1.00 | | | | | | | |
| 24 | -0.11 | -0.25 | -0.44* | -0.13 | -0.26 | -0.05 | -0.24 | -0.13 | -0.19 | 0.00 | -0.27* | 0.30* | 0.06 | -0.14 | 0.08 | 0.20 | -0.19 | -0.22 | 0.12 | 0.21 | -0.12 | 0.40* | 0.16 | 1.00 | | | | | | |
| 25 | -0.48* | -0.36* | 0.08 | 0.05 | 0.31* | 0.55* | 0.59* | 0.06 | 0.39* | 0.32 | -0.31* | -0.12 | -0.55* | -0.11 | -0.28* | -0.14 | 0.63* | 0.51* | 0.42* | 0.22 | 0.62* | 0.83* | 0.78* | 0.06 | 1.00 | | | | | |
| 26 | -0.13 | 0.06 | 0.45* | 0.01 | 0.38* | 0.40* | 0.61* | 0.00 | 0.49* | 0.25 | 0.03 | -0.32* | -0.34* | -0.07 | -0.20 | -0.32* | 0.67* | 0.61* | 0.35* | 0.08 | 0.68* | 0.52* | 0.32* | -0.42* | 0.58* | 1.00 | | | | |
| 27 | 0.50* | 0.57* | 0.43* | -0.04 | -0.05 | -0.19 | 0.02 | -0.16 | 0.19 | -0.31* | 0.23 | -0.07 | 0.14 | 0.03 | 0.13 | -0.18 | 0.08 | 0.16 | 0.06 | -0.03 | 0.08 | -0.31* | -0.63* | -0.42* | -0.42* | 0.30* | 1.00 | | | |
| 28 | -0.37* | -0.25 | 0.12 | 0.16 | 0.38* | 0.33* | 0.49* | 0.11 | 0.46* | 0.32* | -0.12 | -0.09 | -0.41* | -0.14 | -0.17 | -0.31* | 0.49* | 0.40* | 0.25 | 0.05 | 0.64* | 0.65* | 0.54* | -0.21 | 0.75* | 0.66* | -0.07 | 1.00 | | |
| 29 | -0.47* | -0.49* | -0.29* | 0.05 | -0.19 | 0.51* | 0.26 | 0.12 | 0.01 | 0.31* | -0.54* | -0.06 | -0.21 | 0.06 | 0.04 | 0.02 | 0.37* | 0.19 | 0.39* | 0.41* | 0.46* | 0.53* | 0.78* | 0.28* | 0.57* | 0.09 | -0.53* | 0.39* | 1.00 | |
| 30 | -0.15 | 0.07 | 0.50* | -0.28* | 0.20 | 0.39* | 0.58* | 0.19 | 0.45* | -0.04 | 0.20 | -0.25 | -0.54* | 0.08 | -0.33* | -0.19 | 0.36* | 0.41* | 0.29* | -0.01 | 0.30* | 0.31* | 0.35* | -0.17 | 0.36* | 0.42* | -0.08 | 0.29* | 0.02 | 1.00 |

^a Numbers in columns and rows heads are described below;

- | | | | |
|------------------------------|--|-------------------------------|------------------------|
| 1. Days to 50% tasseling | 9. Ear damage | 17. Plant height | 25. Kernel length |
| 2. Days to 50% silking | 10. Shape of uppermost ear | 18. Ear height | 26. Kernel width |
| 3. Anthesis-Silking Interval | 11. Shape of upper surface of kernel | 19. Leaf length | 27. Kernel thickness |
| 4. Sheath pubescence | 12. Kernel row arrangement | 20. Leaf width | 28. 1000 kernel weight |
| 5. Leaf pubescence | 13. Kernel type | 21. Ear length | 29. Yield per plant |
| 6. Foliage | 14. Kernel colour | 22. Ear diameter | 30. MLND score |
| 7. Tassel size | 15. Endosperm colour | 23. Number of kernels per row | |
| 8. Cob colour | 16. Number of leaves above uppermost ear | 24. Number of kernel row | |

Significant negative correlation was found with sheath pubescence (-0.28), kernel type (-0.54) and endosperm colour (-0.33). The significant correlations with high values included anthesis-silking interval (0.50), tassel size (0.58), ear damage (0.45) and kernel type (0.54).

5.4 Discussion

The standardized area under the disease progress curve (rAUDPC) was utilized to assess the cumulative disease progression within the growing process of the crops as explained by Luitel *et al.* (2016). The rAUDPC was higher at Naivasha (0.76) than at Mlangarini (0.26) because the extent of disease pressure at Naivasha was higher due to artificial inoculation. The inoculation procedure was done in such a way that maximum pathogen inoculum was applied to the plants. The inoculum application was carefully done twice at Naivasha while maize accessions were subjected to the natural disease pressure at Mlangarini with the extent of general occurring disease progression being low. The rAUDPC evaluated how the genetic differences among the accessions would determine the way they react to the extent of disease development on plants (Skelsey and Newton, 2014). The accessions with the lowest rAUDPC values were considered to be resistant to the MLN disease and vice versa (Safavi *et al.*, 2010). The tolerance performance is on the other hand determined by comparing the performance of accessions with known susceptible and resistant check materials (Massa *et al.*, 2015). Therefore, accessions that were able to show the least disease progression trend in both locations could be considered promising for resistance against MLND. In this case, accessions TZA2793 and TZA3544 may be considered for further screening to ascertain their resistance characteristics against MLND. The artificial inoculation of MLND pathogens at Naivasha realized the highest level of disease expression and the highest level of response by the accessions. The occurrence of the disease at Naivasha enhanced high heritability (0.86) as compared with the low heritability (0.18) at Mlangarini. This is due to the fact that uniform environmental conditions ensure high level of heritability while variable environments lower the level of heritability (Brunda *et al.*, 2014). The amount of inoculum applied on the plants at Naivasha was actually uniform such that genetic variation of the accessions in this study exerted an influence responsible for the variation in response against MLND. On the other hand, the Mlangarini situation was different because the amount of pathogens occurred naturally with no regular inoculum pressure distributed on plants.

Heritability measures phenotypic variance as caused by genetic variation, it provides prediction for plant breeding strategies. Heritability also determines the extent that a particular trait is transmitted to the next generation. It gives an opportunity for obtaining choices through selection process towards crop improvement (Bello *et al.*, 2012b).

The correlation between sheath pubescence and the occurrence of MLN disease was significantly negative. That means, an increase of pubescence is associated with lowering the disease (enhance resistance) to the crop plants. The resistance might directly be the close link of the pubescent marker and the gene controlling resistance or else the indirect link through the prevention of the insect pests carrying the pathogen causing the disease. Mmbaga *et al.* (1996) found the presence of dense pubescence to be linked to the resistance of rust on common bean and the pubescence trait was merely inherited. Endosperm colour is another character which was negatively correlated to the occurrence of MLN disease. The descriptor for endosperm colour used for scoring, ranged from white, cream, pale yellow, yellow, orange and white cap. A large descriptor number of endosperm colour indicated presence of more coloured endosperm such as yellow or orange, while the lower number were associated with whitish colour. Therefore, as the colour becomes more yellow or orange, it is associated with resistance and the white endosperm relate to susceptibility. Endosperm colour arises from the colour of the nutritious materials covering an embryo. Scott (1989) studied the link between endosperm colour and potyvirus resistance and found that the gene for yellow endosperm colour was linked to the resistance on potyviral disease. Also the promising inbred lines resistant to MLND that had been developed so far have yellow endosperm colour (Semagn *et al.*, 2015). Anthesis-silking interval (ASI) was significantly and positively correlated to the occurrence of the MLN disease, which means the longer the ASI the higher the chance for possible susceptibility, while short ASI indicated the possible resistance. Ngugi *et al.* (2013) established that ASI was significant as well as positively correlated with stress susceptible index. ASI is to a large extent determined by the variation in number of days to silking where a strong association of their Quantitative Trait Loci (QTLs) confirms the close link of the two traits (Gemenet *et al.*, 2010). This implies that short ASI depends on the short duration of the silking from the planting date (Magorokosho *et al.*, 2003) and a genotype with short days to silking would have a high chance of being able to tolerate or

resist growing stresses like diseases and drought. Another positive relationship was for MLND scores and tassel size, where large size of tassels associated with susceptibility and small tassels with resistance. Kernel type was also significantly correlated with the disease scores such that flint kernel type was found to associate with resistance traits. The indication of kernel type being correlated to disease resistance traits was also found by Marcon *et al.* (1996), but they found that dent kernel types had the relative resistance potential against high plains virus while sweet corn (sweet kernel type) expressed high susceptibility.

The cluster analysis grouped the accessions into groups depending on their performance against MLND in the two experiments that were set according to different ways of disease occurrence. That is, the disease established through artificial inoculation in the field (Naivasha, Kenya) with the highest pathogenic pressure and the natural disease occurrence in the field (Mlangarini, Tanzania) with low disease pressure. The clusters brought a landrace TZA2793 grouped together with the check varieties that are moderately resistant CKDHL0500, CKSBL10205, CKDHL120552 and resistant CLRCY039, CLYCN261, CLRCY034. The landrace could be considered for other tests to confirm its potential for MLND resistance. Cluster III had only TZMI 730 which was as well a check variety for moderate resistance but it isolated itself from the rest of the check varieties because it performed worse under natural infestation than even the landraces. The performance of TZMI 730 could have been caused by its adaptability and the exposure to a different environment which could not be verified for the rest of the CIMMYT lines that had the same source as itself. As expected, the rest of the 49 landrace accessions were grouped together in cluster II except for two accessions from Tanga regions TZA 2813 and TZA 5200 that had grouped themselves in cluster I. The two accessions isolated themselves from the other 47 accessions because they uniquely performed well at Naivasha and completely different at Mlangarini with worse performance. The way accessions distribute and group themselves on dendrogram reflects their performance (Lavanya *et al.*, 2008). However, some of the accessions may be named differently and yet they originated from the same geographic region (Biniam *et al.*, 2015). This also means that the current set of accessions especially landraces presents the existence of duplications from farmers who embrace a varied system of naming and managing their local materials (Tairo *et al.*, 2008).

5.5 Conclusion

The objective of this study was to evaluate the response of maize landraces from Tanzania against *maize lethal necrosis disease* (MLND) in comparison with commercial varieties and CIMMYT lines as checks. There was a significant variation in disease progression at Naivasha and no significant difference on disease progression was observed on plant response to MLND at Mlangarini. The artificial inoculation of MLND pathogens at Naivasha enhanced high heritability of 0.86 as compared to the low heritability (0.18) at Mlangarini. This was due to the fact that uniform disease pressure at Naivasha ensured high level of heritability while variable pathogenic distribution lowered the level of heritability at Mlangarini. The phenotypic variance (response against MLND) at Naivasha as caused by genetic variation was higher as compared with Mlangarini. The average relative area under disease progress curve (rAUDPC) was higher at Naivasha (0.76) than at Mlangarini (0.26) due to the extent of disease pressure at Naivasha being higher as compared with Mlangarini. The two locations presented the two different conditions that enabled evaluation of the response of accessions under investigation to the extreme situations. There is also a possibility that the two locations were infected with different strains of the virus causing MLND that is *Maize chlorotic mottle virus* (MCMV) and *Sugarcane mosaic virus* (SCMV) although not confirmed. Regarding, the assessment of accessions' performance on MLND progression over time, accessions TZA2793 and TZA3544 had the lowest rAUDPC in both locations and may be considered stable and promising for resistance characteristics against the disease. Phenotypic variations are important components to consider because they are linked to some genes related to disease resistance. Consequently, Pearson correlation matrix showed sheath pubescence, kernel type and endosperm colour to be negatively correlated with MLN disease scores. Also anthesis-silking interval, tassel size, ear damage and kernel type were positively correlated with MLND scores. Cluster analysis associated TZA2793 with moderate and resistant variety checks and grouped them in the same cluster.

CHAPTER SIX

Analyses of genetic diversity of maize landraces in Tanzania using SSR markers for maize lethal necrosis disease resistance⁵

Mujuni Sospeter Kabululu^{1*}, Tileye Feyissa^{1,2}, Pavithravani Enkataramana¹, Patrick Alois Ndakidemi¹

¹Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Arusha, Tanzania

² Institute of Biotechnology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

* **For correspondence:** kabululum@nm-aist.ac.tz

Abstract

Maize is a staple crop as well as cash crop which is very important for millions of people in sub-Saharan Africa and other parts of the world. Its production has been adversely confounded by several diseases that cause significant yield loss. A new disease known as *maize lethal necrosis* (MLN) has recently hit East Africa and added even more challenges in maize production. Breeding for genetically resistant varieties is the most preferred and reliable solution against MLN disease. Genetic diversity provides key information that aid in identifying important traits against maize production constraints. The aim of this study was to evaluate the genetic diversity of maize landraces from Tanzania and other accessions using SSR markers for resistance against *maize lethal necrosis disease*. A total of 63 alleles with an average of 3.15 per locus were detected using 20 simple sequence repeats (SSR) loci distributed in 9 chromosomes of maize. The polymorphism information content (PIC) ranged from 0.17 to 0.79, with an average of 0.52. Analysis of molecular variance showed that more than half of the total variation (71%) was accounted for the variation within accessions and the rest (29%) was divided to among accessions (17%) and among populations (12%). Cluster and principal coordinate analysis clearly clustered and isolated CIMMYT inbred lines from landraces and commercial varieties because they were obtained from the same source with similar breeding programs different from the other groups. The distribution and frequencies of alleles in SSR markers phi029 on

⁵ Submitted to Indian Journal of Biotechnology

chromosomal bin location 3.04 and phi062 on bin location 10.04 previously identified to be associated with quantitative trait loci (QTL) for *Sugarcane mosaic virus* (SCMV) resistance were calculated and found more with landraces. Thus, the obtained genetic details associated with resistance against SCMV provided an insight towards establishing the utility function of the occurring genetic diversity within studied accessions for MLND resistance.

Key words: *Maize, landraces, SCMV, MCMV, MLND, SSR markers, genetic diversity, Tanzania*

6.1 Introduction

Maize (*Zea mays* L.) which was domesticated about 9000 years ago in Mexico (Warburton *et al.*, 2008) belongs to the Poaceae family (Abdellatif and Khidr, 2010). It is the third most important cereal crop in the world after rice and wheat (Al-Badeiry *et al.*, 2014, Legesse *et al.*, 2007). In Tanzania, maize is the major food as well as cash crop where its supply is normally equated to the national food security (Kabululu *et al.*, 2017; Katinila *et al.*, 1998). However, the production of maize has been challenged by a number of diseases which cause serious grain yield loss (Anjichi, 2005; Pechanova and Pechan, 2015). In 2011, a devastating disease called *maize lethal necrosis disease* (MLND) emerged in East Africa through Kenya (Wangai *et al.*, 2012b). The disease was found to be caused by synergistic interaction between *Maize chlorotic mottle virus* (MCMV) and *Sugarcane mosaic virus* (SCMV). The two viruses together inflict serious damage to an extent that farmers in the affected areas have been reported to experience extensive to total crop loss (Mahuku *et al.*, 2015; Adams *et al.*, 2014; Wangai *et al.*, 2012b). The control of the disease depends on several methods, while generating varieties that are genetically resistant being important and economically feasible (Saleh *et al.*, 2016; Mahuku *et al.*, 2015; Wisser *et al.*, 2006). Thus, there is a strong need to discover germplasm with resistance traits and eventually establishing genomic regions (markers) map linked to MLN disease resistance. The identified candidates with those markers would then be crossed through breeding with MLND susceptible varieties carrying other desirable traits such as yield to obtain potential varieties against MLND (Semagn *et al.*, 2015; Ragimekula *et al.*, 2013; Ali and Yan, 2012).

Genetic diversity in maize germplasm helps to have reliable information to facilitate breeding programs for genetic improvement and reduction of genetic vulnerability to pests and diseases (Lee, 1998; Abdellatif and Khidr, 2010). It is the basis for survival and adaptation for plants (Rao and Hodgkin, 2002) which provides adequate variation that underscores the possibility for further manipulation to attain a certain goal (Rahman *et al.*, 2008). Different germplasm such as landraces, cultivated and new elite materials have always been screened to look for the plant materials that would express desirable traits such as yield and resistance to diseases (ESA, 2010). Furthermore, genetic diversity at molecular level using markers is critical as a resource to find new alleles for important traits through potential sources of resistances and tolerances to biotic and abiotic stresses (Taba *et al.*, 2004; Rauf *et al.*, 2010). It can be categorized in terms of the number of different alleles existing in different populations, distribution of those alleles in the chromosomes, the impact they have on performance and the general variability among different populations under various environmental conditions (Rao and Hodgkin, 2002; Mondini *et al.*, 2009; Bindroo and Moorthy, 2014). A molecular marker is a variant of DNA or a protein which can be detected and whose inheritance can be monitored reliably (Jones *et al.*, 1997). Polymerase chain reaction (PCR) based molecular marker techniques have made it possible for breeders and other scientists to make genetic diversity estimates as generated through different molecular markers (Arif *et al.*, 2010; Poczai *et al.*, 2013). Some of the PCR based molecular techniques that have been applied in molecular studies include RAPDs (Brown-Guedira *et al.*, 2000; Mondini *et al.*, 2009), AFLPs (Mondini *et al.*, 2009), Simple Sequence Repeats (SSRs) (Kumari *et al.*, 2005; Beyene *et al.*, 2005; Mondini *et al.*, 2009; Aci *et al.*, 2013; Abraha *et al.*, 2014). However, these different marker techniques emphasize on different features (Abdel-Mawgood, 2012) and different aspects of genetic diversity (Matsuoka *et al.*, 2002; Mondini *et al.*, 2009). SSRs have been the marker system of choice for population genetic studies, because it combines many desirable properties including co-dominance, robustness, hypervariability, rapid and simple assays, and it is uniformly dispersed in plant genomes (Powell *et al.*, 1996; Beyene *et al.*, 2005; Prasanna and Hoisington, 2003).

George *et al.* (2004) screened 102 inbred lines using 76 SSR markers for downy mildew resistance and the frequencies of alleles in SSR loci linked to previously identified quantitative

trait loci (QTL) were calculated. The association of alleles with disease and pest resistance identified five inbred lines through allelic patterns (George *et al.*, 2004). Resistance to MLND requires a good understanding of the genetic structure of the viruses that cause the disease so as to enhance efficient generation of resistant germplasm. Apparently, the genetics of resistance to *Sugarcane mosaic virus* (SCMV) as one of the two MLND causing viruses has been investigated very well, where the major quantitative trait loci (QTL) against SCMV has repeatedly been detected on the short arm of chromosome 3 and on the long arm of chromosome 10 (Mahuku *et al.*, 2015; Zhang *et al.*, 2003). The markers linked to the SCMV resistance QTLs can be used in the marker assisted selection to attain germplasm improvement towards obtaining solution against MLND. Utilization of the underlying variation emphasizes the use of landraces and wild relatives of crop plants because they carry desirable genes that grant resistance to pests and diseases (Beyene *et al.*, 2005). However, landraces and wild relatives present a group of germplasm which are less exploited with limited agronomic and genetic data that renders difficulties to use them (Hoisington *et al.*, 1999; Warburton *et al.*, 2002; Molin *et al.*, 2013). The objective of this study was to evaluate the genetic diversity of maize landraces from Tanzania and other accessions through the use of SSR markers as relates to the resistance against *maize lethal necrosis disease*.

6.2 Materials and methods

6.2.1 Plant materials

A total of 96 accessions were used for this study comprising of 51 landraces from Tanzania, 11 elite lines from CIMMYT, Kenya and 34 commercial varieties in Tanzania (Table 14). The landraces from Tanzania were collected from different regions representing all agroecological zones and were kept at the national plant genetic resource center (NPGRC) in Arusha. The elite lines from CIMMYT, Kenya were included in this study as checks for resistance, moderately resistance and susceptibility to MLND (Table 14). In order to enhance genetic variability, commercial varieties were also included as they have potential for other agronomic traits.

6.2.2 Extraction of DNA

Fifteen seeds were planted in greenhouse for each accession and at 3 - 4 leaf stage, an approximate 0.75 g healthy young leaf tissue was taken from 15 bulked plants. The total genomic DNA from the leaf samples was extracted using CTAB procedure by Dellaporta *et al.* (1983) with slight modification. DNA pellets were washed once with 70% ethanol and then air dried for 1 hour before dissolving them in 100 µl of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). The concentration of DNA samples was determined through 0.8% agarose in 0.5xTBE buffer (Tris base + Boric acid + 0.5M EDTA), containing a gel staining solution of EZ – Vision (Amresco, fountain parkway solon, OH USA).

6.2.3 Simple Sequence Repeats markers

Thirty SSR primers shown in Table 15 were selected as of the previous study (Warburton *et al.*, 2002) and from the public Maize GDB (http://www.agron.missouri.edu/ssr_probes/ssr.htm) based on their good polymorphism information content (data not shown) and coverage of 10 chromosomes.

6.2.4 Polymerase Chain Reaction and gel electrophoresis

The reactions were carried out in a thermal cycler with 10 µl reaction mixture consisting of 20 ng template DNA with a volume of 2µl, then 0.5 µl of 1 mM dNTPs, 2 µl of 2.5 mM SSR primers (forward and reverse), 0.05 µl of 5 U/ µl Taq polymerase, 2.5 µl of 10x Taq buffer and 0.95 µl of autoclaved distilled water. PCR reactions were performed in a BIO-RAD thermocycler with the following conditions: an initial denaturation at 95°C for 5 min, followed by 35 cycles of 1min at 95°C denaturation, 1min annealing (52 to 60°C depending on the requirement of specific primer), and 1min extension at 72°C and the final extension step of 72°C for 5 min was performed. The amplified products were separated by electrophoresis in 2.5% agarose in 0.5xTBE buffer (Tris base + Boric acid + 0.5M EDTA), containing a gel staining solution of EZ – Vision (Amresco, fountain parkway solon, OH USA). The gels were photographed under UV light in BIO-RAD Gel Doc EZ Imager and the images were transferred to a computer for scoring and documentation. A 100 base pairs DNA ladder was used as standard molecular weight.

Table 14: List of maize accessions and their sources as used in the evaluation of genetic diversity for maize lethal necrosis disease resistance.

| SNo. | Accession name | Status | Collection source | Population/ zone | SNo. | Accession name | Status | Collection source | Population/ zone |
|------|---------------------------|------------|-------------------|---------------------|------|----------------|----------|-------------------|---------------------|
| 1 | TZA597 | Landrace | Singida | Central zone | 49 | TZA2731 | Landrace | Morogoro | Eastern zone |
| 2 | TZA599 | Landrace | Singida | Central zone | 50 | TZA2793 | Landrace | Morogoro | Eastern zone |
| 3 | TZA608 | Landrace | Singida | Central zone | 51 | TZA2813 | Landrace | Tanga | Eastern zone |
| 4 | CKDHL0500 ^{MR} | Elite line | CIMMYT | CIMMYT | 52 | TZA2843 | Landrace | Tanga | Eastern zone |
| 5 | CKDHL120552 ^{MR} | Elite line | CIMMYT | CIMMYT | 53 | TZA3536 | Landrace | Morogoro | Eastern zone |
| 6 | CKSBL10205 ^{MR} | Elite line | CIMMYT | CIMMYT | 54 | TZA3544 | Landrace | Morogoro | Eastern zone |
| 7 | CLRCY034 ^R | Elite line | CIMMYT | CIMMYT | 55 | TZA5101 | Landrace | Tanga | Eastern zone |
| 8 | CLRCY039 ^R | Elite line | CIMMYT | CIMMYT | 56 | TZA5102 | Landrace | Tanga | Eastern zone |
| 9 | CLYN261 ^R | Elite line | CIMMYT | CIMMYT | 57 | TZA5129 | Landrace | Tanga | Eastern zone |
| 10 | CML440 ^{MR} | Elite line | CIMMYT | CIMMYT | 58 | TZA5138 | Landrace | Tanga | Eastern zone |
| 11 | CML442 ^S | Elite line | CIMMYT | CIMMYT | 59 | TZA5162 | Landrace | Tanga | Eastern zone |
| 12 | CML443 ^{MR} | Elite line | CIMMYT | CIMMYT | 60 | TZA5169 | Landrace | Tanga | Eastern zone |
| 13 | CML544 ^{MR} | Elite line | CIMMYT | CIMMYT | 61 | TZA5170 | Landrace | Tanga | Eastern zone |
| 14 | TZMI730 | Elite line | CIMMYT | CIMMYT | 62 | TZA5186 | Landrace | Tanga | Eastern zone |
| 15 | CML445 | Comm Var | Comm Var | Comm Var | 63 | TZA5200 | Landrace | Tanga | Eastern zone |
| 16 | CML444 | Comm Var | Comm Var | Comm Var | 64 | TZA3914 | Landrace | Mara | Lake zone |
| 17 | CML489 | Comm Var | Comm Var | Comm Var | 65 | TZA3926 | Landrace | Mara | Lake zone |
| 18 | DH04 | Comm Var | Comm Var | Comm Var | 66 | TZA3958 | Landrace | Mara | Lake zone |
| 19 | DK8031 | Comm Var | Comm Var | Comm Var | 67 | TZA3971 | Landrace | Mara | Lake zone |
| 20 | Phb3253 | Comm Var | Comm Var | Comm Var | 68 | TZA4020 | Landrace | Mwanza | Lake zone |
| 21 | SC403 | Comm Var | Comm Var | Comm Var | 69 | TZA4203 | Landrace | Mwanza | Lake zone |
| 22 | SITUKA 1 | Comm Var | Comm Var | Comm Var | 70 | TZA4320 | Landrace | Kagera | Lake zone |
| 23 | SITUKA M1 | Comm Var | Comm Var | Comm Var | 71 | TZA4351 | Landrace | Kagera | Lake zone |
| 24 | TMV-1 | Comm Var | Comm Var | Comm Var | 72 | TZA4574 | Landrace | Mwanza | Lake zone |
| 25 | DKC9089518 | Comm Var | Comm Var | Comm Var | 73 | TZA4667 | Landrace | Mwanza | Lake zone |
| 26 | H519 | Comm Var | Comm Var | Comm Var | 74 | TZA5618 | Landrace | Manyara | Nothern zone |
| 27 | H614D | Comm Var | Comm Var | Comm Var | 75 | TZA5619 | Landrace | Manyara | Nothern zone |
| 28 | KH600-95A | Comm Var | Comm Var | Comm Var | 76 | TZA5620 | Landrace | Manyara | Nothern zone |
| 29 | P2859W | Comm Var | Comm Var | Comm Var | 77 | TZA1723 | Landrace | Njombe | Southern Highland |
| 30 | SC513 | Comm Var | Comm Var | Comm Var | 78 | TZA1724 | Landrace | Njombe | Southern Highland |
| 31 | STAHA | Comm Var | Comm Var | Comm Var | 79 | TZA1745 | Landrace | Njombe | Southern Highland |
| 32 | TZH536 | Comm Var | Comm Var | Comm Var | 80 | TZA1757 | Landrace | Mbeya | Southern Highland |
| 33 | DK8053 | Comm Var | Comm Var | Comm Var | 81 | TZA212 | Landrace | Mbeya | Southern Highland |
| 34 | TZM523 | Comm Var | Comm Var | Comm Var | 82 | TZA2910 | Landrace | Ruvuma | Southern Highland |
| 35 | ZAMS606 | Comm Var | Comm Var | Comm Var | 83 | TZA93 | Landrace | Rukwa | Southern Highland |
| 36 | H628 | Comm Var | Comm Var | Comm Var | 84 | TZA163 | Landrace | Mtwara | Southern zone |
| 37 | KILIMA | Comm Var | Comm Var | Comm Var | 85 | TZA2263 | Landrace | Lindi | Southern zone |
| 38 | PAN15 | Comm Var | Comm Var | Comm Var | 86 | TZA2264 | Landrace | Lindi | Southern zone |
| 39 | PAN3M-01 | Comm Var | Comm Var | Comm Var | 87 | TZA2330 | Landrace | Lindi | Southern zone |
| 40 | PAN4M-19 | Comm Var | Comm Var | Comm Var | 88 | TZA2338 | Landrace | Mtwara | Southern zone |
| 41 | PAN4M-21 | Comm Var | Comm Var | Comm Var | 89 | TZA3585 | Landrace | Mtwara | Southern zone |
| 42 | Phb30G19 | Comm Var | Comm Var | Comm Var | 90 | TZA3614 | Landrace | Mtwara | Southern zone |
| 43 | SC627 | Comm Var | Comm Var | Comm Var | 91 | TZA3837 | Landrace | Mtwara | Southern zone |
| 44 | LUBANGO | Comm Var | Comm Var | Comm Var | 92 | TZA3167 | Landrace | Kigoma | Western zone |
| 45 | TZH538 | Comm Var | Comm Var | Comm Var | 93 | TZA3171 | Landrace | Kigoma | Western zone |
| 46 | DK9089 | Comm Var | Comm Var | Comm Var | 94 | TZA3181 | Landrace | Kigoma | Western zone |
| 47 | CZL0616/CZL097 | Comm Var | Comm Var | Comm Var | 95 | TZA3206 | Landrace | Tabora | Western zone |
| 48 | H625 | Comm Var | Comm Var | Comm Var | 96 | TZA3310 | Landrace | Tabora | Western zone |

^R stands for resistant check variety; ^{MR} stands for moderate resistant check variety; ^S stands for susceptible check variety; ^a **Comm Var** stands for Commercial variety; ^b **SNo.** stands for Serial number

Table 15: The 30 primers used in this study with their names, repeat units, bin location and sequence details.

| Serial No. | Primer | | Bin ^b | Sequence ^c |
|------------|-----------|---------------------|------------------|---|
| | name | Repeat ^a | | |
| 1 | phi011 | Tri | 1.09 | For ^e : TGTTGCTCGGTCACCATAACC Rev ^f : GCACACACACAGGACGACAGT |
| 2 | phi015 | Tetra | 8.09 | For : GCAACGTACCGTACCTTTCCGA Rev : ACGCTGCATTCAATTACCGGAAG |
| 3 | phi029 | Comp. ^d | 3.04 | For : TTGTCTTTCTTCTCCACAAGCAGCGAA Rev : ATTTCCAGTTGCCACCGACGAAGAAGT |
| 4 | phi031 | Tetra | 6.04 | For : GCAACAGGTTACATGAGCTGACGA Rev : CCAGCGTGCTGTTCCAGTAGTT |
| 5 | phi062 | Tri | 10.04 | For : CCAACCCGCTAGGCTACTTCAA Rev : ATGCCATGCGTTCGCTCTGTATC |
| 6 | phi065 | Penta | 9.03 | For : AGGGACAAATACGTGGAGACACAG Rev : CGATCTGCACAAAGTGGAGTAGTC |
| 7 | phi072 | Tetra | 4.01 | For : ACCGTGCATGATTAATTTCTCCAGCCTT Rev : GACAGCGCGCAAATGGATTGAAGT |
| 8 | phi083 | Tetra | 2.04 | For : CAAACATCAGCCAGAGACAAGGAC Rev : ATTCATCGACGCGTCCAGTCTACT |
| 9 | phi100175 | Tetra | 8.06 | For : TATCTGACGAATCCCATTCCC Rev : TACGTAACGGACGGACGG |
| 10 | phi102228 | Tetra | 3.04 | For : ATTCGACGCAATCAACA Rev : TTCATCTCTCCAGGAGCCTT |
| 11 | phi109642 | Tetra | 2.00 | For : CTCTCTTTCCTTCCGACTTTCC Rev : GAGCGAGCGAGAGAGATCG |
| 12 | phi233376 | Tri | 8.03 | For : CCGGCAGTCGATTACTCC Rev : CGAGACCAAGAGAACCCTCA |
| 13 | phi229852 | Tri | 6.08 | For : GATGTGGGTGCTACGAGCC Rev : AGATCTCGGAGCTCGGCTA |
| 14 | phi420701 | Tri | 8.01 | For : GATGTTTCAAACCACCCAGA Rev : ATGGCACGAATAGCAACAGG |
| 15 | phi453121 | Tri | 3.00 | For : ACCTTGCTGTCTTCTTTCT Rev : CAAGCAAGACTTTTGATCAGCC |
| 16 | phi96342 | Tetra | 10.02 | For : GTAATCCCACGTCCTATCAGCC Rev : TCCAACTTGAACGAACTCCTC |
| 17 | umc1109 | Tri | 4.10 | For : GCAACACAGGACCAATCATCTCT Rev : GTTCGGTCCGTAGAAGAAGTCTCA |
| 18 | umc1122 | Tri | 1.06 | For : CACAACTCCATCAGAGGACAGAGA Rev : CTGCTACGACATACGCAAGGC |
| 19 | umc1136 | Tri | 3.10 | For : CTGCATACAGACATCCAACCAAAG Rev : CTCTCGTCTCATCACCTTTCCCT |
| 20 | umc1143 | Penta | 6.00 | For : CGTGGTGGGATGCTATCCTTT Rev : GACACTAGCAATGTTCAAACCCC |
| 21 | umc1152 | Tetra | 10.01 | For : CCGAAGATAACCAAAACAATAATAGTAGG Rev : ACTGTACGCCTCCCCTTCTC |
| 22 | umc1153 | Tri | 5.09 | For : CAGCATCTATAGCTTGCTTGCAAT Rev : TGGGTTTTGTTGTTGTTGTTGTTG |
| 23 | umc1279 | Tri | 9.00 | For : CAATCCAATCCGTTGCAGGTC Rev : GATGAGCTTGACGACGCTG |
| 24 | umc1304 | Tetra | 8.02 | For : GCCAACTAGAACTACTGCTGCTCC Rev : CATGCAGCTCTCCAAATTAATCC |
| 25 | umc1545 | Tetra | 7.00 | For : GAAAATGCATCAACAACAAGCTG Rev : ATTGGTTGGTTCTTGCTTCCATTA |
| 26 | phi006 | Tri | 4.11 | For : AGGCGGCGTGCTGAACACCT Rev : CGCTTCATCTCCCGTGACAATG |
| 27 | phi034 | Tri | 7.02 | For : TAGCGACAGGATGGCCTTCT Rev : GGGGAGCACGCCTTCGTTCT |
| 28 | phi063 | Tetra | 10.02 | For : GGCGGCGGTGCTGGTAG Rev : CAGCTAGCCGTAGATATACGCT |
| 29 | phi064 | Tetra | 1.11 | For : CCGAATTGAAATAGCTGCGAGAACCT Rev : ACAATGAACGGTGGTTATCAACACGC |
| 30 | phi227562 | Tri | 1.12 | For : TGATAAAGCTCAGCCACAAGG Rev : ATCTCGGCTACGGCCAGA |

^a **Repeat** indicates the repeat units of a primer

^b **Bin** indicates chromosomal location

^c **Sequence**, the first sequence represents the Forward and the second Reverse

^d **Comp.** represents a compound repeat, containing more than one repeat type

^e **For** stands for Forward and ^f **Rev** stands for Reverse

6.2.5 Data analysis

Band profiles were binary coded 1 for presence or 0 for absence within each locus. The discriminatory power of each locus generated by all accessions, were established by the polymorphism information content (PIC) given by the following formula;

$$PIC = 1 - \sum f_i^2$$

where f_i is the frequency of the i th allele (Smith *et al.* 1997). Nei's genetic distances between groups of population, Nei's gene diversity and Shannon's information index were analysed using POPGENE software version 1.31 (Nei and Li 1979). The analysis of the molecular variance (AMOVA) among the populations, among accession and within accessions were done by GenAlEx 6.5 Excel package software. Cluster analysis was performed on the genetic dissimilarity matrix using unweighted pair group arithmetic average method (UPGMA). The dendrogram and principal coordinate analysis were generated using the PAST statistical software.

6.3 Results

6.3.1 Simple Sequence Repeats polymorphism

Among the thirty primers utilized in this study, only twenty were selected for further analysis (Table 16) while the other ten that did not amplify or produced monomorphic bands were excluded. The allelic diversity measured at each SSR locus is presented in Table 16. A total of 63 alleles with an average of 3.15 per locus were detected using 20 simple sequence repeat (SSR) loci distributed on 9 chromosomes of maize. The number of alleles ranged from five (phi109642 and phi065) to two (phi011, umc1122, phi031, phi420701, phi062) . The polymorphism information content (PIC) ranged from 0.17 (phi031) to 0.79 (phi065), with an average of 0.52. Eleven primers had the PIC greater than the overall average of 0.52 while the other 9 primers had the PIC less than the average (Table 16).

Table 16: The selected 20 primers with their number of alleles, bin location, repeat units and polymorphism information content (PIC).

| Primer | Repeat ^a | Bin ^b | Total number of alleles | PIC ^c |
|----------------------|---------------------|------------------|-------------------------|------------------|
| phi011 | Tri | 1.09 | 2 | 0.50 |
| umc1122 | Tri | 1.06 | 2 | 0.49 |
| phi083 | Tetra | 2.04 | 3 | 0.56 |
| phi109642 | Tetra | 2.00 | 5 | 0.63 |
| phi029 | Comp. ^d | 3.04 | 3 | 0.26 |
| phi453121 | Tri | 3.00 | 4 | 0.72 |
| umc1136 | Tri | 3.10 | 3 | 0.50 |
| phi006 | Tri | 4.11 | 3 | 0.56 |
| umc1153 | Tri | 5.09 | 3 | 0.53 |
| phi031 | Tetra | 6.04 | 2 | 0.17 |
| umc1143 | Penta | 6.00 | 3 | 0.54 |
| phi015 | Tetra | 8.09 | 3 | 0.66 |
| phi233376 | Tri | 8.03 | 3 | 0.42 |
| phi420701 | Tri | 8.01 | 2 | 0.32 |
| umc1304 | Tetra | 8.02 | 4 | 0.65 |
| phi065 | Penta | 9.03 | 5 | 0.79 |
| umc1279 | Tri | 9.00 | 3 | 0.61 |
| phi062 | Tri | 10.04 | 2 | 0.44 |
| phi96342 | Tetra | 10.02 | 3 | 0.54 |
| umc1152 | Tetra | 10.01 | 3 | 0.49 |
| Average/Total | | | 63 | 0.52 |

^a **Repeat** indicates the repeat unit of a primer

^b **Bin** indicates chromosomal location

^c **PIC** stands for Polymorphism Information Content

^d **Comp.** represents a compound repeat, containing more than one repeat type

6.3.2 Analysis of molecular variance (AMOVA)

The summary of AMOVA as given in Table 17 was done to estimate the population genetic variation among and within the groups of accessions used in this study. The source of variation included among populations of accessions, among accessions and within accessions. There were highly significant differences ($p < 0.05$) of molecular variance within and among the accessions used in this study (Table 17). The 71% of the total genetic variation was significantly ($p = 0.001$) observed within accessions while 17% of the total variance was found among accessions and 12% was generated among populations at $p = 0.03$ and 0.005 respectively (Table 17 and Fig. 9).

Table 17: Analysis of molecular variance (AMOVA) for the 96 accessions used in this study.

| Source | Degrees of freedom | Sum of Squares | Mean Square | Estimated Variance | Molecular variance (%) | Probability value |
|-------------------|--------------------|----------------|-------------|--------------------|------------------------|-------------------|
| Among populations | 2 | 4.146 | 2.073 | 0.032 | 12 | 0.001 |
| Among accessions | 93 | 25.833 | 0.278 | 0.045 | 17 | 0.030 |
| Within accessions | 96 | 18.000 | 0.188 | 0.188 | 71 | 0.005 |
| Total | 191 | 47.979 | | 0.265 | 100 | |




Figure 9: Pie chart showing the distribution of genetic variation generated by 96 accessions divided into three groups viz landraces, commercial varieties and CIMMYT inbred lines.

6.3.3 Genetic relationship between groups of accessions

The observed Nei's pairwise population's differentiation (between landraces, commercial varieties and CIMMYT inbred lines) were shown in Table 18. The highest pairwise differentiation between populations (0.347), which explains the lowest similarity was between landraces and CIMMYT inbred lines. On the other hand, the lowest differentiation (0.086) which corresponds to the highest similarity was between landraces and commercial varieties.

Table 18: Pairwise genetic differentiation among the three populations, Nei's genetic identity (above diagonal) and genetic distance (below diagonal).

| Populations | CIMMYT lines | Commercial varieties | Landraces |
|-----------------------------|---------------------|-----------------------------|------------------|
| CIMMYT lines | **** | 0.834 | 0.707 |
| Commercial varieties | 0.181 | **** | 0.918 |
| Landraces | 0.347 | 0.086 | **** |

6.3.4 Genetic variation and diversity

The Nei's gene diversity (h) and Shannon's information index (I) analysis for the three groups of accessions are presented in Table 19. The Nei's gene diversity ranged from 0.212 (CIMMYT lines) to 0.475 (commercial varieties) with an average of 0.379. On the other hand, the Shannon index was highest in commercial varieties (0.667) and the lowest in CIMMYT lines (0.325) while the average was 0.544.

Table 19: Genetic diversity indices for the three groups of accessions.

| Populations | Nei's gene diversity (h) | Shannon's Information index (I) |
|-----------------------------|---------------------------------|--|
| CIMMYT lines | 0.212 | 0.325 |
| Commercial varieties | 0.475 | 0.667 |
| Landraces | 0.449 | 0.640 |
| Average | 0.379 | 0.544 |

6.3.5 Cluster and principal coordinate analyses

The cluster analysis was used to show more associations in terms of the genetic diversity among different groups of maize accessions through classical dendograms using Ward's algorithm method. The 96 accessions were clustered into five major groups (Fig. 11). The first (I) group consisted of 26 accessions with subgroup "a" containing a mixture of landraces and commercial varieties, while subgroup "b" contained only commercial varieties. The second (II) cluster consisted of only 14 landraces. Cluster three (III) contained 25 accessions which were mixed up with landraces, commercial varieties and just one accession (CKSBL10205) from CIMMYT in subgroup "b". The fourth (IV) group contained 18 accessions that were again a mixture of landraces and commercial varieties both in subgroup "a" and "b". The last group (V) clustered 13

accessions from CIMMYT inbred lines. However, within those 13 CIMMYT inbred lines, CML 489, CML 444, CML 445 were among the commercial varieties used in this study. On the other hand, the first two principal coordinates (Coordinate 1 and Coordinate 2), expressed 20.4% and 12.9% of the total variation in the SSR data, respectively (Fig. 10). The CIMMYT inbred lines clearly isolated themselves in the lower right quadrant, these lines clustered in one group different from the rest of other accessions. The other two groups (landraces and commercial varieties) were mixed and scattered along the four quadrants of the scatter plot plane (Fig. 10).

3.3.6 Genetic diversity associated with MLND resistance

The distribution and frequencies of alleles in two SSR loci phi029 (on chromosome 3) and phi062 (on chromosome 10) that are associated with QTLs against SCMV (Zhang *et al.*, 2003) were calculated for a group of accessions *viz* landraces, commercial varieties and CIMMYT inbred lines. The three alleles detected at phi029 locus had frequencies of 0.07, 0.36 and 0.07, respectively in landraces (Table 20). On the other hand, only one allele was detected at phi029 locus with frequencies of 0.15 and 0.35 for CIMMYT inbred lines and commercial varieties respectively. Furthermore, two alleles were detected at phi062 locus with frequencies of both 0.33 in landraces. The frequencies of 0.25 and 0.08 were observed in CIMMYT inbred lines and commercial varieties only in one allele at phi062 locus respectively (Table 20). The frequencies of all alleles detected at both phi029 and phi062 were not significantly ($p > 0.05$) different between accessions (Table 20).

Table 20: Allelic distribution and frequency of the three groups of accessions along with the two loci associated with SCMV resistance QTL's.

| <u>Locus</u> | | <u>phi029</u> | | | | <u>phi062</u> | | |
|-----------------------------|------------------|---------------------------|-----------------|-----------------|--------------|---------------------------|-----------------|--------------|
| | | <u>Allele 1</u> | <u>Allele 2</u> | <u>Allele 3</u> | <u>Total</u> | <u>Allele 1</u> | <u>Allele 2</u> | <u>Total</u> |
| Landraces | <i>Amplicons</i> | 4 | 20 | 4 | | 4 | 4 | |
| | <i>Frequency</i> | 0.07 | 0.36 | 0.07 | 0.51 | 0.33 | 0.33 | 0.67 |
| CIMMYT lines | <i>Amplicons</i> | 0 | 8 | 0 | | 0 | 3 | |
| | <i>Frequency</i> | 0.00 | 0.15 | 0.00 | 0.15 | 0.00 | 0.25 | 0.25 |
| Commercial varieties | <i>Amplicons</i> | 0 | 19 | 0 | | 0 | 1 | |
| | <i>Frequency</i> | 0.00 | 0.35 | 0.00 | 0.35 | 0.00 | 0.08 | 0.08 |
| Total | | | | | 1.00 | | | 1.00 |
| F - Statistics | | 3.103^{ns} | | | | 5.571^{ns} | | |




Figure 10: Relationships among the 96 accessions displayed by principal coordinate analysis using SSR data




Figure 11: Dendrogram for cluster analyses based on Ward's algorithm method for genetic distance among the 96 accessions analyzed using SSR.

6.4 Discussion

It is very important to have the details and knowledge about genetic diversity and genetic relationships within, between and among germplasm for crop improvement programs. The variation between germplasm revealed at DNA level through molecular markers grant an opportunity to have an effective means for germplasm utilization. In this study SSR markers were used to study genetic diversity among different groups of maize accessions *viz* landraces, commercial varieties and CIMMYT inbred lines as relates to MLND resistance. A total of 63 alleles with polymorphic bands were observed in 96 accessions evaluated in this study using 20 SSR markers. All these 20 SSR loci expressed polymorphism with an average of 3.15 alleles per primer. Other studies reported allelic detection that was either lower or higher than what was obtained in this study. For instance, lower number of alleles per locus of 2.87 and 2.9 with higher number of SSR markers of 30 and 47 than in the current study were obtained by Lopes *et al.* (2015) and Mollin *et al.* (2013), respectively. On the other hand, Reid *et al.* (2011) and Enoki *et al.* (2002) generated higher number of alleles per locus that is 3.62 and 7.3, respectively than in the current study using 105 and 60 SSR markers. Legesse *et al.* (2007) stated that genetic diversity is the most important aspect that limits the number of alleles detected at each SSR locus. On the other hand, Terra *et al.* (2011) gave a reason for different levels of allele number detection being the types of gel used during electrophoresis process. The use of agarose gels is associated with low number of alleles, while polyacrylamide gels provide a very high detection power for even small fragments (Mollin *et al.*, 2013; Terra *et al.*, 2011; Legesse *et al.*, 2007). In this study, we employed agarose gel for running electrophoresis as used by Lopes *et al.* (2015) and Mollin *et al.* (2013). Reid *et al.* (2011) and Enoki *et al.* (2002) on the other hand, detected the amplification using polyacrylamide gels. Furthermore, polymorphism information content (PIC) expresses the ability of the SSR markers to detect variations within germplasm as caused by the number of alleles per locus and the distribution of those alleles (Nyaligwa *et al.*, 2015). The PIC in the current study ranged from 0.17 to 0.79, with an average of 0.52 which was as well reported by Terra *et al.* (2011). Also Nyaligwa *et al.* (2015) obtained an average of 0.51 PIC. Legesse *et al.* (2007) reported 0.55 and Lopes *et al.* (2015) obtained a PIC of 0.41. PIC indicates the discriminatory power of primers which is associated with the variation occurring at the chromosome region of the accessions evaluated. Therefore, expression of lower PIC indicate

that the chromosomal region marked with a certain primer is conserved in a group of germplasm under study (Terra *et al.*, 2011). On the other hand, loci that generate higher PIC indicate their ability to discriminate between germplasm and hence chromosomal region is variable between individuals within a group of germplasm (Lopes *et al.*, 2015). The PIC in this study provided a room for discrimination and expression of genetic diversity within germplasm.

Analysis of molecular variance in this study showed that 71% of the total variation was accounted for by the variation within accessions and the rest (29%) was divided between among accessions (17%) and among populations (12%). This is explained by the characteristics of cross pollination expressed by plant species like maize which exhibit enormous heterogeneity that may cause SSR variability within as well as among accessions (Hu *et al.*, 2007). For self-pollinating plant species, considerable genetic diversity is found among cultivars while little is found within a cultivar (Jensen *et al.*, 2006). The low level of variation among populations indicate the occurrence of gene flow which is generally caused by seed mixing and pollen contamination (Nie *et al.*, 2014; Louette *et al.*, 1997). Thus, accessions used in this study represent the three groups from different sources and improvement status but also there is a certain level of interaction between them. This was evidenced by the general low level of differentiation between the three groups, where the highest 0.347 pairwise genetic distance in this study was between landraces and CIMMYT inbred lines which was actually still low. The exchange and movement of seeds between research institutions, breeding programs, seed companies, as well as between farmers still occur at regional level and in Africa at large (Bøhn *et al.*, 2016). The Nei's gene diversity within each group was highest in commercial varieties and closely followed by landraces and lastly was the CIMMYT lines. The commercial varieties were comprised of varieties from different companies and different breeding programs, it also included the OPVs that were selected and developed before being commercialized. The landraces that formed the largest group of accessions had also higher diversity as expected and they were collected from different regions in Tanzania where at each region the diversity is constituted within regions' independent selection for specific desired traits (Beyene *et al.*, 2006). However, instead of landraces possessing the highest gene diversity than the rest of the group as expected due to their nature of heterogeneity, they had the diversity slightly lower than the commercial varieties

probably because of seed mixing done by farmers through exchange as their strategy to improve crop performance (Louette *et al.*, 1997). As for the CIMMYT inbred lines having the lowest gene diversity, this might be because the accessions in this group were obtained from the same source with possibly similar breeding programs. This finding is supported by Lu *et al.* (2009) who characterized 394 CIMMYT global maize and could not differentiate them into groups. The cluster analysis generated through UPGMA using Ward's algorithm method grouped the accessions into five clusters. However, the clustering of accessions was not independent to the designated group of accessions as were collected from different sources. It was only the CIMMYT lines that exhibited a unique pattern of clustering on dendrogram as well as on scatter plot with principal coordinates analysis. The grouping may express associations related to their pedigree records (Legesse *et al.*, 2007), but also could be due to the effect of selection, drift and mutation (Warburton *et al.*, 2002).

The extent of *Sugarcane mosaic virus* (SCMV) resistance that might give an indication for MLND resistance was examined with accessions in this study. The accessions were examined following the presence of alleles associated with QTL's on resistance to SCMV. The distribution and frequencies of alleles in two SSR loci phi029 (on chromosome 3 bin 3.04) and phi062 (on chromosome 10 bin 10.04) that are associated with those QTL's against SCMV (Zhang *et al.*, 2003) were calculated for a group of accessions *viz* landraces, commercial varieties and CIMMYT inbred lines. Those two loci are located on genes and QTLs that are clustered in chromosomal regions associated with multiple viral pathogen resistance and in particular *Sugarcane mosaic virus* (SCMV) resistance (Wiser *et al.*, 2006; Zhang *et al.*, 2003). Therefore, it was worth contemplating that these loci provide a room for promising alleles that link to resistance traits against *Sugarcane mosaic virus* (SCMV) as well as other viral pathogens and diseases. The detection and distribution of all alleles at both loci were observed in landraces while in other groups of accessions only one allele at each locus was detected. Also the detection frequency was high in landraces as compared with the other groups. The associations with alleles attached to the loci within the chromosomal region of the QTLs for SCMV resistance are expected to enhance adaptability to a varying level of stresses caused by the virus. There was also no significant difference in allele frequency and distribution between both landraces and

commercial varieties with CIMMYT inbred lines which had phenotypically resistant accessions to MLND. However, SCMV constitute just a part of a synergistic interaction with *Maize chlorotic mottle virus* (MCMV) which is the major pathogen in triggering *maize lethal necrosis* (MLN) disease (Mahuku *et al.*, 2015). Furthermore, little is understood about MCMV's genetics for resistance in maize, though promising tolerant genotypes have been grown and identified (Mahuku *et al.*, 2015; Redinbaugh and Zambrano, 2014; Nelson *et al.*, 2011). In addition, prevention or resistance to one virus that synergistically interact with another is anticipated to reduce the extent of the occurring disease. Thus establishment of genetic details associated with resistance against SCMV provided an insight towards understanding the utility function of the occurring genetic diversity of the studied accessions for MLND resistance.

6.5 Conclusion

Maize supports millions of people in sub-Saharan Africa and other parts of the world as an important staple food as well as a cash crop. However, maize production is challenged by a number of diseases which cause significant yield loss. Recently, maize production in East Africa has been brought into even more challenge due to an outbreak of MLND which is reported to cause up to complete loss of the crop in the field. The method to control the disease which is most reliable and economically feasible is through breeding for genetically resistant varieties. Successful maize breeding requires the availability of reliable genetic diversity to help in the identification and estimation of the level of expected heterosis from germplasm as well as the level of variability during breeding. Therefore, good understanding of genetic diversity within and among maize accessions ensures effective utilization of the genetic resource available for the fight against the current major challenge (MLND) in maize production. A total of 63 alleles and a mean of 3.15 alleles per locus observed with the 20 SSR markers as well as the average PIC of 0.52 provided a room for discriminating and realizing genetic diversity within accessions. More than half of the total molecular variation (71%) was recorded within accessions and the rest (29%) was divided among accessions and among populations. That means accessions used in this study had high diversity within themselves. The Nei's gene diversity was highest with commercial varieties and closely followed by landraces and lastly was the CIMMYT lines. The locus phi029 on chromosomal bin location at 3.04 and phi062 on bin 10.04 are the loci

previously identified to be associated with quantitative trait loci (QTL) responsible for *Sugarcane mosaic virus* (SCMV) resistance. The distribution and frequencies of alleles in those two SSR loci were calculated and found higher with landraces than the rest of the group. However, SCMV constitute just a part of a synergistic interaction with *Maize chlorotic mottle virus* (MCMV) which is the major pathogen in triggering for maize lethal (MLN) disease. Therefore, establishment of the genetic details associated with resistance against SCMV provided just an insight towards understanding the utility function of the occurring genetic diversity within studied accessions for MLND resistance.

CHAPTER SEVEN

General discussion, conclusion and recommendations

7.1 General discussion

Maize continues to be the most preferred staple food as well as a cash crop in Tanzania and in other parts of the world (Romay *et al.*, 2013). However, maize is affected by many pathogens and some of them cause significant yield loss (Ali and Yan, 2012). The outbreak of a new disease in East Africa, MLND, presents immediate concerns as well as uncertain long-term consequences (Kabululu *et al.*, 2017). MLND infection rate reaches 100% and yields severely affected up to a complete loss of the crop (Adams *et al.*, 2013). Genetic diversity (morphological and molecular) study provides key information that may help in identifying important traits against several production constraints (Enoki *et al.*, 2002). Hence, good understanding of genetic diversity within and among maize accessions ensures effective utilization of the genetic resource available for the fight against the current major challenge (MLND) in maize production. A total of 51 maize landraces from Tanzania were evaluated in different experiments along with 34 commercial varieties and 13 elite lines from CIMMYT Kenya to study their potential genetic diversity and how the diversity responds against MLND. The significant differences expressed among those accessions and their interactions with environments suggest that the accessions were different (variable) from each other and they could respond differently in different environments and stresses. This variability is what we desire to use in breeding programs to establish improved maize varieties required by the community for specific traits and even for specific location (Kabululu *et al.*, 2017). Successful plant breeding then requires careful choices of genotypes that possess desirable traits for best combinations. The correlations between traits are also another important knowledge that helps on predicting the required performance in terms of certain traits. For example, the negative correlation between yield and flowering parameters help to have an indirect selection of higher yielding genotypes through flowering character.

Results in this current study showed that an OPV Situka 1 and a hybrid DH 04 were generally the best performing genotypes in terms of grain yield and stability as well as for other related parameters across all the three locations. However, TZA 2793 was a local cultivar that expressed

promising performance for yield. The maize landraces have always been considered less productive than other improved varieties, but they present an important source of genetic variability that can be exploited to search for genes against biotic and abiotic stresses. The flowering character defines the maturity differences among accessions (Olaoye, 2009) and it can also be connected to the yielding ability that early maturing accessions could generate high grain yield while those which are late maturing produce low yield (Lafitte *et al.*, 1997). However, the opposite also holds true under different circumstances where late maturing varieties produce high yield due to their ability to have longer periods to accumulate maximum assimilates into grains than the early maturing varieties (Wang *et al.*, 2011; Bello *et al.*, 2012a). The analysis in this study showed a significant negative correlation between both days to 50% anthesis and days to 50% silking to all the yield related parameters. A wide range of variation in flowering characters could signify the potential variability within accessions that helps on developing genotypes adaptable to different areas with different characteristics (Cömertpay, 2012).

Quantitative morphological traits that highly contributed to the total variation expressed by the accessions under study included 1000 kernel weight, plant height, ear height, yield per plant and days to 50% silking. These traits could be used to characterize several maize accession populations and discover potential candidates as parents for generating elite materials. The study identified distinction of the three groups of accessions used in this study where commercial varieties were discriminately identified by high yield related parameters and early flowering characteristics. CIMMYT elite lines were characterized by significant small plant structures and landraces were extensively diverse. For qualitative morphological traits, the percent frequency distribution of accessions within traits differentiated the three groups in terms of foliage, tassel size, shape of upper surface of kernel and kernel type. Other traits of stem colour, sheath pubescence, tassel type, cob colour, shape of upper most ear, kernel row arrangement, kernel colour and endosperm colour characterized the three groups similarly though with different percent distribution. The former traits were able to discriminate between and within the three groups while the later identified differences just within each group. Traits that had higher percentage distribution of accessions towards one class within a trait include stem colour (green), sheath pubescence (intermediate), tassel type (primary-secondary), cob colour (white), shape of

uppermost ear (conical), shape of kernel upper surface (rounded), kernel row arrangement (regular), kernel colour (white) and endosperm colour (white). This trend of accessions being distributed towards a certain class of trait reflects farmers' or consumers' preferences through successive selection (Ntundu *et al.*, 2006; Louette and Smale, 2000). On the other hand, the grouping of the accessions through cluster analysis reflected individual performance and type of accessions. Most of the landraces lacked consistent grouping in terms of collection sites. This was related to the finding by Sun *et al.* (2016) who observed geographically close populations of Chinese sweetgum in different clusters. The implication in this finding is that the landraces involved in this study were comprised of a heterogeneous group that occurred through repeated exchange and selection of germplasm executed by farmers (Ntundu *et al.*, 2006).

Having the distinct and diverse groups of accessions in this study, tests were made to see how this diversity results into responding against MLND. The standardized relative area under the disease progress curve (rAUDPC) was utilized to assess the cumulative disease progression within the growing process of the crop plants as explained by Luitel *et al.* (2016). The rAUDPC was higher at Naivasha than at Mlangarini because the extent of disease pressure at Naivasha was high, due to the artificial inoculation which generated maximum pathogen inoculum applied on plants. On the other hand, natural disease pressure at Mlangarini generated low disease pressure. The rAUDPC evaluated how the genetic differences among the accessions determine the way they react on the extent of disease development on plants (Skelsey and Newton, 2014). The accessions with the lowest rAUDPC were considered to be resistant to the disease and vice versa (Safavi *et al.*, 2010). Also, the resistant performance is determined by comparing the accessions with the performance of the established susceptible and resistant check materials (Massa *et al.*, 2015). Therefore, accessions that were able to show the least disease progression trend in both locations could be considered promising for resistance indication against MLND. In this study, accessions TZA2793 and TZA3544 had among the lowest rAUDPC, but across the two locations and may be considered for further screening to ascertain their resistance characteristics against MLND. The occurrence of the disease at Naivasha enhanced high heritability (0.86) as compared with the low heritability (0.18) at Mlangarini. This was due to the fact that uniform environmental conditions ensure high level of heritability while variable

environments lower the level of heritability (Brunda *et al.*, 2014). The amount of inoculum applied on the plants at Naivasha was actually uniform such that genetic variation of the accessions in this study exerted an influence responsible for the variation in response against MLND. On the other hand, the Mlangarini situation was different because the amount of pathogens occurred naturally with no regular inoculum pressure distributed on plants. Heritability measures phenotypic variance as caused by genetic variation, it provides prediction for plant breeding strategies.

The correlation between sheath pubescence and the occurrence of MLN disease was significantly negative. That means, an increase of pubescence is associated with lowering the disease (enhance resistance) to the crop plants. The resistance might directly be the close link of the pubescent marker and the gene controlling resistance or else the indirect link through the prevention of the insect pests carrying the pathogen causing the disease. Mmbaga *et al.* (1996) found the presence of dense pubescence to be linked to the resistance of rust on common bean and the pubescence trait was merely inherited. Endosperm colour is another character which was negatively correlated to the occurrence of MLN disease. The descriptor for endosperm colour used for scoring, ranged from white, cream, pale yellow, yellow, orange and white cap. The higher number for endosperm colour signified for more coloured endosperm such as yellow or orange, while the lower number associated with whitish colour. Therefore, as the colour become more yellow or orange, it is associated with resistance, and the white endosperm is related to susceptibility. Scott (1989) studied the link between endosperm colour and potyvirus resistance and found that the gene for yellow endosperm colour was linked to the resistance on potyviral disease. The promising inbred lines resistant to MLND that had been developed so far have yellow endosperm colour (Semagn *et al.*, 2015). Anthesis-silking interval (ASI) was significant and positively correlated to the occurrence of the MLN disease, which means the longer the ASI the higher the chance for possible susceptibility, while short ASI indicated the possible resistance. Ngugi *et al.* (2013) established the facts that relate to this finding, where Anthesis-Silking Interval was significant and positively correlated with stress susceptible index. ASI is to a large extent determined by the variation in silking number of days where a strong association of their Quantitative Trait Loci (QTLs) confirms the close link of the two traits (Gemenet *et al.*,

2010). This imply that short ASI depends on the short duration of the silking from the planting date (Magorokosho *et al.*, 2003) and a genotype with short days to silking have a chance to be able to tolerate or resist growing stresses like diseases and drought. Kernel type was also significantly correlated with the disease scores such that flint kernel type was found to associate with resistance traits. The indication of kernel type being correlated to disease resistance traits was also found by Marçon *et al.* (1996), but here however, they learned that dent kernel types had the relative resistance potential against high plains virus while sweet corn (sweet kernel type) expressed high susceptibility. On testing the maize accessions in this study, the cluster analysis brought a landrace TZA2793 grouped together with the check varieties that are moderately resistant CKDHL0500, CKSBL10205, CKDHL120552 and resistant CLRCY039, CLYCN261, CLRCY034. The landrace could be considered for other tests to confirm its potential for MLND resistance.

It is very important to have the details and knowledge about genetic diversity and genetic relationships within, between and among germplasm for crop improvement programs. The variation between germplasm revealed at DNA level using molecular markers grant an opportunity to have effective means for germplasm utilization. In this study we used SSR markers to study genetic diversity among different groups of maize accessions *viz* landraces, commercial varieties and CIMMYT inbred lines as related to MLND resistance. A total of 63 alleles were observed within 96 accessions evaluated in this study using 20 SSR markers. All these 20 SSR loci expressed polymorphism with an average number of 3.15 alleles per primer. The Polymorphism Information Content ranged from 0.17 to 0.79, with an average of 0.52 was comparable to that reported by Terra *et al.* (2011). Analysis of molecular variance in this study showed that more than half of the total variation (71%) was accounted for the variation within accessions where 17% variations was attributed to among accessions and 12% of variation to among populations. This is explained by the characteristics of cross pollination expressed by plant species like maize which exhibit enormous heterogeneity that may cause SSR variability within as well as among accessions (Hu *et al.*, 2007). The low level of variations among populations indicates the occurrence of gene flow that is generally caused by seed mixing and pollen contamination (Nie *et al.*, 2014; Louette *et al.*, 1997). Thus, accessions used in this study

represent the three groups from different sources and improvement status but also a certain level of interaction between them. This was evidenced by the general low level of differentiation among the three groups, where the highest 0.347 pairwise genetic distance in this study between landraces and CIMMYT inbred lines were actually still low. The exchange and movement of seeds between research institutions, breeding programs, seed companies, as well as between farmers still occur at regional level and in Africa at large (Bøhn *et al.*, 2016). The cluster analysis generated through UPGMA using Ward's similarity matrix grouped the accessions with no independency to the designated group of accessions collected from different sources. Only CIMMYT lines grouped themselves together. The grouping may express associations related to their pedigree records (Legesse *et al.*, 2007), but also could be due to the effect of selection, drift and mutation (Warburton *et al.*, 2002).

The extent of *Sugarcane mosaic virus* (SCMV) resistance that might give an indication for MLND resistance was examined with accessions in this study. The accessions were examined following the presence of alleles associated with QTL's on resistance to SCMV. The distribution and frequencies of alleles in two SSR loci phi029 and phi062 were calculated for the accessions evaluated in this study. Those SSR loci phi029 on chromosome 3 bin 3.04 and phi062 on chromosome 10 bin 10.04 are associated with QTL's against SCMV (Zhang *et al.*, 2003). The loci are located on genes and QTLs that are clustered in chromosomal regions associated with multiple viral pathogen resistance and in particular *Sugarcane mosaic virus* (SCMV) resistance (Wiser *et al.*, 2006; Zhang *et al.*, 2003). Therefore, it was worth contemplating that these loci provide a room for promising alleles that link to resistance traits against *Sugarcane mosaic virus* (SCMV) as well as other viral pathogens and diseases. The detection and distribution of all alleles at both loci were observed in landraces while in other groups of accessions only one allele at each locus was detected. The detection frequency was high in landraces as compared with the other groups. The association with alleles attached to the loci within the chromosomal region of the QTLs for SCMV resistance is expected to enhance adaptability to a varying level of stresses caused by the virus. However, SCMV constitute just a part of a synergistic interaction with *Maize chlorotic mottle virus* (MCMV) which is little understood about its genetics for resistance in maize (Mahuku *et al.*, 2015; Redinbaugh *et al.*, 2014; Nelson *et al.*, 2011). Thus,

establishment of genetic details associated with resistance against SCMV provided an insight towards understanding the utility function of the occurring genetic diversity for MLND resistance.

7.2 Conclusion

Maize genome harbors potential amount of morphological and molecular diversity that can be sourced and invested for maize crop improvements. Therefore, good understanding of genetic diversity within and among maize accessions ensures effective utilization of the genetic resource available for resistance against the current major challenge (MLND) in maize production. The results in this study have revealed a significant range of genetic diversity in maize accessions evaluated. This might provide a source of variation required for breeding programs to hold back the genetic vulnerability as a result of recurrent outbreaks of new strains of pest and diseases. It also offers an opportunity to widen the genetic background of the available maize germplasm because the materials that are currently at disposal for several breeding programs are composed of narrow genetic base. Agronomic evaluation revealed that TZA 2793 is the most promising accession on yield. Cluster analysis also identified two landraces, TZA 2793 and TZA 5170 grouped with commercial varieties that are mostly higher yielding. The cluster analysis also disclosed the expression of landraces lacking regular pattern in clustering within their major group. This elucidate the fact that farmers select cultivars based on their preferences and also exchange seed crop materials with fellow farmers even from very distant regions. Farmers play a significant role in shaping the structure of landrace population existing in a certain area.

The assessment of accessions' performance on MLND progression over time, showed accessions TZA2793 and TZA3544 to have the lowest rAUDPC in both locations and may be considered stable and promising for resistance characteristics against the disease. Pearson correlation matrix showed sheath pubescence, kernel type and endosperm colour to be negatively correlated with MLN disease scores while anthesis-silking interval, and kernel type were positively correlated with MLND scores. Cluster analysis associated TZA2793 with moderate and resistant variety checks and grouped them in the same cluster. Molecular genetic diversity study revealed that 71% of the total molecular variation was recorded within accessions and the rest (29%) was

divided among accessions with 17% and 12% among populations. That means accessions used in this study had the high level of diversity within themselves.

The distribution and frequencies of alleles in phi029 and phi062 loci previously identified to be associated with quantitative trait loci (QTL) responsible for *Sugarcane mosaic virus* (SCMV) resistance were found to be higher with landraces. Thus, the established genetic detail associated with resistance against SCMV provides an insight towards understanding the utility function of the occurring genetic diversity for MLND resistance.

7.3 Recommendations

Lack of regular clustering pattern in terms of regions of collection calls for systematic involvement of farmers in breeding and selection processes through participatory breeding in order to have an organized process of population structuring.

The significant range of genetic diversity revealed in maize landraces evaluated in this study showed the need to establish extensive molecular and morphological genetic details of a vast germplasm materials kept in the Genebanks, research institutions and farmers in order to have the required allelic information for crop improvements.

Accessions TZA 2793, TZA 5170 and TZA 3544 expressed promising performance on yield and response against MLND, hence could further be investigated to utilize their potentials for future and imminent maize crop improvement especially resistance against MLND.

REFERENCES

- Abdellatif, K. F. and Khidr, Y. A. (2010). Genetic diversity of new maize hybrids based on SSR markers as compared with other molecular and biochemical markers. *Journal of Crop Science and Biotechnology*. **13**(3): 139 - 145.
- Abdel-Mawgood, A. L. (2012). *DNA based techniques for studying genetic diversity, genetic diversity in microorganisms*. Prof. Mahmut Caliskan (Ed.), ISBN: 978-953-51-0064-5, *InTech*. <http://cdn.intechopen.com/pdfs-wm/28889.pdf>. Accessed on 25 May 2016.
- Abraha, T., Githiri, S. M., Kasili, R. W., Skilton, R. A., Solomon, M. and Nyende, A. B. (2014). Genetic diversity analysis of Eritrean Sorghum (*Sorghum bicolor* (L.) Moench) germplasm using SSR markers. *Molecular Plant Breeding*. **5**(13): 1 - 12. <http://mpb.biopublisher.ca>.
- Aci, M. M., Revilla, P., Morsli, A., Djemel, A., Belalia, N., Kadri, Y, ... Khelifi, L. (2013). Genetic diversity in Algerian maize (*Zea mays* L) landraces using SSR markers. *Maydica*. **58**: 304 - 310.
- Adams, I. P., Harju, V. A., Hodges, T., Hany, U., Skelton, A., Rai, S, ... Boonham, N. (2014). First report of maize lethal necrosis disease in Rwanda. *New Disease Reports*. **29**: 22. <http://dx.doi.org/10.5197/j.2044-0588.2014.029.022>.
- Adams, I. P., Miano, D. W., Kinyua, Z. M., Wangai, A., Kimani E., Phiri N, ... Boonham, N. (2013). Use of next generation sequencing for the identification and characterization of *Maize chlorotic mottle virus* and *Sugarcane mosaic virus* causing maize lethal necrosis in Kenya. *Plant Pathology*. **62**: 741 - 749. (Published Online: Doi: 10.1111/j.1365-3059.2012.02690.x).
- Al-Badeiry, N. A., Al-Saadi, A. H. and Merza, T. K. (2014). Analysis of genetic diversity in maize (*Zea mays* L.) varieties using simple sequence repeat (SSR) markers. *Journal of Babylon University*. **22** (6): 1768 - 1774.
- Ali, R. and Yan, J. (2012). Disease resistance in maize and the role of molecular breeding in defending against global threat. *Journal of Integrative Plant Biology*. **54** (3): 134 - 151.
- Anjichi, V. E., Muasya, R. M., Gohole, L. S., Rao, N. K. and Muui, C. W. (2005). Genetic biodiversity assessment and local seed systems of maize landraces among smallholder

- farmers in Western Kenya. *African Crop Science Conference Proceedings*. **7**: 1335 - 1340.
- Anumalla, M., Roychowdhury, R., Geda, C. K., Mazid, M. and Rathoure, A. K. (2015). Utilization of plant genetic resources and diversity analysis tools for sustainable crop improvement with special emphasis on rice. *International Journal of Advanced Research*. **3**: 1155 - 1175.
- Arif, I. A., Bakir, M. A., Khan, H. A., Al Farhan, A. H., Al Homaidan, A. A., Bahkali, A. H, ... Shobrak, M. (2010). A Brief review of molecular techniques to assess plant diversity. *International Journal of Molecular Science*. **11**: 2079 - 2096.
- Asare, S., Tetteh, A. Y., Twumasi, P., Adade, K. B. and Akromah, R. A. (2016). Genetic diversity in lowland, midaltitude and highland African maize landraces by morphological trait evaluation. *African Journal of Plant Science*. **10**(11): 246 - 257.
- Ashimogo, G. and Rukulantile, H. (2000). Danish International Development Agency (DANIDA). Agricultural Sector Programme Support. Consultancy on seed market assessment in Dodoma, Iringa and Morogoro regions. Final report. June 2000. Faculty of Agriculture, Sokoine University of Agriculture. Morogoro, Tanzania. pp. 37.
- Bello, O. B., Abdulmalik, S. Y., Ige, S. A., Mahamood, J., Oluleye, F., Azeez, M. A. and Afolabi M. S. (2012a). Evaluation of early and late/intermediate maize varieties for grain yield potential and adaptation to a Southern Guinea Savanna agro-ecology of Nigeria. *International Journal of Plant Research*. **2**(2): 14 - 21
- Bello, O. B., Ige, S. A., Azeez, M. A., Afolabi, M. S., Abdulmalik, S. Y. and Mahamood, J. (2012b). Heritability and genetic advance for grain yield and its component characters in maize (*Zea mays* L.). *International Journal of Plant Research*. **2**(5): 138 - 145.
- Benson, J. M., Poland, J. A., Benson, B. M., Stromberg, E. L. and Nelson, R. J. (2015). Resistance to Gray leaf spot of maize: Genetic architecture and mechanisms elucidated through nested association mapping and Near-Isogenic Line analysis. *PLoS Genetics*. **11**(3): 1 - 23. doi:10.1371/journal.pgen.1005045.
- Beyene, Y., Botha, A. and Myburg, A. A. (2005). A comparative study of molecular and morphological methods of describing genetic relationships in traditional Ethiopian highland maize. *African Journal of Biotechnology*. **4**: 586 - 595.

- Beyene, Y., Botha, A. and Myburg, A. A. (2006). Genetic diversity among traditional Ethiopian highland maize accessions assessed by simple sequence repeat (SSR) markers. *Genetic Resources and Crop Evolution*. **53**:1579 - 1588.
- Bindroo, B. B. and Moorthy, S. M. (2014). Genetic divergence, implication of diversity, and conservation of Silkworm, *Bombyx mori* . *International Journal of Biodiversity*. **2014**: 1 - 15.
- Biniam, M. G., Githiri, S. M., Tadesse, M. and Kasili, R. W. (2015). Analysis of Diversity among Potato Accessions Grown in Eritrea Using Single Linkage Clustering. *American Journal of Plant Sciences*. **6**: 2122 - 2127.
- Bøhn, T., Aheto, D. W., Mwangala, F. S., Fischer, K., Bones, I. L., Simoloka, C, ... Breckling, B. (2016). Pollen-mediated gene flow and seed exchange in small-scale Zambian maize farming, implications for biosafety assessment. *Science Report*. **6** (34483): 1-12; doi: 10.1038/srep34483.
- Bolafios, J. and Edmeades, G. O. (1996). The importance of the anthesis-silking interval in breeding for drought tolerance in tropical maize. *Field Crops Research*. **48**: 65 - 80.
- Brown-Guedira, G. L., Thompson, J. A., Nelson, R. L. and Warburton, M. L. (2000). ‘Evaluation of genetic diversity of soybean introductions and North American ancestors using RAPD and SSR markers’. *Crop Science*. **40**: 815 - 823.
- Brunda, S. M., Kamatar, M.Y., Naveenkumar, K. L. and Hundekar, R. (2014). Study of genetic variability, heritability and genetic advance in foxtail millet in both rainy and post rainy season. *IOSR Journal of Agriculture and Veterinary Science*. **7**(11): 34 - 37.
- Bucheyeki, T. L. (2012). Characterization and genetic analysis of maize germplasm for resistance to northern corn leaf blight disease in Tanzania. Unpublished Dissertation for Award of PhD Degree at University of KwaZulu-Natal, Pietermaritzburg. <http://hdl.handle.net/10413/8730>. Accessed on 02 February 2017.
- Camacho, V. T. C., Maxted, N., Scholten, M. and Ford-Lloyd, B. (2005). Defining and identifying crop landraces. *Plant Genetic Resources*. **3**: 373 - 384.
- Carvalho, V. P., Ruas, C. F., Josué, M., Ferreira, J. M., Moreira, R. M. P. and Ruas, P. M. (2004). Genetic diversity among maize (*Zea mays* L.) landraces assessed by RAPD markers. *Genetics and Molecular Biology*. **27**(2): 228 - 236.

- CGIAR. (2012). Research Program MAIZE Annual Report. Mexico, D.F.: CIMMYT.. <http://libcatalog.cimmyt.org/download/cim/98018.pdf>. Accessed on 29 March 2017.
- CIMMYT. (2013). Maize lethal necrosis (MLN) disease in Kenya and Tanzania: Facts and actions. [http://www.fao.org/fileadmin/user_upload/drought/docs/1%20%20Maize%20Lethal%20Necrosis-\(MLN\)%20Facts%20and%20Actions.pdf](http://www.fao.org/fileadmin/user_upload/drought/docs/1%20%20Maize%20Lethal%20Necrosis-(MLN)%20Facts%20and%20Actions.pdf). Accessed on 27 March 2017.
- Cömertpay, G., Baloch, F. S., Kilian, B., Ahmet, C. Ü. and Hakan, Ö. (2012). Diversity assessment of Turkish maize landraces based on fluorescent labelled SSR markers. *Plant Molecular Biology Reporter*. **30**: 261 - 274.
- De Groote, H., Oloo, F., Tongruksawattana, S. and Das, B. (2016). Community-survey based assessment of the geographic distribution and impact of maize lethal necrosis (MLN) disease in Kenya. *Crop Protection*. **82**: 30 - 35.
- De Jaramillo, E. H. (2009). *Ecological aspects of biosafety*. In: *Biosafety of genetically modified organisms: Basic concepts, methods and issues*. [Chowdhury, M. K. A, Hoque, M. I. and Sonnino, A. (Eds.)]. © FAO 2009, pp 51 - 105.
- Dellaporta, S. L., Wood, J. and Hiskis, J. B. (1983). A plant DNA mini-preparation. Version II. *Plant Molecular Biology Reporter*. **4**: 19 - 21.
- Dogan, Y., Ekinici, M. B., Togay, N. and Togay, Y. (2015). Determination of suitable nitrogen doses for growing second product maize (*Zea mays* L.) varieties in chickpea planting fields and its economic analysis. *Indian Journal of Agricultural Research*. **49** (2): 125 - 133.
- Dowswell, C. R., Paliwal, R. L. and Cantrell, R. L. (1996). *Maize in the Third World*. Westview Press, USA.
- Drinic, S. M., Andjelkovic V. and Micic, D. I. (2012). *Genetic diversity of maize landraces as sources of favorable traits. The molecular basis of plant genetic diversity*. Prof. Mahmut Caliskan (Ed.), ISBN: 978-953-51-0157-4, *InTech*.
- DSMZ. (2014). Data sheet on Maize Lethal Necrosis (MLN) Disease. DS-MLND_V1.0 04/2014. https://www.dsmz.de/fileadmin/Bereiche/PlantVirusesAndAntisera/Dateien/Data-MLND_V1-0.pdf. Accessed on 27 March 2017.

- Dubreuil, P., Warburton, M., Chastanet, M., Hoisington, D. and Charcosset, A. (2006). 'More on the introduction of temperate maize into Europe: large-scale bulk SSR genotyping and new historical elements'. *Maydica*. **51**: 281 - 291.
- Durga, K. K., Ankaiah, R. and Ganesh, M. (2015). Characterization of horse gram cultivars using plant morphological characters. *Indian Journal of Agricultural Research*. **49**: 215 - 221.
- Edmeades, G.O., Bolanos, J., Hernandez, M. and Bello, S. (1993). Causes for silk delay in lowland tropical maize. *Crop Science*. **33**: 1029 - 1035.
- Enoki, H., Sato, H. and Koinuma, K. (2002). SSR analysis of genetic diversity among maize inbred lines adapted to cold regions of Japan. *Theoretical and Applied Genetics*. **104**: 1270–1277.
- European Seed Association (ESA). (2010). *Biodiversity and plant breeding*.
- Forbes, G., Pérez, W. and Andrade, P. J. (2014). Field assessment of resistance in potato to *Phytophthora infestans*. Lima (Peru). International Potato Center (CIP). 35 p. DOI: 10.4160/9789290604402. <http://cipotato.org/wp-content/uploads/2014/06/006154.pdf>. Accessed on 29 March 2017.
- Frese, L., Bjorn, G. K., Branca, F., Ford-Lloyd, B. V., Germeier, C. U., Iriondo, J. M., ... de Carvalho, M. A. A. P. (2012). *Genetic reserve conservation of European crop wild relative and landrace diversity*. In: *Agrobiodiversity conservation* by Maxted, N., Ehsan, D. M., Ford-Lloyd, B. V., Frese, L., Iriondo, J. M., Pinheiro de Carvalho, M. A. A. (Eds.). CABI, Wallingford, UK, **1**: 1 - 6.
- Gemenet, D. C., Wachira, F. N., Pathak, R. S. and Munyiri, S. W. (2010). Identification of molecular markers linked to drought tolerance using bulked segregant analysis in Kenyan maize (*Zea mays* L.) landraces. *Journal of Animal and Plant Sciences*. **9**(1): 1122 - 1134.
- George, M. L., Regalado, E., Warburton, M., Vasal, S. and Hoisington, D. (2004). Genetic diversity of maize inbred lines in relation to downy mildew. *Euphytica*. **135**: 145 - 155.
- Govindaraj, M., Vetriventhan, M. and Srinivasan, M. (2015). Importance of genetic diversity assessment in crop plants and its recent advances: An overview of its analytical perspectives. *Genetics Research International*. 2015: 1 - 14.

- Gowda, M., Das, B., Makumbi, D., Babu, R., Semagn, K., Mahuku, G, ... Prasanna, B. M. (2015). Genome-wide association and genomic prediction of resistance to maize lethal necrosis disease in tropical maize germplasm. *Theoretical and Applied Genetics*. **128**:1957 - 1968.
- Gudu, S. O., Okalebo, J. R., Othieno, C. O., Obura, P. A., Ligeyo, D. O., Shulze D. and Johnston, C. (2005). Response of five maize genotypes to nitrogen, phosphorus and lime on acid soils of western Kenya. *African Crop Science Conference Proceedings*. **7**: 1109-1115.
- Guruprasad, M., Sridevi, V., Vijayakumar, G., and Kumar. M. S. (2016). Plant regeneration through callus initiation from mature and immature embryos of maize (*Zea mays* L.). *Indian Journal of Agricultural Research*. **50**(2): 135 - 138.
- Hagenblad, J., Oliveira, H. R., Nils E. G., Forsberg, N. E. G. and Leino, M. W. (2016). Geographical distribution of genetic diversity in *Secale landrace* and wild accessions. *BMC Plant Biology*. **16**: 1 - 20.
- Hartings, H., Berardo, N., Mazzinelli, G. F., Valoti, P., Verderio, A and Motto, M. (2008). Assessment of genetic diversity and relationships among maize (*Zea mays* L.) Italian landraces by morphological traits and AFLP profiling. *Theoretical and Applied Genetics*. **117**: 831–42.
<http://www.fao.org/docrep/013/i1500e/United%20Republic%20Tanzania.pdf>.
- Hodgkin, T., Roviglionni, R., de Vicente, M. C. and Dudnik, N. (2001). Molecular methods in the conservation and use of plant genetic resources. *Acta Horticulturae*. **546**: 107 - 118.
- Hoisington, D., Khairallah, M., Reeves, T., Ribaut, J. M., Skovmand, B., Taba, S. and Warburton, M. L. (1999). Plant genetic resources: What can they contribute toward increased crop productivity? *Proceedings of the National Academy of Science of the United States of America*. **96**: 5937 - 5943.
- Hu, J., Mou, B. and Vick, B. (2007). Genetic diversity of 38 spinach (*Spinacia oleraceae* L.) germplasm accessions and 10 commercial hybrids by TRAP markers. *Genetic Resources and Evolution*. **54**: 1667 - 1671.
- Hugos, H. G. and Abay, F. (2013). AMMI and GGE biplot analysis of bread wheat genotypes in the northern part of Ethiopia. *Journal of Plant Breeding and Genetics*. **1**: 12 - 18.

- IBPGR. (1991). Descriptors for maize. Mexico City (Mexique) / Rome (Italy), CIMMYT / IPGRI. at:
http://archive-cpgr.cgiar.org/fileadmin/bioversity/publications/pdfs/104_Descriptors_for_maize.Descriptores_para_maiz.Descripteurs_pour_le_mais-cache=1415188810.pdf. Accessed on 24 December 2016
- Ignjatovic-Micic, D., Risitc, D., Babic, V., Anelkovic, V., Markovic, K. and Vancetovic, J. (2013). Genetic assessment of maize landraces from former Yugoslavia. *Genetika*. **45**: 405 - 417.
- Ihsan, H., Khalil, I. H., Rehman, H. and Iqbal, M. (2005). Genotypic variability for morphological traits among exotic maize hybrids. *Sarhad Journal of Agriculture*. **21**: 599 - 602.
- Isabirye, B. E. and Rwomushana, I. (2016). Current and future potential distribution of maize chlorotic mottle virus and risk of maize lethal necrosis disease in Africa. *Crop Protection*. **5**: 215 - 228.
- Jensen J., Verbakel, H., Peleman, J. and van Hintum, T. J. (2006). A note on the measurement of genetic diversity within genebank accessions of lettuce (*Lactuca sativa* L.) using AFLP markers. *Theoretical and Applied Genetics*. **112**: 554 - 561.
- Jerry, H. (2000). *Number cruncher statistical systems (NCSS)*. statistical software, Utah.
- Jones, C.J., Edwards, K.J., Castiglioni, S., Winfield, M.O., Sala, F., Van de Wiel, C, ... Karp, A. (1997). Reproducibility testing of RAPD, AFLP, and SSR markers in plants by a network of European laboratories. *Molecular Breeding*. **3**: 381 - 389.
- Kabululu, M. S., Ndunguru, J., Ndakidemi, P. and Feyissa, T. (2017). Genetic diversity of maize accessions for Maize Lethal Necrosis Disease resistance. *Indian Journal of Agricultural Research*. **51**(1): 17 - 24.
- Kagoda, F., Gidoi, R. and Isabirye, B. E. (2016). Status of maize lethal necrosis in eastern Uganda. *African Journal of Agricultural Research*. **11**: 652 - 660.
- Katinila, N., Verkuijl, H., Mwangi, W., Anandajayasekeram, P. and Moshi, A. J. (1998). Adoption of maize production technologies in southern Tanzania. Mexico, D.F.: International maize and wheat improvement center (CIMMYT), the United Republic of

- Tanzania, and the Southern Africa Centre for Cooperation in Agricultural Research (SACCAR).
- Kiruwa, F. A., Feyissa, T. and Ndakidemi, P. A. (2016). Insights of maize lethal necrotic disease: A major constraint to maize production in East Africa. *African Journal of Microbiology Research*. **10**: 271 - 279.
- Kirway, T. N., Ulotu, H. A., Lyimo, S. D., Lema, N. M., Mduruma, Z. O., Semgalawe, Z. M, ... Nyaki, A. S. (2000). The role of technology in poverty alleviation: farm household financial profitability of maize/beans intercropping technological packages.
- Kumari, J., Gadag, R. N. and Prasanna, B. M. (2005). Molecular profiling of maize (*Zea mays* L.) inbred lines using SSR markers. *Indian Journal of Genetic Plant Breeding*. **65**: 249 - 252.
- Lafitte, H. R., Edmeades, G. O. and Taba, S. (1997). Adaptive strategies identified among tropical maize landraces for nitrogen-limited environments. *Field Crops Research*. **49**: 187 - 204.
- Lavanya, G. R., Srivastava, J. and Ranade, S. A. (2008). Molecular assessment of genetic diversity in Mungbean germplasm. *Journal of Genetics*. **87**(1): 65 - 74. PubMed PMID: 18560176.
- Lee, M. (1998). Genome projects and gene pools: New germplasm for plant breeding? *Proceedings of the National Academy of Science of the United States of America*. **95**: 2001 - 2004.
- Legesse, B. W., Myburg, A. A., Pixley, K. V. and Botha, A. M. (2006). Genetic diversity of maize inbred lines revealed by SSR markers. *Hereditas*. **00**: 1 - 8.
- Legesse, B. W., Myburg, A. A., Pixley, K. V. and Botha, A. M. (2007). Genetic diversity of African maize inbred lines revealed by SSR markers. *Hereditas*. **144**: 10 - 17.
- Liao, B., Weng, P. and Zhang, J. (2010). Contrasting genetic paths to morphological and physiological evolution. *Proceedings of the National Academy of Sciences of the United States of America*. **16**: 7353 - 7358.
- Lisuma, J. B., Semoka, J. M. R. and Semu, E. (2006). Maize yield response and nutrient uptake after micronutrient application on a volcanic soil. *Agronomy Journal*. **98**: 402 - 406.

- Liu, Z., Xia, X., Yang, C. and Huang, J. (2016). Colorimetric detection of Maize chlorotic mottle virus by reverse transcription loop-mediated isothermal amplification (RT-LAMP) with hydroxynaphthol blue dye. *RSC Advances*. **6**: 73.
- Lopes, A. D., Scapim, C. A., Machado, M. F. P. S., Mangolin, C. A., Silva, T. A., Cantagali, L. B., ... Mora, F. (2015). Genetic diversity assessed by microsatellite markers in sweet corn cultivars. *Scientia Agricola*. **72**(6): 513 - 519.
- Louette, D. and Smale M. (2000). Farmers' seed selection practices and traditional maize varieties in Cuzalapa, Mexico. *Euphytica*. **113**: 25 - 41.
- Louette, D., Charrier, A. and Berthaud, J. (1997). In situ conservation of maize in Mexico: genetic diversity and maize seed management in a traditional community. *Economic Botany*. **51**: 20 - 34.
- Lu, Y., Yan, J., Guimaraes, G. T., Taba, S., Hao, Z., Gao, S, ... Xu, Y. (2009). Molecular characterization of global maize breeding germplasm based on genome-wide single nucleotide polymorphisms. *Theoretical and Applied Genetics*. **120**: 93 - 115.
- Luitel, B. P., Kim, S. G., Sung, J. S., Hur, O., Yoon, M., Rhee, J, ... Ko, H. (2016). Screening of Pumpkin (*Cucurbita spp.*) germplasm for resistance to Powdery Mildew at various stages of seedlings growth. *Research in Plant Disease*. **22**(3): 133 - 144.
- Magorokosho, C., Pixley, K. V. and Tongoona, P. (2003). Selection for drought tolerance in two tropical maize populations. *African Crop Science Journal*. **11**(3): 151 - 161.
- Mahuku, G., Lockhart, B. E., Wanjala, B., Jones, M. W., Kimunye, J. N., Stewart, L. R, ... Redinbaugh, M. G. (2015). Maize lethal necrosis (MLN), an emerging threat to maize-based food security in Sub-saharan Africa. *Phytopathology*. **105**: 956 - 965.
- Makone, S. M., Menge, D. and Basweti, E. (2014). Impact of Maize Lethal Necrosis Disease on Maize yield: a case of Kisii, Kenya. *International Journal of Agricultural Extension*. **02**: 211 - 218.
- Makumbi, D. and Wangai, A. (2013). Maize lethal necrosis (MLN) disease in Kenya and Tanzania: *Facts and actions*. CIMMYT/KARI.
http://www.fao.org/fileadmin/user_upload/drought/docs/1%20%20Maize%20Lethal%20Necrosis-%28MLN%29%20Facts%20and%20Actions.pdf. Accessed on 26 May 2016.

- Makurira, H., Savenije, H. H. G., Uhlenbrook, S., Rockstrom J. and Senzanje, A. (2007). Towards a better understanding of water partitioning processes for improved smallholder rainfed agricultural systems: A case study of Makanya catchment, Tanzania. *Physics and Chemistry of the Earth*. **32**: 1082 - 1089.
- Marçon, A., Kaeppler, S. M. and Jensen, S. G. (1997). Genetic variability among maize inbred lines for resistance to the high plains virus–wheat streak mosaic virus complex. *Plant Disease*. **81**: 195 - 198.
- Massa, A. N., Norma, C., Manrique-Carpintero., Coombs, J. J., Zarka, D. G., Boone, A. E., ... Douches, D. S. (2015). Genetic linkage mapping of economically important traits in cultivated tetraploid potato (*Solanum tuberosum* L.). *G3: Genes, Genomes, Genetics*. **5**: 2357 - 2364.
- Matsuoka, Y., Mitchell, S. E., Kresovich, S., Goodman, M. and Doebley, J. (2002). Microsatellites in maize variability, patterns of mutations, and use for evolutionary studies. *Theoretical and Applied Genetics*. **104**: 436 - 450.
- Maxted, N., Magos, B. J. and Kell, S. (2013). Resource book for preparation of national conservation plans for crop wild relatives and landraces. http://www.fao.org/fileadmin/templates/agphome/documents/PGR/PubPGR/ResourceBook/TEXT_ALL_2511.pdf. Accessed on 23 December 2016.
- Mehari, M., Alamerew S. and Lakew, B. (2015). Genotype by environment interaction for protein content of malt Barley genotypes using the additive main effect and multiplicative interaction effect model (AMMI) and genotype plus genotype by environment interaction (GGE) Biplot. *Asian Journal of Agricultural Sciences*. **9**(6): 305 - 314.
- Mercer, K. L. and Wainwright, J. D. (2007). Gene flow from transgenic maize to landraces in Mexico: An analysis. *Agriculture, Ecosystems and Environment*. **3068**: 1- 7.
- Mezzalama, M., Das, B. and Prasanna, B. M. (2015). MLN pathogen diagnosis, MLN-free seed production and safe exchange to non-endemic countries. CIMMYT brochure. Mexico, D.F.: CIMMYT. <http://repository.cimmyt.org/xmlui/bitstream/handle/10883/4284/56880.pdf?sequence=3>. Accessed on 23 December 2016.

- Mitawa, G. M. and Marandu, W. Y. F. (1996). Tanzania: Country report to the FAO international technical conference on plant genetic resources (Leipzig 1996). <http://www.fao.org/pgrfa-gpa-archive/tza/docs/tanzania.pdf>. Accessed on 26 May 2016.
- Mmbaga, M. T., Steadman, J. R. and Stavely, J. R. (1996). The use of host resistance in disease management of rust in common bean. *Integrated Pest Management Reviews*. **1**: 191 - 200.
- Molin, D., Coelho, C. J., Máximo, D. S., Ferreira, F. S., Gardingo, J. R. and Matiello, R.R. (2013). Genetic diversity in the germplasm of tropical maize landraces determined using molecular markers. *Genetics and Molecular Research*. **12**(1): 99 - 114.
- Mondini, L., Noorani, A. and Pagnotta, M. A. (2009). Assessing plant genetic diversity by molecular tools. *Diversity*. **1**: 19 - 35; doi:10.3390/d1010019.
- Moshi, A. J., Mduruma, Z. O., Lyimo, N. G., Marandu, W. F. and Akonnay, H. B. (1990). *Maize breeding for target environments in Tanzania*. In Moshi, A. J. and Ransom, J. K. (eds.), *Maize Research in Tanzania: Proceedings of the First Tanzania National Maize Research Workshop*. Dar-Es-Salaam, Tanzania: TARO. Pp. 11 - 16.
- Mwakalinga, H. A. and Massawe, W. (2007). Report on output market support. Agricultural council of Tanzania.
- N'Da, H. A., Akanvou, L., Bi, A. I. Z. and Kouakou, C. K. (2015). Phenotypic diversity of farmer's traditional maize (*Zea mays* L) varieties in Cote d'Ivoire. *Maydica*. **60**: 1 - 9.
- Nass, L. L., Pellicano I. T. and Valois, A. C. C. (1993). Utilization of genetic resources for maize and soybean breeding in Brazil. *Brazilian Journal of Genetics*. **16**: 983 - 988.
- Nei, M. and Li, W. H. (1979). Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Science of the United States of America*. **76**: 5269 - 5273.
- Nelson, S., Brewbaker, J. and Hu, J. (2011). *Maize chlorotic mottle virus*. pp. 1–6. Honolulu, HI: College of tropical agriculture and human resources, University of Hawaii at Manoa.
- Nestory, S. and Reuben, S. O. W. M. (2016). Diversity of maize landraces collected at farm level in Northern Tanzania. *International Journal of Agricultural Science Research*. **5**(1): 019-026.

- Ngugi, K., Collins, J. O. and Muchira, S. (2013). Combining, earliness, short anthesis to silking interval and yield based selection indices under intermittent water stress to select for drought tolerant maize. *Australian Journal of Crop Science*. **7**(13):2014 - 2020.
- Ngwediagi, P., Maeda, E., Kimomwe, H., Kamara, R., Massawe, S., Akonaay, H. B. and Mapunda, L. N. D. (2009). Tanzania report on the state of plant genetic resources for food and Agriculture.
<http://www.fao.org/docrep/013/i1500e/United%20Republic%20Tanzania.pdf>. Accessed on 02 February 2017.
- Nie, G., Zhang, X., Huang, L., Xu, W., Wang, J., Zhang, Y., Ma, X. and Yan, H. (2014). Genetic variability and population structure of the potential bioenergy crop *Miscanthus sinensis* (Poaceae) in southwest China on SRAP Markers. *Molecules*.12881 - 12897.
- Ntundu, W. H., Shillah, S. A., Marandu, W. Y. F. and Christiansen, J. L. (2006). Morphological diversity of bambara groundnut [*Vigna subterranea* (L.) Verdc.] landraces in Tanzania. *Genetic Resources and Crop Evolution*. **53**: 367 - 378.
- Nyaligwa, L., Hussein, S., Amelework, B. and Ghebrehiwot, H. (2015). Genetic diversity analysis of elite maize inbred lines of diverse sources using SSR markers. *Maydica*. **60**: 1 - 8.
- Obeng-Antwi, K., Craufurd, P. Q., Menkir, A., Ellis, R. H. and Sallah, P. Y. K. (2012). Phenotypic diversity in maize landraces in Ghana. *International Journal of Science and Advanced Technology*. **2**(5): 39 - 70.
- Ogwu, M. C., Osawaru, M. E. and Ahana, C. M. (2014). Challenges in conserving and utilizing plant genetic resources (PGR). *International Journal of Genetics and Molecular Biology*. **6**: 16 - 22.
- Olaoye, G. (2009). Evaluation of new generations of *Maize streak virus* (MSV) resistant varieties for grain yield, agronomic potential and adaptation to a southern guinea savanna ecology of Nigeria. *Journal of Tropical Agriculture Food, Environment and Extension*. **8** (2): 104 - 109.
- Olson, M. B., Morris, K. S. and Méndez, V. (2012). Cultivation of maize landraces by small-scale shade coffee farmers in western El Salvador. *Agricultural Systems*. **111**: 63 - 74.

- Omondi, E. C., Norton J. B. and Ashilenje, D. S. (2014). Performance of local open pollinated maize variety and a common hybrid variety under intensive small scale farming practices. *African Journal of Agricultural Research*. **9**(11): 950 - 955.
- Osawaru, M. E., Ogwu, M. C. and Aiwansoba, R. O. (2015). Hierarchical approaches to the analysis of genetic diversity in plants: A systematic overview. *University Of Mauritius Research Journal*. **21**: 92 - 127.
- Otunge, D., Muchiri, N., Wachoro, G. and Kullaya, A. (2010). Mitigating the impact of drought in Tanzania: the WEMA intervention. *Policy brief: African agricultural technology foundation (AATF) and the Tanzania commission for science and technology (COSTECH)*. <http://www.aatf-africa.org/userfiles/WEMA-TZ-policy-brief1.pdf>. Accessed on 25 December 2016.
- Pechanova, O. and Pechan, T. (2015). Maize-pathogen interactions: An ongoing combat from a proteomics perspective. *International Journal of Molecular Sciences*. **16**: 28429 - 28448.
- Pineda-Hidalgo, K. V., Méndez-Marroquín, K. P., Alvarez, E. V., Chávez-Ontiveros, J., Sánchez-Peña, P., Garzón-Tiznado, J. A., ... López-Valenzuela, J. A.(2013). Microsatellite-based genetic diversity among accessions of maize landraces from Sinaloa in México. *Hereditas*. **150**: 53 - 59.
- Poczai, P., Varga, I., Laos, M., Cseh, A., Bell, N., Valkonen, J. P. T and Hyvönen, J. (2013). Advances in plant gene targeted and functional markers: a review. *Plant Methods*. **9**: 1 - 31.
- Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S. and Rafalski, A. (1996). The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Molecular Breeding*. **2**: 225 - 238.
- Prasanna, B. M. (2010). Phenotypic and molecular diversity of maize landraces: characterization and utilization. *Indian Journal of Genetics and Plant Breeding*. **70**(4): 315 - 327.
- Prasanna, B. M. (2012). Diversity in global maize germplasm: Characterization and utilization. *Journal of Biological Science*. **37**: 843 - 855.
- Prasanna, B. M. and Hoisington, D. (2003). Molecular breeding for maize improvement: An Overview. *Indian Journal of Biotechnology*. **2**: 85 - 98.

- Pressoir, G. and Berthaud, J. (2004). Patterns of population structure in maize landraces from the Central Valleys of Oaxaca in Mexico. *Heredity*. **92**: 88 - 94.
- Ragimekula, N., Varadarajula, N. N., Mallapuram, S. P., Gangimani, G., Reddy, R. K. and Kondreddy, H. R. (2013). Marker assisted selection in disease resistance breeding. *Journal of Plant Breeding and Genetics*. **01**(02): 90 - 109.
- Rahman, H., Ali, S., Shah, S. M. A., Shah, S. S., Rahman, N., Khalil, I. A., Hussain, I. and Afzal, F. (2008). Diversity for morphological and maturity traits in maize populations from upper Dir. *Sarhad Journal of Agriculture*. **24**(3): 439 - 443.
- Rao, V. R. and Hodgkin, T. (2002). Genetic diversity and conservation and utilization of plant genetic resources. *Plant Cell, Tissue and Organ Culture*. **68**: 1 - 19.
- Rauf, S., Teixeira da Silva, J. A., Khan, A. A. and Naveed, A. (2010). Consequences of plant breeding on genetic diversity. *International Journal of Plant Breeding*. **4**(1): 1 - 21.
- Redinbaugh, M. G., José L. and Zambrano, J. L. (2014). Control of virus diseases in maize. *Advances in Virus Research*. **90**: 391 - 429.
- Reid, L. M., Xiang, K., Zhu, X., Baum, B. R. and Molnar, S. J. (2011). Genetic diversity analysis of 119 Canadian maize inbred lines based on pedigree and simple sequence repeat markers. *Canadian Journal of Plant Sciences*. **91**: 651 - 661.
- Reif, J. C., Warburton, M. L., Xia, X. C., Hoisington, D. A., Crossa, J., Taba, S., ... Melchinger, A. E. (2006). Grouping of accessions of Mexican races of maize revisited with SSR markers. *Theories of Applied Genetics*. **113**: 177 - 185.
- Ristic, D., Babic, V., Anđelkovic, V., Vancetovic, J., Drinic, S. M. and Dragana I. (2013). Genetic diversity in maize dent landraces assessed by morphological and molecular markers. *Genetika*. **45**(3): 811 - 824.
- Ristic, D., Kostadinovic, M., Kravic, N., Andjelkovic, V., Vancetovic, J. and Ignjatovic-Micic, D. (2014). Genetic Diversity in maize flint landraces assessed by morphological and SSR Markers. *Journal of International Scientific Publications: Agriculture and Food*. **2**: 206-213.
- Rodriguez, M. G., Miguel-Chavez, R. S. and Larque-Saavedra, A. (1998). Physiological aspects in Tuxpeno maize with improved drought tolerance. *Maydica*. **43**: 137 - 141.

- Romay, M. C., Millard, M. J., Glaubitz, J. C., Peiffer, J. A., Swarts, K. L., Casstevens, T. M, ... Gardner, C. A. (2013). Comprehensive genotyping of the USA national maize inbred seed bank. *Genome Biology*. **14**:R55.
- Saad, M. S. and V. Rao, V. R. (2001). Establishment and management of field genebank, a training manual. IPGRI-APO, Serdang. <https://www.cbd.int/doc/case-studies/ttcc/fieldgenebankmanual.pdf>. Accessed on 26 May 2016.
- Safavi, S. A., Ahari, A. B. and Arzanlou, M. (2010). Slow rusting resistance in 19 promising Wheat lines to yellow rust in Ardabil, Iran. *Pakistan Journal of Biological Science*. **13**(5): 240 - 244.
- Saghai-Marouf, M. A., Soliman, K. M., Jorgensen, R. A. and Allard, R. W. (1984). Ribosomal DNA spacer length polymorphisms in barley: Mendelian inheritance, chromosomal location and population dynamics. *Proceedings of the National Academy of Sciences of the United States of America*. **81**: 8014 - 8018.
- Saleh., N., Milad, S. I., El- Rouby, M. M. and Barakat, M. N. (2016). Identification of new molecular markers linked to maize stalk rot disease resistance (*Fusarium moniliforme*) in maize. *Plant Omics Journal*. **9**(1): 12 - 18.
- Salami, H. A., Adjanohoun, A., Padonou, W., Yacoubou, A. M., Aly, D., Yallou, C., Sina, H. and Baba-Moussa, L. (2015). Morphological diversity of corn's (*Zea mays* L.) local cultivar and improved varieties in Central and North of Benin. *American Journal of Plant Sciences*. **6**: 2867 - 2877.
- Salami, A. E., Adegoke S. A. O. and Adegbite, O. A. (2007). Genetic variability among maize cultivars grown in Ekiti-State, Nigeria-East. *Middle-East Journal of Scientific Research*. **2** (1): 9 - 13.
- Saleh, B. K., Kasili, R. W., Mamati, E. G., Araia, W. and Nyende, A. B. (2016). Diversity of pepper (*Capsicum* spp.) genotypes from Eritrea assessed by morphological traits. *Journal of Agriculture Science*. **8**: 156 - 168.
- Sao, K. M., Nair, S. K., Verulkar, S. B., Saxena R. R. and Nanda, H. C. (2015). Molecular profiling and genetic diversity of mungbean (*Vigna radiata* L.) genotypes using ISSR and SSR markers. *Indian Journal of Agricultural Research*. **48**: 373 - 376.

- Scheets, K. (1998). Maize chlorotic mottle Machlomovirus and wheat Streak Mosaic Rymovirus concentrations increase in the synergistic disease corn lethal Necrosis. *Virology*. **242**: 28 - 38.
- Scott, G. E. (1989). Linkage between maize dwarf mosaic virus resistance and endosperm color in maize. *Crop Science*. **29**(6): 1478 - 1480.
- Semagn, K., Beyene, Y., Babu, R., Nair, S., Gowda, M., Das, B, ... Prasanna, B. M. (2015). Quantitative trait loci mapping and molecular breeding for developing stress resilient maize for Sub-Saharan Africa. *Crop Science Society of America*. **55**: 1 - 11.
- Semagn, K., Bjørnstad, Å. and Ndjioudjop, M. N. (2006). An overview of molecular marker methods for plants. *African Journal of Biotechnology*. **5**: 2540 - 2568.
- Senior M. X., Murphy J. P., Goodman M. M. and Stuber C. W. (1998). Utility of SSRs for determining genetic similarities and relationships in maize using an agarose gel system. *Crop Science*.**38**: 1088 - 1098.
- Sharma, L., Prasanna, B. M. and Ramesh, B. (2010). Analysis of phenotypic and microsatellite-based diversity of maize landraces in India, especially from the North East Himalayan region. *Genetica*. **138**: 619 - 631.
- Sharma, R., Maloo, S. R., Joshi, A. and Choudhary, S. (2015). Assessment of genetic divergence in diverse maize (*Zea mays* L.) genotypes. *Indian Journal of Agricultural Research*. **49**: 476 - 478.
- Shiri, M. R., Choukan, R. and Aliyev, R. T. (2014). Study of genetic diversity among maize hybrids using SSR markers and morphological traits under two different irrigation conditions. *Crop Breeding Journal*. **4**(1): 65 - 72.
- Skelsey, P. and Newton A. C. (2014). Scale-dependent assessment of relative disease resistance to plant pathogens. *Agronomy*. **4**: 178 - 190.
- Smith, J. S. C., Chin, E. C. L., Shu, H., Smith, O. S., Wall, S. J., Senior, M. L, ... Ziegler, J. (1997). An evaluation of the utility of SSR loci as molecular markers in maize (*Zea mays* L.): Comparisons with data from RFLPs and pedigree. *Theoretical and Applied Genetics* **95**:163 - 173. Troyer, F. 1999. Background of U.S. hybrid corn. *Crop Science*. **39**: 601 - 626.

- Sood, S., Flint-Garcia, S., Martha, C., Willcox, M. C. and Holland, J. B. (2014). *Mining natural variation for maize improvement: Selection on phenotypes and genes*. Tuberosa, R., Graner, A., Frison, E (eds.). *Genomics of Plant Genetic Resources*. Springer, Netherlands. P 615 - 649. DOI 10.1007/978-94-007-7572-5_25.
- Sun, R., Lin, F., Huang, P. and Zheng, Y. (2016). Moderate genetic diversity and genetic differentiation in the Relict Tree *Liquidambar formosana* Hance Revealed by genic Simple Sequence Repeat Markers. *Frontiers in Plant Science*. **7**: 1 - 11.
- Taba, S., van Ginkel, M., Hoisington, D. and Poland, D. (2004). Wellhausen-Anderson Plant Genetic Resources Center: *Operations manual*. El Batan, Mexico: CIMMYT.
- Tairo, F., Mnene, E. and Kullaya, A. (2008). Morphological and agronomical characterization of Sweet potato [*Ipomoea batatas* (L.) Lam.] germplasm collection from Tanzania. *African Journal of Plant Science*. **2** (8): 77 - 85.
- Terra, T. F., Wiethölter, P., Almeida, C. C. S., Silva, S. D. A., Bered, F., Sereno, M. J. C. M. and Neto, J. F. B. (2011). Genetic variability in maize and teosinte populations estimated by microsatellites markers. *Ciência rural, Santa Maria*. **41**(2): 205 - 211.
- Tokatlidis, I. and Vlachostergios, D. (2016). Sustainable Stewardship of the Landrace Diversity. *Diversity*. **8** (29) 1 - 15.
- Toshimenla., Singh, J. and Changkija, S. (2016). Genetic divergence studies on upland rice grown in Nagaland, India. *Indian Journal of Agricultural Research*. **50**(6): 555 - 560.
- Uphoff, N., Fasoula, V., Iswandi, A., Kassam, A. and Thakur, A. K. (2015). Improving the phenotypic expression of rice genotypes: Rethinking “intensification” for production systems and selection practices for rice breeding. *The Crop Journal*. **3**: 174 - 189.
- Wan, Q., Wu, H., Fujihara, T. and Fang, S. (2004). Which genetic marker for which conservation genetics issue?. *Electrophoresis*. **25**: 2165 - 2176.
- Wang X., Chang, J., Qin, G., Zhang, S., Cheng, X. and Li, C. (2011). Analysis on yield components of elite maize variety Xundan 20 with super high yield potential. *African Journal of Agricultural Research*. **6**(24): 5490 - 5495.
- Wangai, A., Kinyua, Z. M., Otipa, M. J., Miano, D. W., Kasina, J. M., Leley, P. K. and Mwangi, T. N. (2012a). Maize (corn) lethal necrosis (MLN) disease. *KARI Information Brochure*.

- http://www.fao.org/fileadmin/user_upload/drought/docs/1%20%20Maize%20Lethal%20Necrosis%20KARI.pdf. Accessed on 26 May 2016.
- Wangai, A. W., Redinbaugh, M. G., Kinyua, Z. M., Miano, D. W., Leley, P. K., Kasina, M, ... Jeffers, D. (2012b). First report of maize chlorotic mottle virus and maize lethal necrosis in Kenya. *Plant Disease Journal*. **96**: 1582 - 1583.
<http://apsjournals.apsnet.org/doi/abs/10.1094/PDIS-06-12-0576-PDN>. Accessed on 29 March 2017.
- Warburton, M. L., Reif, J. C., Frisch, M., Bohn, M., Bedoya, C., Xia, X. C, ... Melchinger, A. E. (2008). Genetic diversity in CIMMYT nontemperate maize germplasm: landraces, open pollinated varieties, and inbred lines. *Crop Science*. **48**: 617 - 624.
- Warburton, M. L., Xianchun, X., Crossa, J., Franco, J., Melchinger, A. E., Frisch, M., Bohn, M. and Hoisington, D. (2002). Genetic characterization of CIMMYT inbred maize lines and open pollinated populations using large scale fingerprinting methods. *Crop Science*. **42**: 1832 - 1840.
- Wasala, S. K., Guleria, S. K., Sekhar, J. C., Mahajan, V., Srinivasan, K., Parsad R. and Prasanna, B. M. (2013). Analysis of yield performance and genotype × environment effects on selected maize (*Zea mays*) landrace accessions of India. *Indian Journal of Agricultural Science*. **83**(3): 287 - 293.
- Westengen, O. T., Ring, K. H., Berg, P. R. and Brysting, A. K. (2014). Modern maize varieties going local in the semi-arid zone in Tanzania. *BMC Evolutionary Biology*. **14**: 1 - 12.
- Whitt, S. R., Wilson, L. M., Tenaillon, M. I., Gaut, B. S. and Buckler, E. S. (2002). Genetic diversity and selection in the maize starch pathway. *Proceedings of the National Academy of Sciences of the United States of America*. **99**(20): 12959 - 12962.
- Wisser, R. J., Balint-Kurti, P. J. and Nelson, R. J. (2006). The genetic architecture of disease resistance in maize: A synthesis of published studies. *Phytopathology*. **96**: 120 - 129.
- Wisser., R. J., Kolkman, J. M., Patzoldt, M. E., Holland, J. B., Yu, J., Nelson, R. J. and Balint-Kurti, P. J. (2011) Multivariate analysis of maize disease resistances suggests a pleiotropic genetic basis and implicates a GST gene. *Proceedings of National Academy of Sciences*. **108**(18): 7339 - 7344.

- Xia, Z., Zhao, Z., Chen, L., Li, M., Zhou, T., Deng, C., Zhou, Q. and Fan, Z. (2016). Synergistic infection of two viruses MCMV and SCMV increases the accumulations of both MCMV and MCMV-derived siRNAs in maize. *Scientific Reports*. 6: 20520; doi: 10.1038/srep20520.
- Yan, W. (2001). GGE biplot-a Windows application for graphical analysis of multi-environment trial data and other types of two-way data. *Agronomy Journal*. **93**(5): 1111-1118.
- Yan, W. and Tinker, N. A. (2006). Biplot analysis of multi-environment trial data: Principles and applications. *Canadian Journal of Plant Science*. **86**: 623 - 645.
- Yan, W. K., Hunt, L. A., Sheng, Q. L. and Szlavnic, Z. (2000). Cultivar evaluation and mega-environment investigation based on the GGE Biplot. *Crop Science*. **40**: 597 - 605.
- Yan, W., Kang, M. S., Ma, B., Woods, S. P. and Cornelius, L. (2007). GGE biplot vs. AMMI analysis of genotype-by-environment data. *Crop Science*. **47**: 643 - 653.
- Yao, Q., Yang, K., Pan, G. and Rong T. (2007). Genetic Diversity of Maize (*Zea mays* L.) Landraces from Southwest China based on SSR data. *Journal of Genetics and Genomics*. **34**(9): 851 - 860.
- Yılmaz, M., Ozic, C. and Gok, Ý. (2012). *Principles of nucleic acid separation by agarose gel electrophoresis, gel electrophoresis - principles and basics*, Dr. Sameh Magdeldin (Ed.), ISBN: 978-953-51- 0458-2, *InTech*.
- Zarei, B., Kahrizi, D., Aboughadareh, A. P. and Sadeghi, F. (2012). Correlation and path coefficient analysis for determining interrelationships among grain yield and related characters in corn hybrids (*Zea mays* L.). *International Journal of Agriculture and Crop Sciences*. **4**(20): 1519 - 1522.
- Zeven, A. C. (1998). Landraces: A review of definitions and classifications. *Euphytica*. 104: 127 - 139.
- Zhang, S. H., Li, X. H., Wang, Z. H., George, M. L., Jeffers, D., Wang, F. G, ... Yuan, L. X. (2003). QTL mapping for resistance to SCMV in Chinese maize germplasm. *Maydica*. **48**: 307 - 312.
- Zhang, Y., Zhao, W., Li, M., Chen, H., Zhu, S. and Fan, Z. (2011). Real-time TaqMan RT-PCR for detection of maize chlorotic mottle virus in maize seeds. *Journal of Virological Methods*. **171**: 292 - 294.

APPENDICES

Appendix 1: Mean values on grain yield and yield components of all maize accessions used on yield evaluation study

| Genotype name | Yield per plant (g) | 1000 Kernel weight (g) | Days to 50% | Days to 50% | Ear Length (cm) | Ear diameter (cm) | Number of Kernels/row | Kernel rows Number |
|----------------------------|---------------------|------------------------|-------------------|-------------------|--------------------|-------------------|-----------------------|--------------------|
| | | | Tasseling | Silking | | | | |
| SITUKA1 | 116.01±12.28 | 275.51±19.47 | 64.33±2.25 | 70.89±2.67 | 16.02±0.89 | 4.28±0.17 | 31.17±2.08 | 13.05±0.33 |
| DH04 | 115.90±14.48 | 273.74±15.33 | 72.11±2.05 | 75.44±2.01 | 15.58±0.41 | 4.51±0.07 | 32.65±1.75 | 12.78±0.13 |
| PIONEER | 112.73±17.19 | 258.10±25.60 | 69.89±1.76 | 75.89±2.04 | 17.66±0.36 | 4.64±0.05 | 36.07±0.96 | 13.83±0.18 |
| TMV-1 | 108.14±10.84 | 247.27±13.52 | 73.44±1.82 | 77.78±1.88 | 16.50±0.52 | 4.36±0.07 | 31.92±1.71 | 13.92±0.50 |
| DEKALB | 104.74±9.11 | 302.73±19.86 | 66.44±2.29 | 69.89±2.39 | 16.63±0.51 | 4.58±0.09 | 33.09±1.17 | 12.42±0.13 |
| TZA2793 | 100.46±11.22 | 260.68±10.45 | 78.33±2.38 | 84.00±2.74 | 15.86±0.39 | 4.24±0.08 | 29.36±1.47 | 12.45±0.20 |
| TZA5170 | 99.80±14.55 | 285.36±26.12 | 77.44±2.46 | 83.00±2.71 | 15.14±0.54 | 4.38±0.09 | 26.77±1.55 | 11.78±0.26 |
| SITUKAM1 | 99.41±14.94 | 301.99±13.68 | 64.11±2.19 | 71.56±1.89 | 15.95±0.33 | 4.43±0.08 | 30.30±1.16 | 13.19±0.22 |
| TZA2263 | 97.54±17.01 | 260.73±20.96 | 80.33±2.69 | 86.11±3.29 | 15.35±0.69 | 4.37±0.12 | 28.86±1.73 | 12.77±0.14 |
| SC403 | 95.14±14.76 | 313.55±18.15 | 63.67±2.05 | 69.78±1.96 | 16.77±0.59 | 4.52±0.11 | 33.10±1.88 | 12.67±0.16 |
| TZA4203 | 88.19±8.48 | 250.60±16.64 | 82.00±2.79 | 89.11±2.82 | 16.20±0.55 | 3.92±0.08 | 29.06±0.90 | 12.09±0.11 |
| TZA3926 | 79.99±15.50 | 309.37±22.85 | 77.67±3.01 | 87.44±3.86 | 16.31±0.86 | 4.20±0.13 | 25.33±1.69 | 11.30±0.24 |
| TZA212 | 79.88±10.22 | 260.34±15.24 | 74.78±2.25 | 80.33±2.36 | 15.82±0.20 | 3.97±0.05 | 29.56±0.63 | 12.41±0.17 |
| TZA2731 | 78.56±14.39 | 256.75±19.40 | 79.67±3.76 | 87.00±4.27 | 14.05±1.12 | 4.21±0.16 | 27.26±2.60 | 13.21±0.50 |
| TZA3585 | 74.75±12.80 | 256.48±25.25 | 77.11±3.32 | 81.78±3.80 | 14.38±0.82 | 4.18±0.12 | 29.35±1.80 | 12.75±0.30 |
| TZA3971 | 74.20±11.58 | 274.02±9.55 | 73.78±2.48 | 79.00±3.14 | 14.99±0.54 | 4.12±0.08 | 26.25±1.07 | 12.34±0.12 |
| TZA2330 | 74.04±11.67 | 246.04±15.90 | 79.33±2.46 | 86.33±2.81 | 14.75±0.61 | 4.46±0.09 | 27.61±1.45 | 13.85±0.30 |
| CML442 | 72.93±6.68 | 212.98±16.96 | 75.89±2.10 | 81.67±2.09 | 14.87±0.34 | 3.91±0.08 | 25.22±0.79 | 13.31±0.23 |
| TZA5205 | 70.85±8.15 | 226.52±20.09 | 71.11±1.96 | 76.22±2.08 | 14.06±0.53 | 4.22±0.07 | 26.07±1.04 | 14.23±0.23 |
| TZA4020 | 70.80±6.73 | 259.01±9.81 | 73.67±2.11 | 77.67±2.32 | 14.35±0.41 | 4.06±0.06 | 28.11±1.15 | 11.69±0.22 |
| TZA2843 | 70.41±9.17 | 236.78±17.97 | 78.00±2.68 | 85.11±2.93 | 13.30±0.58 | 3.96±0.11 | 25.57±1.26 | 11.34±0.16 |
| TZA5169 | 70.07±8.79 | 236.43±14.19 | 71.44±2.52 | 77.33±2.79 | 14.66±0.96 | 3.72±0.12 | 27.89±2.66 | 11.35±0.49 |
| TZA4320 | 69.15±29.26 | 269.92±15.70 | 84.50±3.65 | 93.44±3.29 | 15.31±1.10 | 4.20±0.14 | 24.61±3.12 | 12.00±0.33 |
| TZA3206 | 69.09±8.53 | 296.91±19.14 | 70.44±2.66 | 76.11±2.79 | 14.09±0.59 | 4.06±0.08 | 26.05±1.41 | 11.43±0.25 |
| TZA599 | 66.91±9.82 | 273.42±24.13 | 83.56±2.51 | 92.11±2.49 | 16.61±0.71 | 4.21±0.12 | 27.08±2.20 | 11.24±0.29 |
| TZA3536 | 65.42±9.49 | 235.32±13.75 | 78.67±2.27 | 86.67±3.53 | 15.49±0.33 | 4.30±0.07 | 28.86±1.23 | 14.10±0.22 |
| TZA4667 | 65.02±4.37 | 279.22±11.68 | 78.22±2.95 | 85.67±3.23 | 14.28±0.44 | 4.04±0.10 | 27.23±0.63 | 11.02±0.11 |
| TZA4351 | 64.43±10.84 | 266.86±13.19 | 83.44±3.25 | 92.00±4.38 | 14.54±0.71 | 4.54±0.10 | 24.76±1.48 | 13.41±0.40 |
| TZA5129 | 63.95±8.69 | 267.91±20.21 | 83.67±2.82 | 91.22±2.23 | 15.15±0.41 | 4.28±0.09 | 25.20±1.41 | 12.74±0.19 |
| TZA3544 | 63.34±7.75 | 272.89±15.38 | 78.11±2.75 | 85.67±2.82 | 14.54±0.70 | 4.21±0.06 | 23.41±1.78 | 12.47±0.18 |
| TZA3914 | 61.16±7.01 | 271.82±15.96 | 79.33±2.95 | 82.89±3.28 | 14.83±0.78 | 4.26±0.11 | 22.73±1.59 | 12.25±0.40 |
| TZA4574 | 60.84±8.43 | 221.23±23.26 | 81.44±2.92 | 89.44±3.50 | 13.59±0.81 | 3.67±0.17 | 25.07±2.53 | 10.90±0.36 |
| CLYN261 | 60.46±10.82 | 232.81±24.03 | 84.56±2.75 | 85.89±2.78 | 13.01±0.55 | 3.65±0.13 | 18.96±1.82 | 13.01±0.21 |
| TZA1723 | 60.36±5.02 | 265.78±15.32 | 81.89±2.84 | 89.33±2.87 | 15.11±0.49 | 3.73±0.06 | 23.90±0.96 | 11.10±0.17 |
| TZA5138 | 60.13±11.55 | 213.58±11.39 | 76.78±2.02 | 83.33±2.35 | 14.18±0.48 | 4.14±0.08 | 25.37±1.66 | 13.52±0.41 |
| TZA5162 | 59.20±13.58 | 219.55±25.49 | 86.56±3.31 | 93.78±3.34 | 14.34±0.39 | 3.80±0.10 | 24.17±1.75 | 11.43±0.30 |
| TZA2264 | 58.28±11.09 | 250.20±14.40 | 78.44±2.26 | 84.11±2.73 | 14.03±0.43 | 4.37±0.11 | 26.81±1.28 | 13.05±0.27 |
| TZA3167 | 57.66±6.25 | 272.20±14.34 | 77.33±2.49 | 83.89±2.66 | 14.72±0.63 | 3.98±0.08 | 24.62±1.08 | 11.12±0.12 |
| TZA3310 | 57.39±7.29 | 238.47±13.35 | 82.72±2.52 | 90.72±2.39 | 13.83±0.36 | 3.87±0.08 | 24.57±1.95 | 11.09±0.33 |
| CLRCY034-B | 57.32±8.99 | 172.64±14.64 | 87.00±3.02 | 92.00±2.44 | 13.25±0.13 | 3.37±0.08 | 19.19±1.05 | 12.92±0.40 |
| TZA3958 | 57.06±9.71 | 296.41±25.21 | 73.44±1.50 | 79.56±1.36 | 14.72±0.63 | 4.30±0.08 | 22.29±2.39 | 11.77±0.39 |
| TZA5200 | 54.98±6.40 | 218.86±16.17 | 80.44±3.52 | 90.11±3.20 | 13.36±0.52 | 3.96±0.12 | 25.22±1.25 | 13.03±0.23 |
| TZA2338 | 53.92±7.27 | 265.96±15.04 | 84.67±3.24 | 91.56±2.90 | 13.72±0.72 | 3.78±0.10 | 22.45±2.28 | 11.73±0.32 |
| CLRCY039 | 53.25±9.46 | 181.62±14.14 | 83.56±1.82 | 87.67±1.91 | 13.42±0.44 | 3.48±0.19 | 17.20±1.14 | 12.76±0.22 |
| TZA5102 | 53.00±9.74 | 271.66±18.64 | 88.00±2.77 | 93.78±3.04 | 13.56±0.54 | 4.26±0.10 | 20.06±1.81 | 13.47±0.41 |
| TZA3614 | 52.32±8.08 | 241.38±18.31 | 84.11±2.74 | 93.00±3.57 | 13.81±0.37 | 3.83±0.07 | 22.44±1.23 | 12.50±0.22 |
| TZA3171 | 51.67±5.62 | 276.78±21.54 | 78.56±2.43 | 86.89±2.19 | 14.56±0.62 | 3.91±0.10 | 22.58±0.61 | 10.90±0.28 |
| TZA4164 | 51.39±8.76 | 239.93±21.74 | 80.72±3.28 | 90.44±3.66 | 12.69±0.80 | 3.93±0.11 | 20.55±1.82 | 11.44±0.34 |
| TZA3837 | 50.72±10.34 | 254.11±17.02 | 83.89±2.83 | 90.89±3.45 | 13.68±0.86 | 4.28±0.14 | 22.66±2.04 | 13.16±0.20 |
| TZA1745 | 50.65±8.04 | 266.43±21.68 | 85.67±3.08 | 94.56±3.30 | 15.16±0.50 | 3.83±0.12 | 23.91±1.35 | 10.36±0.23 |
| CKDHL0500 | 50.54±4.18 | 229.46±31.53 | 82.44±2.21 | 87.67±3.04 | 12.79±0.31 | 3.89±0.07 | 21.18±0.55 | 14.30±0.24 |
| TZA163 | 49.02±8.86 | 254.16±13.35 | 83.56±2.60 | 94.56±3.25 | 15.32±0.62 | 3.80±0.10 | 21.95±2.19 | 11.59±0.32 |
| TZA2904 | 48.99±5.08 | 279.01±22.62 | 69.67±4.59 | 78.56±3.10 | 15.22±0.42 | 3.67±0.11 | 22.23±1.20 | 10.84±0.29 |
| TZA608 | 48.64±7.16 | 295.61±12.53 | 79.89±3.08 | 87.67±2.58 | 15.37±0.44 | 4.33±0.07 | 27.62±1.98 | 11.58±0.38 |
| TZA5619 | 44.60±8.04 | 218.18±33.93 | 90.89±3.84 | 102.33±4.39 | 14.30±1.42 | 3.74±0.22 | 19.41±3.51 | 10.63±0.93 |
| CKDHL120552 | 43.85±6.31 | 199.49±13.85 | 80.56±2.93 | 87.56±4.10 | 11.21±0.90 | 3.29±0.21 | 18.80±3.06 | 10.72±1.11 |
| CML443 | 43.57±7.29 | 203.42±15.03 | 74.67±2.56 | 79.00±2.29 | 15.95±2.26 | 3.19±0.10 | 19.80±0.99 | 11.64±0.60 |
| TZA1753 | 41.70±5.45 | 228.84±24.48 | 78.22±3.33 | 84.89±3.52 | 15.06±0.75 | 4.24±0.12 | 22.43±1.34 | 12.65±0.30 |
| TZA3181 | 41.26±6.98 | 269.77±27.96 | 81.22±3.78 | 89.67±4.32 | 14.25±0.89 | 3.80±0.14 | 23.15±1.47 | 10.90±0.82 |
| TZA93 | 40.45±2.91 | 271.26±18.52 | 77.89±2.20 | 87.33±1.98 | 16.46±0.48 | 4.05±0.09 | 25.27±0.99 | 11.37±0.18 |
| TZA5618 | 36.69±5.03 | 219.87±24.64 | 85.78±2.99 | 94.44±3.31 | 15.41±0.67 | 4.01±0.08 | 21.50±1.33 | 11.99±0.26 |
| CKSBL10205 | 35.95±11.31 | 185.63±15.87 | 79.56±2.22 | 81.22±2.17 | 10.80±0.55 | 3.28±0.18 | 17.72±1.91 | 12.53±0.96 |
| TZA1757 | 35.72±8.24 | 284.61±24.16 | 85.11±3.46 | 93.00±3.73 | 14.59±0.64 | 4.23±0.17 | 23.35±1.31 | 11.61±0.40 |
| CML544 | 29.20±5.18 | 192.49±19.20 | 80.56±2.81 | 80.56±2.84 | 12.72±0.50 | 3.69±0.09 | 18.84±0.73 | 12.94±0.37 |
| CML440 | 29.03±4.80 | 174.27±10.31 | 72.11±1.98 | 76.22±2.52 | 11.48±0.30 | 3.36±0.11 | 17.88±0.75 | 12.82±0.37 |
| TZA1724 | 24.47±3.27 | 248.36±25.28 | 90.67±2.98 | 101.33±2.97 | 16.00±0.52 | 3.72±0.13 | 20.18±2.43 | 10.12±0.40 |
| TZMI730 | 21.14±3.85 | 204.93±21.38 | 87.78±1.88 | 91.00±2.10 | 10.32±0.73 | 3.34±0.19 | 15.44±2.45 | 12.01±0.63 |
| TZA2813 | 11.52±2.16 | 174.11±15.90 | 88.22±3.10 | 100.67±3.41 | 11.39±0.80 | 3.37±0.10 | 13.18±1.81 | 11.19±0.62 |
| Grand mean | 62.94±1.48 | 249.59±2.59 | 78.98±0.40 | 85.70±0.45 | 14.52±0.10 | 4.01±0.02 | 24.59±0.26 | 12.21±0.06 |
| Location (L) | 49.31*** | 155.64*** | 1342.61*** | 1321.48*** | 131.61*** | 129.77*** | 113.69*** | 26.93*** |
| Genotypes (G) | 6.59*** | 5.48*** | 30.65*** | 36.79*** | 6.92*** | 15.40*** | 10.86*** | 9.43*** |
| Interaction (L x G) | 1.45** | 1.59*** | 1.41** | 1.64*** | 1.12 ^{ns} | 1.50** | 1.60*** | 1.98*** |

