

2015

# Challenges in Diagnosis of Febrile illnesses in Tanzania in the Era of Declining Malaria Epidemiology

Seth, Misago

American Journal of Research Communication

---

<http://dspace.nm-aist.ac.tz/handle/123456789/477>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274699289>

# Challenges in diagnosis of febrile illnesses in Tanzania in the era of declining malaria epidemiology

Article · January 2015

CITATIONS

4

READS

73

1 author:



[Misago Seth](#)

National Institute for Medical Research (NIMR)

14 PUBLICATIONS 86 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Bioethics [View project](#)



Fish Biology [View project](#)

## Challenges in Diagnosis of Febrile illnesses in Tanzania in the Era of Declining Malaria Epidemiology

Misago Seth<sup>1,4</sup>, Daniel Mdetele<sup>2</sup>, Scott T. Phillips<sup>3</sup> and Joram Buza<sup>1</sup>

<sup>1</sup>School of Life Sciences and Bioengineering, Nelson Mandela African Institute of Science and Technology NM-AIST), P. O. Box 447, Arusha, Tanzania

<sup>2</sup>Ministry of Livestock Development & Fisheries, Directorate of Veterinary Services, Central Zone Veterinary Investigation Centre, P.O. Box 1752, Dodoma, Tanzania

<sup>3</sup>Department of Chemistry, Pennsylvania State University, University Park, PA 16802, United States

<sup>4</sup>National Institute for Medical Research (NIMR), Tanga Medical Research Centre, P. O. Box 5004 Tanga, Tanzania

Misago Seth (**Corresponding** Author): [sethm@nm-aist.ac.tz](mailto:sethm@nm-aist.ac.tz); [sethmdj2002@yahoo.com](mailto:sethmdj2002@yahoo.com)

### Abstract

Malaria and other febrile illnesses are very common especially in children in developing countries. Due to reliance on clinical algorithms for diagnosis in resource-poor settings, most febrile episodes have always been attributed to malaria. However, continuous malaria monitoring and recent improvements in malaria diagnosis have revealed a progressive decline in malaria and significant involvement of non-malarial etiologies in most febrile cases. This paper highlights the situation of malarial and non-malarial fevers, challenges facing the health sector, and possible approaches to addressing these challenges for better diagnosis of non-malarial febrile illnesses in Tanzania.

**Key words:** Non-malarial fevers, febrile illness, malaria, diagnostic challenges, Tanzania

{**Citation:** Misago Seth, Daniel Mdetele, Scott T. Phillips, Joram Buza. Challenges in diagnosis of febrile illnesses in Tanzania in the era of declining malaria epidemiology. American Journal of Research Communication, 2015, 3(5):} [www.usa-journals.com](http://www.usa-journals.com), ISSN: 2325-4076.

## Introduction

Malaria is ranked among the leading causes of morbidity and mortality in Africa and other malarial endemic regions with fever being the predominant clinical manifestation (Chandramohan *et al.* 2002; Luxemburger *et al.* 1998; Muhe *et al.* 1999; Snow *et al.* 2005; WHO 2009). Diagnosis of febrile episodes using fever as the sole criteria is somewhat cumbersome as many illnesses are accompanied by elevated axillary body temperatures. Studies conducted over 5—10 years ago have estimated the number of annual fever episodes among African children to be as high as 870 million (Gething *et al.* 2010b; Gething *et al.* 2011; Snow *et al.* 2003; Snow *et al.* 2005), a large proportion of which has been associated with malaria. However, these data have been found to overestimate malaria burden due to the lack of specificity of a purely clinical diagnosis. Studies have revealed a diminishing contribution of malaria to the febrile illnesses whose prevalence remains high (D'Acremont *et al.* 2010; Bisoffi & Buonfrate 2013; Crump *et al.* 2013; D'Acremont *et al.* 2014a). With the current decline in malaria transmission in many of the malaria endemic regions, the preponderance of febrile episodes could therefore be attributed to non-malarial febrile illnesses (NMFIs) (Acestor *et al.* 2012; Bisoffi & Buonfrate 2013; Okiro & Snow 2010). Since most fevers without any other obvious cause were in the past almost considered malarial, identification of alternative causes of fever has always been perceived as unnecessary and causes of NMFIs have remained mostly unknown. Today, the world has made a significant step in terms of malaria diagnosis. With deployment of malaria rapid diagnostic tests (mRDTs), over-diagnosis, and consequently over-treatment of malaria has been reduced and the contribution of non-malarial etiologies to febrile illnesses has now been recognized (D'Acremont *et al.* 2010). Some studies reported a proportion of non-malarial fevers ranging between 57% to above 90% (D'Acremont *et al.* 2010; Gething *et al.* 2010a). With malaria having been ruled out and without sufficient means to specifically reach

etiologic diagnosis of the febrile conditions (Petti *et al.* 2006), clinicians often find themselves in a dilemma under such circumstances. Consequently, clinical management is often driven by syndrome-based guidelines employing empiric treatment (Baltzell *et al.* 2013;WHO 2011). In the next sections, we discuss the situation of malarial and non-malarial fevers, challenges facing the health sector, and possible approaches to addressing these challenges for better diagnosis of NMFIs in Tanzania.

### **Febrile illnesses among children and adults in Tanzania**

Febrile illnesses among Tanzanian children and adults are due to a wide range of etiologies including malaria and other non-malarial causes. A recent study on malaria passive case detection (PCD) in communities of Korogwe District in Tanga, north-eastern Tanzania, has revealed that despite a significant decline in malaria prevalence in these communities, fever prevalence remains high (Rutta *et al.* 2012). In this study, the malaria positivity rate in 15,729 febrile cases was 20% in 2006, but progressively declined to below 10% in 2009 while fever cases remained unchanged (over 40% for children under five, and over 20% for children older than five (Rutta *et al.* 2012). This study did not identify the causes of non-malarial fevers, but its findings underscore the fact that even though malaria has declined, the prevalence of fever cases remains high. For example, a 2013 hospital-based study in the same district investigating causes of fever in children aged 2 to 59 months showed multiple etiology of febrile illnesses with respiratory infections occurring in more than 60% of cases (Mahende *et al.* 2014). Other syndromes include gastroenteritis (>21%), bacteremia and urinary tract infections (>21%) and malaria (8%). More than 26% of patients were found to be co-infected with more than one pathogen in this study (Mahende *et al.* 2014).

A separate study in rural and urban Tanzania investigated causes of fevers in children up to 10 years of age. As in other studies, acute respiratory infections were by far the most

common etiology (62%) followed by malaria (11%), gastroenteritis (10%) and urinary tract infection (6%). Others included typhoid fever, mucosal infections, meningitis, and some with no known causative organism (D'Acremont *et al.* 2014b). Furthermore, hospital-based studies investigating all cases admitted as severe malaria cases at Bombo (Tanga), Magunga (Korogwe), and KCMC (Kilimanjaro) hospitals revealed that a considerable proportion of admissions were malaria negative on microscopy (Reyburn *et al.* 2004; Msangeni *et al.* 2011). Specific tests on bacteremia patient samples revealed infection with *Salmonella typhi*, *Streptococcus spp*, *Staphylococcus spp*, non typhi *Salmonella*, brucellosis, rickettsiosis, leptospirosis and others (Biggs *et al.* 2011; Biggs *et al.* 2013; Biggs *et al.* 2014; Crump 2012; Crump *et al.*, 2013; D'Acremont *et al.* 2014a; D'Acremont *et al.* 2014b; Mahende *et al.* 2014; Msangeni *et al.* 2011; Prabhu *et al.* 2011). It should also be noted that not all cases were positive on blood culture, indicating that most of the patients were infected with other groups of pathogens such as viruses, protozoans, fungi as well as other bacteria that do not grow well on blood cultures and whose detection methods were beyond the scope(s) or objective(s) of these studies (Msangeni *et al.* 2011). It is apparent that febrile illnesses are still highly prevalent in Tanzania despite the reported decline in malaria. However, unlike the situation in the past, frequent causes of current febrile illnesses are non-malarial.

### **Diagnostic dilemma of non-malarial fevers**

Accurate diagnosis is a prerequisite for appropriate management of any disease or condition. Therefore, diagnostic facilities should be strengthened in order to make sure that appropriate treatment is instituted for the right disease. Failure to meet the above standards usually leads to wrong treatments, poor outcomes, and an increased risk for development of drug resistance. While the above is true, the Tanzanian health sector is still facing a challenge of poor quality diagnostic facilities, which are contributing to difficulties in disease management

(Ishengoma *et al.* 2009; Ishengoma *et al.* 2010). Some of the ongoing challenges facing health systems in Tanzania with respect to diagnosis of malarial and non-malarial fevers are discussed below.

### ***Submicroscopic malaria infection***

Microscopic examination of Giemsa-stained blood slides remains the gold standard and the most widely-used laboratory tool for diagnosis of malaria. Although mRDTs provide a more convenient option at point-of-care, and nucleic acid-based detection and identification techniques offer more sensitivity and specificity, it is microscopy that has continued to be the mainstay for malaria diagnosis in settings where resources are available. This is because, while mRDTs are rapid and user-friendly and require no specialized training, equipment, or power to read the results, they are relatively new in application and some of the original versions lacked the ability to distinguish different species of plasmodium. Most mRDTs also are qualitative, and products from different suppliers vary in sensitivity and specificity (Forney *et al.* 2003; Wongsrichanalai *et al.* 2003a; Wongsrichanalai *et al.* 2007). Nucleic acid-based detection methods are more sensitive and specific than microscopy, but require expensive and highly specialized equipment, highly trained personnel and continuous power supplies (Coleman *et al.* 2006; Santana-Morales *et al.* 2012; Nicastri *et al.* 2009). Most of these essential conditions are limited or missing in most low-resource (developing) countries where malaria is endemic.

Despite the usefulness of microscopy as a tool for malaria diagnosis, its relatively high limit of detection is a concern. The established theoretical limits of detection of malaria parasites by microscopy lie between 5—20 parasites per microliter of blood (Babiker & Schneider 2008; Bruce-Chwatt 1984; Dowling & Shute 1966; Payne 1988). However, due to a number of factors including sample collection, processing, slide reading, equipment, and malaria

incidence and prevalence, the practical limit of detection is over 100 parasites per microliter of blood (WHO 1988). This detection limit implies that microscopy may miss lower density infections, especially in endemic areas and in areas where malaria is declining. For example, it has been found that about 88% of all infections are missed in areas where malaria prevalence measured by PCR is less than 10% (Okell *et al.* 2009). This proportion of patients with parasite levels below the limit of detection by microscopy are referred to as patients with submicroscopic malaria infection (parasitemia).

Studies have demonstrated that submicroscopic parasites contribute substantially to malaria transmission (Karl *et al.* 2011; Babiker *et al.* 2008). Although the proportion of febrile infections as a result of submicroscopic parasite density has not been well characterized, the possibility has been highlighted (Snounou 1993; Cohee *et al.* 2014). The contribution of submicroscopic parasites to transmission intensity as well as febrile episodes is likely to significantly increase in areas of declining malaria burden as a result of diminishing naturally acquired immunity to malaria (Doolan *et al.* 2009). It has also been shown that, apart from acting as reservoirs of transmission, submicroscopic infections could have an impact on hemoglobin levels (Mockenhaupt *et al.* 2000; Cohee *et al.* 2014). Febrile cases due to submicroscopic parasites are expected to be few, but whenever they occur, they are likely to be missed by conventional microscopic diagnosis. Such cases will be considered to be non-malarial and patients will receive inappropriate treatment, hence minimizing their chances of timely and complete recovery, and thus further propagating not only the malaria transmission cycle, but also promoting antimicrobial resistance.

### ***The use of malaria rapid diagnostic tests (mRDTs)***

Along with microscopic examination of Giemsa-stained thick and thin blood smears, mRDTs are currently the best available and affordable alternative means of diagnosing malaria,



especially in resource-poor settings (Minja *et al.* 2012; Ishengoma *et al.* 2011; Mubi *et al.* 2011; Santana-Morales *et al.* 2012; Ochola *et al.* 2006). The use of mRDTs significantly facilitates targeted malaria treatment and reduces over-treatment with anti-malarials (Ishengoma *et al.* 2011; Mubi *et al.* 2011; Rutta *et al.* 2012). These tests are useful because they do not require sophisticated equipment or complicated procedures or training, and can be stored at room temperature, which enables their use in areas where refrigeration is lacking.

However, most mRDTs are not ideal. Their sensitivity and specificity under different manufacturing, storage, and operating conditions vary (Iqbal *et al.* 2003; Ishengoma *et al.* 2011). Varying performance implies that under certain circumstances, mRDTs may fail to accurately diagnose malaria, thus compromising the choice of treatment. Low sensitivity results in failure to detect malaria parasites in an infected patient (false negative), while low specificity results into over-diagnosis, hence overtreatment of malaria. It is paramount that mRDTs are sufficiently accurate for proper diagnosis, which is crucial in effecting optimal management of fever patients. Fortunately, in most cases their sensitivity and specificity are sufficiently high for *Plasmodium falciparum* (Pattanasin *et al.* 2003; Iqbal *et al.* 2003).

Some brands of mRDTs are designed to detect only one Plasmodium species and may be the best choice in areas where that species is the predominant cause of malaria. For example, most cases in Sub-Saharan Africa are caused by *Plasmodium falciparum* (Gething *et al.* 2011; Snow *et al.* 2005; Hay *et al.* 2009). In other parts of the world like Asia, North and South America, the Middle East, North Africa, and the South Pacific, *P. vivax* is the predominant malaria parasite (Li *et al.* 2001; Oliveira-Ferreira *et al.* 2010). However, between these extremes, other minor species such as *P. ovale*, and *P. malariae* also are present and cannot be ignored. For this reason, some of the mRDTs being developed today are capable of detecting non-falciparum malaria (Ashton *et al.* 2010). This diversity in malaria etiology poses a challenge to health care workers in developing countries, and

sometimes leads to improper diagnosis. In Tanzania for instance, mRDTs are usually provided by the Ministry of Health and Social Welfare (MoHSW) through the Medical Stores Department (MSD) (Chipwaza *et al.* 2014a). If the stock delivered to health facilities is capable of detecting *P. falciparum* only, then any non-falciparum malaria will be missed and may be treated symptomatically as a non-malarial febrile case. Therefore, an ideal mRDT would be highly sensitive, specific, and capable of detecting all common causes of malaria in a particular epidemiological and clinical setting (Mouatcho & Goldring 2013).

Despite favorable mRDT performance parameters, malaria diagnosis still poses a substantial challenge resulting from potential concomitant infection with other causes of fever (Chipwaza *et al.* 2014a). When other pathogens are involved, mere demonstration of malaria parasites may not be conclusive as to what treatment should be applied to the patient (Wongsrichanalai *et al.* 2003b; Agwu *et al.* 2009; Brouqui *et al.* 2005; Keong & Sulaiman 2006; Uneke 2008). In malaria endemic countries, infection with *Plasmodium spp* does not necessarily result in malaria, since individuals may harbor malaria parasites without becoming ill. Therefore, for every febrile case, a thorough check-up should be performed to confirm malaria or rule out non-malaria causes of fever. Unfortunately, in low-resource countries like Tanzania, healthcare facilities are ill-equipped to make this distinction.

### ***Diminishing naturally acquired immunity to malaria and its potential role in fever of malarial origin***

Naturally acquired immunity (NAI) to malaria has been associated with non-sterile protection against malaria disease in individuals who have been infected by malaria parasites (Doolan *et al.* 2009; Pinkevych *et al.* 2012; Richards *et al.* 2010; Douglas *et al.* 2011). This immunity may last for years, but can be overcome. For example, exposure to high infectious doses of

the parasites, changes in the genetics of the parasite resulting in more virulent forms, and compromised immune systems all may overcome and thus result in diminished naturally acquired immunity.

In these situations, endemic stability is likely to be disrupted. In recent years, the world has observed a substantial drop in malaria incidence and prevalence across most of the malaria-endemic regions (Mmbando *et al* 2010; O'Meara *et al.* 2010; WHO 2012; Ishengoma *et al.* 2013; Mboera *et al.*), with the exception of malaria hotspots where no decline has been observed or reported. In certain areas, this decline has been occurring for more than 15 years (Ishengoma *et al.* 2013). If withdrawal of continuous exposure to low-dose infective bites is sustained over a long period, then NAI subsequently diminishes, which will render adults susceptible to malaria (Doolan *et al.* 2009). Unpublished data from Korogwe, Tanzania show a time lag in acquisition of natural immunity to malaria in children from the age of 5 to the age of 10 years and above.

Apart from predisposing the population to rebound epidemics (Doolan *et al.* 2009; Ursing *et al.* 2014), diminished immunity will shift the endemic stability balance. The level of parasite dose to which the population is tolerant will drop in accordance with diminishing natural immunity. As a consequence, certain levels of parasites that previously resulted in asymptomatic infection could eventually cause clinical cases of fever (Doolan *et al.* 2009). Some fever cases may even be caused by parasite levels below detection limits of current rapid diagnostics and microscopy. Although greatly confounded by the non-malarial scenario, this is a possibility that needs to be investigated, and which should be addressed by developing more sensitive tests for both malarial and non-malarial febrile illnesses.

***Knowledge, attitudes and practices of the community and health workers regarding non-malaria fevers***

Although the prime clinical manifestation of malaria is fever, malaria is not the only disease that presents with fever (D'Acromont *et al.* 2014b; D'Acromont *et al.* 2014a; Mayxay *et al.* 2013; Ishengoma *et al.* 2013; Punjabi *et al.* 2012; Baltzell *et al.* 2013; Acestor *et al.* 2012; Bisoffi & Buonfrate 2013). It is therefore important to distinguish all cases of fever and correctly diagnose them to their respective etiologies or groups in order to institute appropriate treatment.

However, the current state of affairs attributes most febrile conditions as malaria. A significant proportion of community members in Tanzania and other countries do not have appropriate knowledge about febrile conditions and most believe that fever and malaria are synonymous (Baltzell *et al.* 2013; Chipwaza *et al.* 2014a; Chipwaza *et al.* 2014b). Only a small proportion were able to associate fever with true causes such as respiratory and urinary tract infections (Chipwaza *et al.* 2014a). Due to this lack of knowledge, most febrile patients or parents and guardians of febrile children expect a positive malaria test. When they are given negative results, sometimes they do not accept them. They would rather have their children given antimalarial drugs even if not diagnosed with malaria. This situation also leads to self-medication with antimalarial drugs, with or without consulting healthcare professionals (Chipwaza *et al.* 2014b; Kunda *et al.* 2007; McCombie 2002). In studies involving community members, most admitted that they did not know other causes of fever apart from malaria (Crump 2012; D'Acromont *et al.* 2014a).

Healthcare workers face a dilemma in diagnosis and management of febrile illnesses. Clinical diagnosis of fever based on symptoms and signs is not sufficient to render appropriate treatment. The use of mRDTs helps in definitive identification of positive malaria cases, but

has no use for negative cases. Non-malarial febrile illnesses due to bacteria, protozoa, fungi and viruses, for example typhoid fever, urinary tract infections (UTI), acute respiratory tract infections (ARIs), rotavirus infection, brucellosis, tick-borne relapsing fever, leptospirosis, dengue fever, and Chikungunya virus infection have been reported in Tanzania (D'Acremont *et al.* 2014b; Crump *et al.* 2013; Mahende *et al.* 2014; Prabhu *et al.* 2011; Schoonman and Swai 2009), however there is limited diagnostic capacity for non-malarial fever etiologies. Thus, healthcare workers always resort to clinical diagnosis or symptomatic treatment of febrile cases (Chipwaza *et al.* 2014a).

### **What should be the way forward?**

#### ***Create community awareness on the existence of non-malarial febrile etiologies***

It is important for patients and the general community to understand the existence of febrile etiologies other than malaria. Healthcare providers should play a leading role in this education. Patients and the general community also should be made aware that a negative malaria test of a febrile person does not necessarily mean that the method, device, or personnel doing the testing is incompetent. It also does not mean that the patient 'is not ill'. Instead, a negative test creates the need for further testing to establish the real cause of the febrile illness in the patient. Such awareness can be accomplished at different levels. First, healthcare providers could educate patients during consultation sessions after receiving test results regardless of the status of the test. Second, the health system should offer alternative diagnosis for non-malarial fever cases. It is certainly not assuring when, having all indications of febrile illness, a patient is told to go home without being treated simply because *only one* test was found to be negative. To meet this requirement, researchers and scientists must continue to explore all possible causes of febrile disease in different epidemiological settings. Third, mass education through local and regional media should be

employed to strengthen awareness and knowledge about the existence of non-malarial febrile illnesses, and the approaches for diagnosing and treating them.

***Strengthen diagnostic systems for malaria and create options for diagnosis of other causes of fever***

Most of what appears to be the non-malarial fever dilemma in Tanzania and other low-resource countries has gradually resulted from over-dependence on clinical algorithm for malaria diagnosis (Craig *et al.* 2010; Font *et al.* 2001; Petti *et al.* 2006; Reyburn *et al.* 2004). Lack of diagnostic infrastructure or resources, trained personnel, instruments such as microscopes, and accompanying reagents has always made it impossible to correctly diagnose malaria in the developing world. Slight improvements in malaria diagnostics coupled with recent declines in malaria infections revealed that not all febrile cases are due to malaria (Mouatcho & Goldring 2013; Murray *et al.* 2003; Reyburn 2010; Snounou *et al.* 1993; Van Den Ende *et al.* 2010; Wongsrichanalai & Miller 2002; Wongsrichanalai *et al.* 2007; Yukich *et al.* 2010). It is therefore proposed that improved diagnostics be extended to remote areas where such capacity is not available. Building malaria diagnostic capacity in health facilities will help discriminate true malarial and non-malarial fevers, thereby enabling accurate, effective and optimal malaria case management.

However, improved malaria diagnostics will be only one step towards success. Capacity needs to be built or strengthened to enable diagnosis of alternative causes of fever in malaria-negative febrile patients (Chappuis *et al.* 2013; Pang & Peeling 2007). To ensure that these tests are useful to those living in remote, low-resource areas in developing countries, universal tests must conform to the requirements set by the World Health Organization (WHO). WHO has developed the ASSURED criteria for ideal diagnostic tests. The test must be *Affordable* by those at risk of infection, *Sensitive* (few false-negative results), *Specific*

(few false-positive results), *User-friendly* (simple to perform by persons with little training), **Rapid and robust**, **Equipment-free** and **Deliverable** to those who need it (Urdea *et al.* 2006; Peeling *et al.* 2006). Current research is directed towards understanding the etiology of non-malarial febrile illnesses in different epidemiological settings. Most of these studies employ advanced molecular and serological techniques that are expensive, time-consuming, requiring trained personnel and in most cases, electric-powered equipment (D'Acromont *et al.* 2014b; Mahende *et al.* 2014). Although these techniques are useful in pathogen identification, most of them are impractical to be used at point-of-care. This underscores the need to develop simple, highly sensitive, and specific tests that meet the WHO ASSURED criteria in order to reach and serve resource-limited parts of the world where they are needed most. An example of such efforts employs the use of microfluidic paper-based analytical devices ( $\mu$ PADs) (Chin *et al.* 2007; Mabey *et al.* 2004; Martinez *et al.* 2010; Yager *et al.* 2006). Using paper ensures cost reduction and flexibility for patterning where photolithography (Martinez *et al.* 2008), plotting (Bruzewicz *et al.* 2008), ink etching (Abe *et al.* 2008), plasma etching (Li *et al.* 2008), cutting (Fenton *et al.* 2009), and wax printing (Carrilho *et al.* 2009; Lu *et al.* 2009) have potential as patterning methods for  $\mu$ PADs. Paper has potential as a diagnostic platform because it is readily available and inexpensive, can wick and distribute aqueous fluids, and has been in use for a long time as an analytical platform. Paper can be modified to incorporate functional groups, can easily be transported and stored, is compatible with most printing technologies, and can easily be disposed by incineration. Paper is therefore a potential starting material for making ASSURED diagnostics (Martinez *et al.* 2010; Phillips & Lewis 2014).

### ***Develop multiplex rapid diagnostic tests for alternative causes of febrile illness***

The WHO recommends rapid diagnostics that are affordable. This means that they should be made using cheap, readily available materials and simple technology so that the overall cost

of a finished product should not deter the poor from using the service. Multiplexing the tests will also help. Even in cases where simple and cheap diagnostic tests for unconventional fever etiologies are available, most of these diseases are so unknown or overlooked in the developing world that it may imply a trial-and-error approach to arrive at the right diagnosis taking one disease at a time after malaria has been excluded. This act of sequential testing may be slow, and during this time the patient's condition may deteriorate. It may also become prohibitively expensive to most people with limited resources, since the cost of each test will quickly reach an unaffordable range. These setbacks may be overcome if multiplexed, low-cost, rapid diagnostic tests that are capable of simultaneous detection of multiple pathogens are developed. Prototype  $\mu$ PADs have shown potential for multiplexing and also may have favorable performance. This type of flexibility should be utilized in designing multiplex tests for common causes of febrile illnesses in Tanzania and other low-resource countries where malarial and non-malarial febrile illnesses are common. Ongoing research provides crucial information about common causes of fever in children and adults and in different geographical and epidemiological settings. This information should be taken into consideration while planning any multiplex prototype. With all other WHO guidelines met, a diagnostic that is capable of multiplexing will greatly enhance affordability by cutting the overall cost of otherwise sequential multiple testing in non-malarial febrile patients.

## **Conclusion**

Febrile illnesses continue to cause significant problems in developing countries, especially in children under five years of age. Despite its historical dominance, malaria parasites are not the only cause of febrile conditions; the actual contribution of malaria has been diminishing over the past decade. Continuous malaria monitoring and improvement in malaria diagnosis



has uncovered significant contributions from non-malarial causes of febrile illnesses. This observation has created a new challenge since most healthcare outlets in low-resource countries, like Tanzania, are not equipped with the capacity to accurately diagnose non-malarial febrile illnesses at point-of-care. Instead, they usually rely on patient or parent history and presenting clinical signs and symptoms. Therefore, while there is still a need to develop sensitive diagnostics for malaria in order to meet new challenges resulting from low parasitemia and decline in naturally acquired immunity to malaria, there also is an urgent need to (i) create knowledge and awareness of the community regarding non-malaria fevers, and (ii) intensify efforts to develop point-of-care diagnostics for non-malarial febrile illnesses in order to improve diagnosis and management of these diseases in developing countries.

### **Acknowledgements**

We thank Daniel Minja and Deborah Sumari for their useful comments. This work is supported in part by the Tanzania Commission for Science and Technology (COSTECH) through the Nelson Mandela African Institution of Sciences and Technology (NM-AIST) Graduate Scholarship, and the National Institute of General Medical Sciences of the National Institutes of Health (R01GM105686).

### **Competing Interests**

We declare no competing interests.

**References**

1. Abe K, Suzuki K, & Citterio D (2008) Inkjet-printed microfluidic multianalyte chemical sensing paper. *Anal. Chem.* 80, p 6928-6934.
2. Acestor N, Cooksey R, Newton PN *et al.* (2012) Mapping the aetiology of non-malarial febrile illness in Southeast Asia through a systematic review--terra incognita impairing treatment policies. *PLoS. One.* 7, p e44269.
3. Agwu E, Ihongbe JC, Okogun GR, & Inyang NJ (2009) High incidence of co-infection with Malaria and Typhoid in febrile HIV infected and AIDS patients in Ekpoma, Edo State, Nigeria. *Braz. J. Microbiol.* 40, p 329-332.
4. Ashton RA, Kefyalew T, Tesfaye G *et al.* (2010) Performance of three multi-species rapid diagnostic tests for diagnosis of Plasmodium falciparum and Plasmodium vivax malaria in Oromia Regional State, Ethiopia. *Malar. J.* 9, p 297.
5. Babiker HA & Schneider P (2008) Application of molecular methods for monitoring transmission stages of malaria parasites. *Biomed. Mater.* 3, p 034007.
6. Babiker HA, Schneider P, & Reece SE (2008) Gametocytes: insights gained during a decade of molecular monitoring. *Trends Parasitol.* 24, p 525-530.
7. Baltzell K, Elfving K, Shakely D *et al.* (2013) Febrile illness management in children under five years of age: a qualitative pilot study on primary health care workers' practices in Zanzibar. *Malar. J.* 12, p 37.
8. Biggs HM, Bui DM, Galloway RL *et al.* (2011) Leptospirosis among hospitalized febrile patients in northern Tanzania. *Am. J. Trop. Med. Hyg.* 85, p 275-281.
9. Biggs HM, Hertz JT, Munishi OM *et al.* (2013) Estimating leptospirosis incidence using hospital-based surveillance and a population-based health care utilization survey in Tanzania. *PLoS. Negl. Trop. Dis.* 7, p e2589.
10. Biggs HM, Lester R, Nadjm B *et al.* (2014) Invasive Salmonella infections in areas of high and low malaria transmission intensity in Tanzania. *Clin. Infect. Dis.* 58, p 638-647.
11. Bisoffi Z & Buonfrate D (2013) When fever is not malaria. *Lancet Glob. Health* 1, p e11-e12.
12. Brouqui P, Rolain JM, Foucault C, & Raoult D (2005) Short report: Q fever and Plasmodium falciparum malaria co-infection in a patient returning from the Comoros archipelago. *Am. J. Trop. Med. Hyg.* 73, p 1028-1030.
13. Bruce-Chwatt LJ (1984) DNA probes for malaria diagnosis. *Lancet* 1, p 795.

14. Bruzewicz DA, Reches M, & Whitesides GM (2008) Low-cost printing of poly(dimethylsiloxane) barriers to define microchannels in paper. *Anal.Chem.* 80, p 3387-3392.
15. Carrilho E, Martinez AW, & Whitesides GM (2009) Understanding wax printing: a simple micropatterning process for paper-based microfluidics. *Anal.Chem.* 81, p 7091-7095.
16. Chandramohan D, Jaffar S, & Greenwood B (2002) Use of clinical algorithms for diagnosing malaria. *Trop.Med.Int.Health* 7, p 45-52.
17. Chappuis F, Alirol E, D'Acremont V, Bottieau E, & Yansouni CP (2013) Rapid diagnostic tests for non-malarial febrile illness in the tropics. *Clin.Microbiol.Infect.* 19, p 422-431.
18. Chin CD, Linder V, & Sia SK (2007) Lab-on-a-chip devices for global health: past studies and future opportunities. *Lab Chip.* 7, p 41-57.
19. Chipwaza B, Mugasa JP, Mayumana I *et al.* (2014a) Community knowledge and attitudes and health workers' practices regarding non-malaria febrile illnesses in eastern Tanzania. *PLoS.Negl.Trop.Dis.* 8, p e2896.
20. Chipwaza B, Mugasa JP, Mayumana I *et al.* (2014b) Self-medication with anti-malarials is a common practice in rural communities of Kilosa district in Tanzania despite the reported decline of malaria. *Malar.J.* 13, p 252.
21. Cohee LM, Kalilani-Phiri L, Boudova S *et al.* (2014) Submicroscopic malaria infection during pregnancy and the impact of intermittent preventive treatment. *Malar J.* 13, p 274.
22. Coleman RE, Sattabongkot J, Promstaporm S *et al.* (2006) Comparison of PCR and microscopy for the detection of asymptomatic malaria in a Plasmodium falciparum/vivax endemic area in Thailand. *Malar.J.* 5, p 121.
23. Craig JC, Williams GJ, Jones M *et al.* (2010) The accuracy of clinical symptoms and signs for the diagnosis of serious bacterial infection in young febrile children: prospective cohort study of 15 781 febrile illnesses. *BMJ* 340, p c1594.
24. Crump JA (2012) Typhoid Fever and the challenge of nonmalaria febrile illness in sub-saharan Africa. *Clin.Infect.Dis.* 54, p 1107-1109.
25. Crump JA, Morrissey AB, Nicholson WL *et al.* (2013) Etiology of severe non-malaria febrile illness in Northern Tanzania: a prospective cohort study. *PLoS.Negl.Trop.Dis.* 7, p e2324.
26. D'Acremont V, Kaiser L, & Genton B (2014a) Causes of fever in outpatient Tanzanian children. *N.Engl.J.Med.* 370, p 2243-2244.

27. D'Acromont V, Kilowoko M, Kyungu E *et al.* (2014b) Beyond malaria--causes of fever in outpatient Tanzanian children.*N.Engl.J.Med.* 370, p 809-817.
28. D'Acromont V, Lengeler C, & Genton B (2010) Reduction in the proportion of fevers associated with Plasmodium falciparum parasitaemia in Africa: a systematic review.*Malar.J.* 9, p 240.
29. Doolan DL, Dobano C, & Baird JK (2009) Acquired immunity to malaria.*Clin.Microbiol.Rev.* 22, p 13-36, Table.
30. Douglas AD, Andrews L, Draper SJ *et al.* (2011) Substantially reduced pre-patent parasite multiplication rates are associated with naturally acquired immunity to Plasmodium falciparum.*J.Infect.Dis.* 203, p 1337-1340.
31. Dowling MA & Shute GT (1966) A comparative study of thick and thin blood films in the diagnosis of scanty malaria parasitaemia.*Bull.World Health Organ* 34, p 249-267.
32. Fenton EM, Mascarenas MR, Lopez GP, & Sibbett SS (2009) Multiplex lateral-flow test strips fabricated by two-dimensional shaping.*ACS Appl.Mater.Interfaces.* 1, p 124-129.
33. Font F, Alonso GM, Nathan R *et al.* (2001) Diagnostic accuracy and case management of clinical malaria in the primary health services of a rural area in south-eastern Tanzania.*Trop.Med.Int.Health* 6, p 423-428.
34. Forney JR, Wongsrichanalai C, Magill AJ *et al.* (2003) Devices for rapid diagnosis of Malaria: evaluation of prototype assays that detect Plasmodium falciparum histidine-rich protein 2 and a Plasmodium vivax-specific antigen.*J.Clin.Microbiol.* 41, p 2358-2366.
35. Gething PW, Kirui VC, Alegana VA *et al.* (2010a) Estimating the number of paediatric fevers associated with malaria infection presenting to Africa's public health sector in 2007.*PLoS.Med.* 7, p e1000301.
36. Gething PW, Patil AP, Smith DL *et al.* (2011) A new world malaria map: Plasmodium falciparum endemicity in 2010.*Malar.J.* 10, p 378.
37. Gething PW, Smith DL, Patil AP *et al.* (2010b) Climate change and the global malaria recession.*Nature* 465, p 342-345.
38. Hay SI, Guerra CA, Gething PW *et al.* (2009) A world malaria map: Plasmodium falciparum endemicity in 2007.*PLoS.Med.* 6, p e1000048.
39. Iqbal J, Muneer A, Khalid N, & Ahmed MA (2003) Performance of the OptiMAL test for malaria diagnosis among suspected malaria patients at the rural health centers.*Am.J.Trop.Med.Hyg.* 68, p 624-628.

40. Ishengoma DR, Derua YA, Rwegoshora RT *et al.* (2010) The performance of health laboratories and the quality of malaria diagnosis in six districts of Tanzania.*Ann.Trop.Med.Parasitol.* 104, p 123-135.
41. Ishengoma DR, Rwegoshora RT, Mdira KY *et al.* (2009) Health laboratories in the Tanga region of Tanzania: the quality of diagnostic services for malaria and other communicable diseases.*Ann.Trop.Med.Parasitol.* 103, p 441-453.
42. Ishengoma DS, Francis F, Mmbando BP *et al.* (2011) Accuracy of malaria rapid diagnostic tests in community studies and their impact on treatment of malaria in an area with declining malaria burden in north-eastern Tanzania.*Malar.J.* 10, p 176.
43. Ishengoma DS, Mmbando BP, Segeja MD *et al.* (2013) Declining burden of malaria over two decades in a rural community of Muheza district, north-eastern Tanzania.*Malar.J.* 12, p 338.
44. Karl S, Gurarie D, Zimmerman PA *et al.* (2011) A sub-microscopic gametocyte reservoir can sustain malaria transmission.*PLoS.One.* 6, p e20805.
45. Keong BC & Sulaiman W (2006) Typhoid and malaria co-infection - an interesting finding in the investigation of a tropical Fever.*Malays.J.Med.Sci.* 13, p 74-75.
46. Kunda J, Fitzpatrick J, Kazwala R *et al.* (2007) Health-seeking behaviour of human brucellosis cases in rural Tanzania.*BMC.Public Health* 7, p 315.
47. Li J, Collins WE, Wirtz RA *et al.* (2001) Geographic subdivision of the range of the malaria parasite *Plasmodium vivax*.*Emerg Infect Dis.* 12, p 35-42.
48. Li X, Tian J, Nguyen T, & Shen W (2008) Paper-based microfluidic devices by plasma treatment.*Anal.Chem.* 80, p 9131-9134.
49. Lu Y, Shi W, Jiang L, Qin J, & Lin B (2009) Rapid prototyping of paper-based microfluidics with wax for low-cost, portable bioassay.*Electrophoresis* 30, p 1497-1500.
50. Luxemburger C, Nosten F, Kyle DE *et al.* (1998) Clinical features cannot predict a diagnosis of malaria or differentiate the infecting species in children living in an area of low transmission.*Trans.R.Soc.Trop.Med.Hyg.* 92, p 45-49.
51. Mabey D, Peeling RW, Ustianowski A, & Perkins MD (2004) Diagnostics for the developing world.*Nat.Rev.Microbiol.* 2, p 231-240.
52. Mahende C, Ngasala B, Lusingu J *et al.* (2014) Aetiology of acute febrile episodes in children attending korogwe district hospital in north-eastern Tanzania.*PLoS.One.* 9, p e104197.
53. Martinez AW, Phillips ST, Whitesides GM, & Carrilho E (2010) Diagnostics for the developing world: microfluidic paper-based analytical devices.*Anal.Chem.* 82, p 3-10.

54. Martinez AW, Phillips ST, Wiley BJ, Gupta M, & Whitesides GM (2008) FLASH: a rapid method for prototyping paper-based microfluidic devices. *Lab Chip*. 8, p 2146-2150.
55. Mayxay M, Castonguay-Vanier J, Chansamouth V *et al.* (2013) Causes of non-malarial fever in Laos: a prospective study. *Lancet Glob. Health* 1, p e46-e54.
56. Mboera LEG MHRSKR (2013) Towards malaria elimination and its implication for vector control, disease management and livelihoods in Tanzania. *Malar World J*. 13.
57. McCombie SC (2002) Self-treatment for malaria: the evidence and methodological issues. *Health Policy Plan*. 17, p 333-344.
58. Minja DT, Schmiegelow C, Oesterholt M *et al.* (2012) Reliability of rapid diagnostic tests in diagnosing pregnancy-associated malaria in north-eastern Tanzania. *Malar.J.* 11, p 211.
59. Mmbando BP, Vestergaard LS, Kitua AY *et al.* (2010) A progressive declining in the burden of malaria in north-eastern Tanzania. *Malar.J.* 9, p 216.
60. Mockenhaupt FP, Rong B, Till H *et al.* (2000) Submicroscopic Plasmodium falciparum infections in pregnancy in Ghana. *Trop.Med.Int.Health* 5, p 167-173.
61. Mouatcho JC & Goldring JP (2013) Malaria rapid diagnostic tests: challenges and prospects. *J.Med.Microbiol.* 62, p 1491-1505.
62. Msangeni HA, Kamugisha ML, Sembuche SH *et al.* (2011) Prospective study on severe malaria among in-patients at Bombo regional hospital, Tanga, north-eastern Tanzania. *BMC.Infect.Dis.* 11, p 256.
63. Mubi M, Janson A, Warsame M *et al.* (2011) Malaria rapid testing by community health workers is effective and safe for targeting malaria treatment: randomised cross-over trial in Tanzania. *PLoS.One.* 6, p e19753.
64. Muhe L, Oljira B, Degefu H, Enquesellasse F, & Weber MW (1999) Clinical algorithm for malaria during low and high transmission seasons. *Arch.Dis.Child* 81, p 216-220.
65. Murray CK, Bell D, Gasser RA, & Wongsrichanalai C (2003) Rapid diagnostic testing for malaria. *Trop.Med.Int.Health* 8, p 876-883.
66. Nicastrì E, Bevilacqua N, Sane SM *et al.* (2009) Accuracy of malaria diagnosis by microscopy, rapid diagnostic test, and PCR methods and evidence of antimalarial overprescription in non-severe febrile patients in two Tanzanian hospitals. *Am.J.Trop.Med.Hyg.* 80, p 712-717.

67. O'Meara WP, Mangeni JN, Steketee R, & Greenwood B (2010) Changes in the burden of malaria in sub-Saharan Africa. *Lancet Infect. Dis.* 10, p 545-555.
68. Ochola LB, Vounatsou P, Smith T, Mabaso ML, & Newton CR (2006) The reliability of diagnostic techniques in the diagnosis and management of malaria in the absence of a gold standard. *Lancet Infect. Dis.* 6, p 582-588.
69. Okell LC, Ghani AC, Lyons E, & Drakeley CJ (2009) Submicroscopic infection in Plasmodium falciparum-endemic populations: a systematic review and meta-analysis. *J. Infect. Dis.* 200, p 1509-1517.
70. Okiro EA & Snow RW (2010) The relationship between reported fever and Plasmodium falciparum infection in African children. *Malar. J.* 9, p 99.
71. Oliveira-Ferreira J, Lacerda MV, Brasil P *et al.* (2010) Malaria in Brazil: an overview. *Malar J.* 9, p 115.
72. Pang T & Peeling RW (2007) Diagnostic tests for infectious diseases in the developing world: two sides of the coin. *Trans. R. Soc. Trop. Med. Hyg.* 101, p 856-857.
73. Pattanasin S, Proux S, Chompasuk D *et al.* (2003) Evaluation of a new Plasmodium lactate dehydrogenase assay (OptiMAL-IT) for the detection of malaria. *Trans. R. Soc. Trop. Med. Hyg.* 97, p 672-674.
74. Payne D (1988) Use and limitations of light microscopy for diagnosing malaria at the primary health care level. *Bull. World Health Organ* 66, p 621-626.
75. Peeling RW, Holmes KK, Mabey D, & Ronald A (2006) Rapid tests for sexually transmitted infections (STIs): the way forward. *Sex Transm. Infect* 82 Suppl 5, p v1-v6.
76. Petti CA, Polage CR, Quinn TC, Ronald AR, & Sande MA (2006) Laboratory medicine in Africa: a barrier to effective health care. *Clin. Infect. Dis.* 42, p 377-382.
77. Phillips ST & Lewis GG (2014) The expanding role of paper in point-of-care diagnostics. *Expert. Rev. Mol. Diagn.* 14, p 123-125.
78. Pinkevych M, Petravic J, Chelimo K *et al.* (2012) The dynamics of naturally acquired immunity to Plasmodium falciparum infection. *PLoS. Comput. Biol.* 8, p e1002729.
79. Prabhu M, Nicholson WL, Roche AJ *et al.* (2011) Q fever, spotted fever group, and typhus group rickettsioses among hospitalized febrile patients in northern Tanzania. *Clin. Infect. Dis.* 53, p e8-15.
80. Punjabi NH, Taylor WR, Murphy GS *et al.* (2012) Etiology of acute, non-malaria, febrile illnesses in Jayapura, northeastern Papua, Indonesia. *Am. J. Trop. Med. Hyg.* 86, p 46-51.



81. Reyburn H (2010) New WHO guidelines for the treatment of malaria. *BMJ* 340, p c2637.
82. Reyburn H, Mbatia R, Drakeley C *et al.* (2004) Overdiagnosis of malaria in patients with severe febrile illness in Tanzania: a prospective study. *BMJ* 329, p 1212.
83. Richards JS, Stanistic DI, Fowkes FJ *et al.* (2010) Association between naturally acquired antibodies to erythrocyte-binding antigens of *Plasmodium falciparum* and protection from malaria and high-density parasitemia. *Clin. Infect. Dis.* 51, p e50-e60.
84. Rutta AS, Francis F, Mmbando BP *et al.* (2012) Using community-owned resource persons to provide early diagnosis and treatment and estimate malaria burden at community level in north-eastern Tanzania. *Malar.J.* 11, p 152.
85. Santana-Morales MA, Afonso-Lehmann RN, Quispe MA *et al.* (2012) Microscopy and molecular biology for the diagnosis and evaluation of malaria in a hospital in a rural area of Ethiopia. *Malar.J.* 11, p 199.
86. Schoonman L & Swai ES (2009) Risk factors associated with the seroprevalence of leptospirosis, amongst at-risk groups in and around Tanga city, Tanzania. *Ann. Trop. Med. Parasitol.* 103, p 711-718.
87. Snounou G (1993) Population dynamics of human malaria parasites. *Parassitologia* 35 Suppl, p 113-116.
88. Snounou G, Pinheiro L, Goncalves A *et al.* (1993) The importance of sensitive detection of malaria parasites in the human and insect hosts in epidemiological studies, as shown by the analysis of field samples from Guinea Bissau. *Trans. R. Soc. Trop. Med. Hyg.* 87, p 649-653.
89. Snow RW, Eckert E, & Teklehaimanot A (2003) Estimating the needs for artesunate-based combination therapy for malaria case-management in Africa. *Trends Parasitol.* 19, p 363-369.
90. Snow RW, Guerra CA, Noor AM, Myint HY, & Hay SI (2005) The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature* 434, p 214-217.
91. Uneke CJ (2008) Concurrent malaria and typhoid fever in the tropics: the diagnostic challenges and public health implications. *J. Vector. Borne. Dis.* 45, p 133-142.
92. Urdea M, Penny LA, Olmsted SS *et al.* (2006) Requirements for high impact diagnostics in the developing world. *Nature* 444 Suppl 1, p 73-79.
93. Ursing J, Rombo L, Rodrigues A, Aaby P, & Kofoed PE (2014) Malaria transmission in Bissau, Guinea-Bissau between 1995 and 2012: malaria resurgence did not negatively affect mortality. *PLoS. One.* 9, p e101167.



94. Van Den Ende J, Jacobs J, & Bisoffi Z (2010) Utility of point-of-care malaria rapid diagnostic tests.*Am.J.Trop.Med.Hyg.* 83, p 207.
95. Wongsrichanalai C, Arevalo I, Laoboonchai A *et al.* (2003a) Rapid diagnostic devices for malaria: field evaluation of a new prototype immunochromatographic assay for the detection of *Plasmodium falciparum* and non-falciparum *Plasmodium*.*Am.J.Trop.Med.Hyg.* 69, p 26-30.
96. Wongsrichanalai C, Barcus MJ, Muth S, Sutamihardja A, & Wernsdorfer WH (2007) A review of malaria diagnostic tools: microscopy and rapid diagnostic test (RDT).*Am.J.Trop.Med.Hyg.* 77, p 119-127.
97. Wongsrichanalai C & Miller RS (2002) Malaria rapid tests: a public health perspective.*Lancet* 359, p 1781.
98. Wongsrichanalai C, Murray CK, Gray M *et al.* (2003b) Co-infection with malaria and leptospirosis.*Am.J.Trop.Med.Hyg.* 68, p 583-585.
99. WHO (1988) Malaria diagnosis: memorandum from a WHO meeting.*Bull World Health Organ* 66, p 575-594.
100. WHO (2009). World Malaria Report 2009. Geneva (Generic Report).
101. World Health Organization. World Malaria Report. 2012. Geneva (Conference Proceeding).
102. WHO (2011) IMAI district clinician manual: hospital care for adolescents and adults: guidelines for the management of illnesses with limited resources.*World Health Organization*.
103. Yager P, Edwards T, Fu E *et al.* (2006) Microfluidic diagnostic technologies for global public health.*Nature* 442, p 412-418.
104. Yukich J, D'Acremont V, Kahama J, Swai N, & Lengeler C (2010) Cost savings with rapid diagnostic tests for malaria in low-transmission areas: evidence from Dar es Salaam, Tanzania.*Am.J.Trop.Med.Hyg.* 83, p 61-68.