

2014

# The impact of malignant catarrhal fever virus challenge on physiological stress in cattle

Daudi, Mark

American Journal of Research Communication

---

<http://dspace.nm-aist.ac.tz/handle/123456789/498>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

## The impact of malignant catarrhal fever virus challenge on physiological stress in cattle

Mark Daudi<sup>1</sup>, Rachel Santymere<sup>2</sup>, Felix Lankester<sup>3</sup> and Joram Buza<sup>1</sup>

1. Nelson Mandela African Institution of Science and Technology
2. Lincoln Park Zoo Park, 2001 N Clark St, Chicago, IL 60614, USA
3. Paul G. Allen School for Global Animal Health, PO Box 647090, Washington State University, Pullman WA 99164-7090

**Corresponding author:** Joram Buza; e-mail: [joram.buza@nm-aist.ac.tz](mailto:joram.buza@nm-aist.ac.tz)

### Abstract

There is a lot of information in literature on animals succumbing to disease(s) after exposure to stressors. But little information is available on what happens to corticosterone hormones when animals are exposed to a pathogen as a stressor. In this study, the effect of Malignant catarrhal fever viral challenge on the kinetics of corticosterone hormone in vaccinated and un-vaccinated cattle was investigated. Animals were randomly allocated into five (5) treatment groups each containing eight (8) animals. The animals were vaccinated against Malignant catarrhal fever using the attenuated Alcelaphine herpes virus AHV1.0 vaccine in combination with either flagellin or emulsigen adjuvants as follows; Group 1 (vaccine +emulsigen), Group 2 (vaccine+flagellin), Group 3 (vaccine+emulsigen+flagellin), Group 4 (emulsigen only) and Group 5 (flagellin only). All animals were challenged with Malignant catarrhal fever virus (virulent C-500 strain AHV-1 virus)  $tcid_{50}$  on day 77 after the primary vaccination. Stress response was determined by measuring fecal corticosterone levels using ELISA. Survival from viral challenge was 75% (Group 1), 50% (Group 2), 37.5% (Group 3), 50% (Group 4) and 12.5% (Group 5). The differences between survival curves was significant ( $p=0.0182$ ).

The 2-way ANOVA was used to determine whether the different treatment regime (vaccine, adjuvant, viral challenge combinations) resulted in different corticosterone responses. Results showed that while there was significant time effect ( $p < 0.001$ ), the group and group $\times$ time interaction did not have significant effect. Initially in all treatment groups, the mean

corticosterone concentrations decreased progressively from baseline to lowest levels on day 56 and then increased sharply to peak levels on day 77 just before viral challenge. However the mean peak levels were significantly higher than baseline levels only in Group 2 (vaccine+flagellin) and Group 5 (flagellin alone). After challenge with live virus, the mean corticosterone levels decreased progressively in all groups from peak levels on day 77 to lowest levels on day 133 and then stabilized. The decrease in corticosterone levels after challenge was however significant in Group 5 only. Compared to peak levels, corticosterone concentrations in group 5 were significantly lower on Day 133 ( $p<0.05$ ), Day 147 ( $p<0.001$ ) and Day 161 ( $p<0.001$ ).

To determine whether there was difference in corticosterone levels between animals that died from the disease and those that survived, the unpaired t-test was used to determine difference. Mean terminal corticosterone hormone metabolites in dead animals ( $43.36 \pm 3.796$ ) was significantly higher ( $p< 0.05$ ) than in survivors ( $29.32 \pm 4.548$ ) indicating higher stress response in animals with severe clinical signs.

Results from this study demonstrate that Malignant catarrhal fever virus challenge in cattle decreases corticosterone levels but in general corticosterone levels are higher in animals with severe clinical signs of the disease.

{**Citation:** Mark Daudi, Rachel Santymere, Felix Lankester, Joram Buza. The impact of malignant catarrhal fever virus challenge on physiological stress in cattle. American Journal of Research Communication, 2014, 2(3):13-29} [www.usa-journals.com](http://www.usa-journals.com), ISSN: 2325-4076.

## Introduction

Animals, both free ranging and domesticated endure many stressors in their day to day lives. For example, those in the wild experience food shortages, dwell in areas where predator or parasite densities are high, engage in conflicts with neighbors or group members, and face fluctuations in food and water availability and temperature fluctuation (Millspaugh *et al.*, 2003; Preisser *et al.*,

2005; Bayazit, 2009). Domesticated animals also encounter stressors ranging from restricted movements, forced proximity to humans, absence of retreat space, routine husbandry, restricted feeding and foraging opportunities and many others (Campbell, 2001; Koch *et al.*, 2009; Pearce and Paterson, 1993; Morgan and Tromborg, 2007).

Vertebrate animals have evolved mechanisms for coping with stress which involves the adrenal gland-produced glucocorticoids hormones (Sapolsky *et al.*, 2000; Harvey *et al.*, 1984). These hormones prepare the animal to a state of emergency by increasing synthesis of glucose from various storage sources (gluconeogenesis) but also suppress the immune functions, which may predispose to succumbing to diseases (Wingfield *et al.*, 1998; Cyr, 2008).

Of late, fecal glucocorticoids metabolites are becoming the preferred target analytes for stress in different animals. It is a minimally invasive method compared to the blood sampling methods (Cook, 2012). The procedure has been applied in many animal species such as sheep (Coburn, 2010), white and black rhinoceros (Turner Jr *et al.*, 2002), elephants (Foley *et al.*, 2001), farm animals (Palme, 2012) and ground squirrels (Bosson, 2009).

There are many publications on how stress predisposes to diseases but little information is available on how infectious agents or disease affect the stress response. It is well established that infectious agents or parasites can be stressors (Muehlenbein, 2006; Frandsen, 1987; Sapolsky *et al.*, 2000; Cook, 2012; Agarwal and Marshall, 2001; Peterson *et al.*, 1991; Martin, 2009).

This study attempted to investigate the level of stress induced by inoculation of cattle with a viral infectious agent and also test whether animals that are protected by vaccination will have diminished stress response.

## **Materials and Methods**

### ***Experimental Animals***

The study recruited 40 Tanzania shorthorn zebu purchased from cattle owners in local markets around Simanjiro district, Arusha, Tanzania. The experimental animal ages ranged from 24-30 months. The Animals were taken to a Holding Area located in Emboret village close to the

Simanjiro plain. Each animal was ear-tagged with unique identification number at the start of the trial. The animals received regular health checks and were administered with Ivermectin, a broad-spectrum anti-parasitic agent to treat gastrointestinal round worms, lungworm, grub and suckling lice. Also cattle were vaccinated against East Coast Fever one week before the trial.

### *Experimental animal groups*

The animals were randomly assigned into five treatment groups, each containing eight animals. Depending on the group, the animals were either vaccinated with Malignant catarrh fever vaccine in combination with adjuvant or adjuvant only as described in Table 1.

**Table 1: Experimental animals groups**

| Group | Treatment (inoculum) |                     |
|-------|----------------------|---------------------|
|       | Vaccine              | Adjuvant            |
| 1     | +                    | Emulsigen           |
| 2     | +                    | Flagellin           |
| 3     | +                    | Emulsigen+Flagellin |
| 4     | -                    | Emulsigen           |
| 5     | -                    | Flagellin           |

### *Vaccination and challenge*

Vaccine and adjuvant were prepared and administered according to the treatment groups. The propagated Malignant catarrhal fever virus C500 strain passaged more than 1000 times in bovine turbinate cells (Haig, 2008) was used as vaccine. The Emulsigen adjuvant used in this study was an licensed adjuvant from MVP laboratories (Omaha, USA) while the Flagellin used

in this study was extracted from bacterial flagellum as previously described (Lee *et al.*, 2006). For primary vaccination on day 0, cattle were arranged according to their respective treatment groups, confined in a crush then injected with 1.2 mL of inoculum intramuscular around the neck region. Booster vaccination was administered in the same way as prime vaccination on day 28. Animals in all groups were challenged with live Malignant catarrhal fever virus virus on day 77 by intranasal inoculation with 10 mLs virus of suspension containing approximately  $10^4$  TCID<sub>50</sub>/mL.

### ***Fecal sample collection***

Fecal samples were collected just before vaccination (day 0), and after vaccination on days 28, 42, 56, 77, 105, 119, 133, 147 and 161 for determination of fecal corticosterone metabolite levels. Fresh fecal sample were collected from each cattle at approximately 9:00 am to avoid diurnal corticosterone sample variation (Touma and Palme, 2005). Cattle were confined in cattle chute then samples were directly collected from rectum after recto-stimulation. Each sample was stored in small plastic bag which was correctly identified with respect to animal ID and date. The fecal samples were stored in cool box with ice blocks then shipped to Veterinary Investigation Center Laboratory in Arusha town and frozen at -20 °C until use.

### ***Monitoring of animals***

The health status of the cattle was monitored through daily health checks and rectal temperatures were recorded every other day. Any cattle showing signs of disease (fever, depression, lack of appetite, ocular lesions, nasal discharge etc.) were monitored and the severity of clinical signs scored based on the Malignant catarrh fever clinical sign score sheet as previously described (Russell *et al.*, 2012). Animals with very severe clinical signs (score above 6), were judged not fit to continue with the trial and were therefore euthanized. Terminal fecal sample was collected before euthanization.

### ***Hormone Extraction***

The corticosterone metabolites were extracted from fecal samples as previously described (Santymire *et al.*, 2012; Loeding *et al.*, 2011). Samples were defrosted and weighted using weight balance. To each 0.5g of sample was added 5 mLs of 90% ethanol. The mixture was homogenized for one (1) minute and centrifuged at 210( $\times$ g) for 20 minutes. The liquid supernatant was transferred into a clean plastic tube from which one (1) mL sample was removed and transferred to a test tube and left to evaporate and dry for one week. Then the dry sample was capped and stored at room temperature ready for corticosterone quantification.

### ***Corticosterone Enzyme Immunoassay***

The corticosterone concentration in samples was quantified using the DetectX® Corticosterone Immunoassay kit (Arborassay Company, USA) according to manufacturer's instructions. The kit comes with plate wells that are already pre-coated with donkey anti sheep IgG. Briefly, dried fecal samples were reconstituted by adding one (1) mL of assay buffer solution. Then 50 $\mu$ L of the sample and standard were added into respective wells in the plate. The DetectX® Corticosterone Conjugate (25 $\mu$ L) was added to wells and plates tapped to ensure adequate mixing. The plates were then covered with plate sealer and shaken for one hour at room temperature to ensure maximum binding.

The plates were then washed four times, by adding 300  $\mu$ L of wash buffer and then taped on dry clean absorbent towels. Then 100  $\mu$ L of TMB substrate were added to each well using micro channel pipette and incubated at room temperature for 30 minutes without shaking. Finally 50  $\mu$ L of stop solution were added to each well. The optical density generated from each wells in the plate were measured by micro plate reader at the wavelength of 450 nm.

Based on the manufacturer's instructions, before samples were tested, validation of assay was carried out. This involved serial dilutions of fecal sample pool along with known concentration of standard dilutions of corticosterone to obtain respective optical density readings.

### *Data analysis*

Datasets were analyzed using Graphpad prism version 6 (La Jolla, Ca, USA). Comparison of temperature responses between groups was done using One-way ANOVA with repeated measures. The Two-way ANOVA with repeated measures was used to determine differences between treatment groups over the experimental period. The student t-test was used to compare differences in corticosterone concentrations between animals that survived the challenge and those who died from challenge. Differences of  $p < 0.05$  were deemed significant.

## **Results**


### *Clinical outcome following vaccination and viral challenge*

After viral challenge, clinical signs of Malignant catarrhal fever infection were first detectable around day 97 (20 days after challenge). The average incubation period was 30 days (Group 1), 51 days (Group 2), 51 days (Group 3), 47 days (Group 4) and 30 days (Group 5).

Animals in all groups developed clinical signs of Malignant catarrh fever but the signs were more severe in Group 5 and least severe in Group 1. The clinical signs included serous nasal discharge, ocular discharges, swelling of pre-scapular lymph nodes, bilateral corneal opacity, shivering, photophobia, aggression in some animals, incoordination, tremor, head pressing, body shivering, and convulsion.

Rectal temperature differences were apparent between groups after challenge with Malignant catarrhal fever. Temperatures were highest in Group 5 and lowest in Group 1 and the differences between groups was significant (Fig. 1).






**Figure 1: Mean temperatures between day 109 and day119. (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ).**

### *Effect of vaccination regimes on protection from MCF challenge*

The proportion of surviving animals was 75% (Group 1), 50% (Group 2), 37.5% (Group 3), 50% (Group 4) and 12.5% (Group 5). The difference between groups was significant ( $p = 0.0182$ ) in survival from challenge were observed

### *Parallelism of the fecal sample and standard*

The biochemical validation to analyze cattle corticosterone metabolites concentration using enzyme immunoassay was carried out. Serial dilutions of fecal sample pools yielded the displacement curve which was parallel to that of corticosterone standard curves with the linear regression  $R^2 = 0.999$  (Fig. 2).




**Figure 2: Parallelism of serial dilutions of fecal corticosterone in samples and corticosterone standard concentrations.**

### *Effect of vaccination regimes and challenge on concentration of fecal corticosterone*

The two-way ANOVA with repeated measures was used to determine whether the different treatment regimens (vaccine, adjuvant, viral challenge combinations) induced different corticosterone responses. Results showed that while there was significant time effect ( $p < 0.001$ ), the group and group $\times$ time interaction did not have significant effect (Figure 3). Initially in all treatment groups, the mean corticosterone concentrations decreased progressively from baseline to lowest levels on 56 day and then increased sharply to peak on 77 day, just before challenge. However the mean peak levels were significantly higher than baseline levels only in two groups; Group 2 (vaccine+flagellin) and Group 5 (flagellin alone).


After challenge with live virus, the mean corticosterone levels decreased progressively in all groups from peak levels on day 77 to lowest levels on day 133 and then stabilized. The decrease in corticosterone levels after challenge was however significant in group 5 only. Compared to peak levels (Day 77), corticosterone concentrations were significantly lower in group 5 on Day 133 ( $p<0.05$ ), Day 147 ( $p<0.01$ ) and Day 161 ( $p<0.01$ ).



**Figure 3: Changes in mean corticosterone concentrations (ng/g wet feces) in the five groups during the course of study.** BLD-Base line data, PV-Primary vaccination, BV-Booster vaccination and VC-Viral challenge.

### *Terminal corticosterone levels in animals that died from viral challenge*

The unpaired t- test was used to determine whether there were differences in corticosterone levels between animal which died from the disease and those that survived. Mean corticosterone hormone metabolites in dead  $43.36 \pm 3.796$  ( $n=20$ ) was significantly higher ( $p< 0.05$ ) than for survivors  $29.32 \pm 4.548$  ( $n=18$ ).



**Figure 4: Corticosterone concentrations and animals that survived and those that died from malignant catarrhal fever challenge. (\* p<0.05).**

## Discussion

This study attempted to investigate the level of stress induced by Malignant catarrhal fever in cattle via the quantification of fecal corticosterone metabolites. Animals were vaccinated with a Malignant catarrhal fever vaccine in combination with either emulsigen or Flegellin adjuvant and subsequently challenged with live Malignant catarrhal fever virus. Clinical signs characteristic of Malignant catarrhal fever (Cleaveland, 2001; Li *et al.*, 2011; Russell *et al.*, 2009; Schultheiss *et al.*, 2000) developed in all groups though with differing severity. Signs were more severe in unvaccinated control group (Group 5) from which 7/8 animals died and least severe in Group 1 (Vaccine +emulsigen) from which only 2/8 animals died. Considering that there is currently no

known vaccine for Malignant catarrhal fever, results from this study are very promising for a vaccine formulated with emulsigen adjuvant which protected 75% of animals from deaths.

Results from this study shows that the mean corticosterone levels in all groups increased sharply from day 56 and peaked on day 77 just before challenge. After challenge, the mean corticosterone levels progressively decreased to baseline levels between day 119 and day 133. It was interesting to note that although the increase in corticosterone levels occurred in all groups (treated and untreated), significant increase was only noted in Group 2 (vaccine +flagellin) and Group 5 (flagellin only) indicating that flagellin adjuvant played a role in the cortosterone increase. However, flagellin was last injected in animals during a booster vaccination which was done on day 28. Since peak levels were detected on day 77, which is 49 days after booster vaccination, it creates some doubts whether the increase was due to the adjuvant or the vaccine. Normally detectable increase in levels of fecal corticosterone appears in a few hours to few days after induction of stress (Santymire *et al.*, 2012; Möstl and Palme, 2002; Millspaugh and Washburn, 2004) which is contrary to what was seen in this study. An alternative explanation for this rise in cortisone is some environmental factors such water shortage, shortage of grass or even presence of predators as was observed in other studies (Laver *et al.*, 2012; Palme, 2005; Scheuerlein *et al.*, 2001; Boonstra, 1998; Millspaugh and Washburn, 2004). This latter explanation is more feasible because animals in all groups experienced increase in corticosterone levels regardless of the treatment administered.

It was observed that the cortisterone levels started decreasing progressively after animals were challenged with live virus and clinical signs associated with Malignant catarrh fever developed. This is in contrast to our hypothesis and what was observed in other studies involving viral challenge to animals where corticosterone levels increased soon after challenge (Sapolsky *et al.*, 2000; Webster and Sternberg, 2004). This may be due to different reasons; firstly, there is paucity of information on how cortisterone levels react to viral challenges in cattle in view of the fact that to our knowledge no previous Malignant catarrhal fever viral challenge experiment has been reported in this species. Secondly, it may be that the pathology associated with Malignant catarrhal fever interferes with the stress sensing or stress response mechanisms in cattle. Malignant catarrhal fever is associated with pathology in various tissue and organs including lymphocyte infiltration in various non-lymphoid organs and this may interfere with a number of

organs including the adrenal gland which secretes the corticosterone hormones. Alternatively since it is known that elevated corticosterone suppresses immune responses, the decrease in corticosterone levels may be due to an unknown inbuilt mechanism to boost the immune response.

We noted however that those animals that exhibited severe clinical signs and subsequently euthanized had significantly higher terminal corticosterone levels compared with those with mild clinical signs who survived the challenge. This is in agreement with our hypothesis and other studies involving viral challenge in different animal species (Webster and Sternberg, 2004). This also agrees with our argument that higher corticosterone levels may have compromised the immune responses in animals with severe clinical signs and left them susceptible to progression of disease.

## **Conclusion**

This study demonstrated for the first time the stress responses in cattle following challenge with virulent Malignant catarrh fever virus. The results and methodology used will set the trend for further studies of stress responses in cattle.

## **Acknowledgements**

The authors would like to thank the Tanzania Commission for Science and Technology (COSTECH) who sponsored the study through the Nelson Mandela African Institution of Sciences and Technology (NM-AIST). Sincere gratitude to Prof. Sarah Cleveland of the University of Glasgow, UK for allowing the use of her experimental animals in this study.

**References**

- Agarwal, S. and Marshall, G. (2001). Stress effects on immunity and its application to clinical immunology. *Clinical & Experimental Allergy*. 31(1): 25-31.
- Bayazit, V. (2009). Evaluation of cortisol and stress in captive animals. *Australian Journal of Basic and Applied Sciences*. 3(2): 1022-1031.
- Boonstra, R., Hik, D., Singleton, G.R., and Tinnikov, A. (1998). The impact of predator-induced stress on the snowshoe hare cycle. *Ecological Monographs*. 68(3): 371-394.
- Bosson, C. O., Palme, R., and Boonstra, R. (2009). Assessment of the stress response in Columbian ground squirrels: laboratory and field validation of an enzyme immunoassay for fecal cortisol metabolites. *Physiological and Biochemical Zoology*. 82(3): 291-301.
- Campbell, T., Meagher, M.W., Sieve, A., Scott, B., Storts, R., Welsh, T.H., and Welsh, C.J.R. (2001). The effects of restraint stress on the neuropathogenesis of Theiler's virus infection: I. Acute disease. *Brain, behavior, and immunity*. 15(3): 235-254.
- Capitanio, J. P.; Mendoza, S. P.; Lerche, N. W. and Mason, W. A. (1998). Social stress results in altered glucocorticoid regulation and shorter survival in simian acquired immune deficiency syndrome. *Proceedings of the National Academy of Sciences*. 95(8): 4714-4719.
- Cook, N. J. (2012). Review: Minimally invasive sampling media and the measurement of corticosteroids as biomarkers of stress in animals. *Canadian Journal of Animal Science*. 92(3): 227-259.
- Cyr, N. E., and Romero, L .M. (2008). Fecal glucocorticoid metabolites of experimentally stressed captive and free-living starlings: implications for conservation research. *General and comparative endocrinology*. 158(1): 20-28.
- Foley, C.; Papageorge, S. andWasser, S. (2001). Noninvasive Stress and Reproductive Measures of Social and Ecological Pressures in Free-Ranging African Elephants. *Conservation Biology*. 15(4): 1134-1142.
- Frandsen, J. C. (1987). Parasites as stressors: Plasma cortisol responses of goats infected with the stomach worm *Haemonchus contortus* to exogenous corticotropin (ACTH). *Veterinary Parasitology*. 23(1): 43-49.
- Haig, D. M., Grant, D., Deane, D., Campbell, I., Thomson, J., Jepson, C., Buxton, D., and Russell, G.C. (2008). An immunisation strategy for the protection of cattle against alcelaphine herpesvirus-1-induced malignant catarrhal fever. *Vaccine*. 26(35): 4461-4468.

- Harvey, S.; Phillips, J.; Rees, A. and Hall, T. (1984). Stress and adrenal function. *Journal of Experimental Zoology*. 232(3): 633-645.
- Koch, M.; Möstl, E.; Steinmetz, H. W.; Clauss, M.; Masello, J. F. and Quillfeldt, P. (2009). Non-invasive measurement of faecal glucocorticoid metabolites in Upland Geese *Chloephaga picta*. *Polar Biology*. 32(2): 281-285.
- Laver, P.; Ganswindt, A.; Ganswindt, S. and Alexander, K. (2012). Non-invasive monitoring of glucocorticoid metabolites in banded mongooses *Mungos mungo* in response to physiological and biological challenges. *General and comparative endocrinology*.
- Lee, S. E.; Kim, S. Y.; Jeong, B. C.; Kim, Y. R.; Bae, S. J.; Ahn, O. S.; Lee, J.-J.; Song, H.-C.; Kim, J. M. and Choy, H. E. (2006). A bacterial flagellin, *Vibrio vulnificus* FlaB, has a strong mucosal adjuvant activity to induce protective immunity. *Infection and immunity*. 74(1): 694-702.
- Li, H.; Cunha, C. W. and Taus, N. S. (2011). Malignant catarrhal fever: understanding molecular diagnostics in context of epidemiology. *International journal of molecular sciences*. 12(10): 6881-6893.
- Loeding, E.; Thomas, J.; Bernier, D. and Santymire, R. (2011). Using fecal hormonal and behavioral analyses to evaluate the introduction of two sable antelope at Lincoln Park Zoo. *Journal of Applied Animal Welfare Science*. 14(3): 220-246.
- Martin, L. B. (2009). Stress and immunity in wild vertebrates: timing is everything. *General and comparative endocrinology*. 163(1): 70-76.
- Millspaugh, J. J. and Washburn, B. E. (2004). Use of fecal glucocorticoid metabolite measures in conservation biology research: considerations for application and interpretation. *General and comparative endocrinology*. 138(3): 189-199.
- Millspaugh, J. J.; Washburn, B. E.; Milanick, M. A.; Slotow, R. and van Dyk, G. (2003). Effects of heat and chemical treatments on fecal glucocorticoid measurements: implications for sample transport. *Wildlife Society Bulletin*: 399-406.
- Morgan, K. N. and Tromborg, C. T. (2007). Sources of stress in captivity. *Applied Animal Behaviour Science*. 102(3): 262-302.
- Möstl, E. and Palme, R. (2002). Hormones as indicators of stress. *Domestic Animal Endocrinology*. 23(1): 67-74.
- Muehlenbein, M. P. (2006). Intestinal parasite infections and fecal steroid levels in wild chimpanzees. *American journal of physical anthropology*. 130(4): 546-550.
- Palme, R. (2005). Measuring fecal steroids: guidelines for practical application. *Annals of the New York Academy of Sciences*. 1046(1): 75-80.



- Palme, R. (2012). Monitoring stress hormone metabolites as a useful, non-invasive tool for welfare assessment in farm animals. *Animal Welfare-The UFAW Journal*. 21(3): 331.
- Pearce, G. and Paterson, A. (1993). The effect of space restriction and provision of toys during rearing on the behaviour, productivity and physiology of male pigs. *Applied Animal Behaviour Science*. 36(1): 11-28.
- Peterson, P. K.; Chao, C. C.; Molitor, T.; Murtaugh, M.; Strgar, F. and Sharp, B. M. (1991). Stress and pathogenesis of infectious disease. *Review of Infectious Diseases*. 13(4): 710-720.
- Preisser, E. L.; Bolnick, D. I. and Benard, M. F. (2005). Scared to death? The effects of intimidation and consumption in predator-prey interactions. *Ecology*. 86(2): 501-509.
- Reeder, D. M. and Kramer, K. M. (2005). Stress in free-ranging mammals: integrating physiology, ecology, and natural history. *Journal of Mammalogy*. 86(2): 225-235.
- Russell, G. C.; Benavides, J.; Grant, D.; Todd, H.; Deane, D.; Percival, A.; Thomson, J.; Connelly, M. and Haig, D. M. (2012). Duration of protective immunity and antibody responses in cattle immunised against alcelaphine herpesvirus-1-induced malignant catarrhal fever. *Veterinary research*. 43(1): 1-11.
- Russell, G. C.; Stewart, J. P. and Haig, D. M. (2009). Malignant catarrhal fever: a review. *The Veterinary Journal*. 179(3): 324-335.
- Santymire, R. M.; Freeman, E. W.; Lonsdorf, E. V.; Heintz, M. R. and Armstrong, D. M. (2012). Using ACTH challenges to validate techniques for adrenocortical activity analysis in various African wildlife species. *International Journal of Animal and Veterinary Advances*. 4.
- Sapolsky, R. M.; Romero, L. M. and Munck, A. U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine reviews*. 21(1): 55-89.
- Scheuerlein, A.; Van't Hof, T. and Gwinner, E. (2001). Predators as stressors? Physiological and reproductive consequences of predation risk in tropical stonechats (*Saxicola torquata axillaris*). *Proceedings of the Royal Society of London. Series B: Biological Sciences*. 268(1476): 1575-1582.
- Touma, C. and Palme, R. (2005). Measuring fecal glucocorticoid metabolites in mammals and birds: the importance of validation. *Annals of the New York Academy of Sciences*. 1046(1): 54-74.
- Turner Jr, J. W.; Tolson, P. and Hamad, N. (2002). Remote assessment of stress in white rhinoceros (*Ceratotherium simum*) and black rhinoceros (*Diceros bicornis*) by

measurement of adrenal steroids in feces. *Journal of Zoo and Wildlife Medicine*. 33(3): 214-221.

Webster, J. I. and Sternberg, E. M. (2004). Role of the hypothalamic-pituitary-adrenal axis, glucocorticoids and glucocorticoid receptors in toxic sequelae of exposure to bacterial and viral products, *Journal of Endocrinology* (Vol. 181, pp. 207-221).

Wingfield, J. C.; Maney, D. L.; Breuner, C. W.; Jacobs, J. D.; Lynn, S.; Ramenofsky, M. and Richardson, R. D. (1998). Ecological Bases of Hormone—Behavior Interactions: The “Emergency Life History Stage”. *American Zoologist*. 38(1): 191-206.