

2011-08

# A preliminary study of T-2 and HT-2 toxins in cereals sold in traditional market in South Korea

Kassim, Neema

Elsevier Ltd.

---

<https://doi.org/10.1016/j.foodcont.2011.02.023>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

## A preliminary study of T-2 and HT-2 toxins in cereals sold in traditional market in South Korea

N. Kassim, K. Kim, A.B. Mtenga, J.-E. Song, Q. Liu, W.-B. Shim, D.-H. Chung

To download full text click that link

DOI: <https://doi.org/10.1016/j.foodcont.2011.02.023>

### Abstract

Fusarium species are responsible for the production of harmful trichothecenes mycotoxins in cereals. These mycotoxins are cytotoxic, potentially immunosuppressive and potent fast-acting inhibitors of protein and nucleic acid synthesis. This study validated a HPLC method for simultaneous detection of T-2 and HT-2 toxins. The method was then used for the detection of T-2 and HT-2 toxins in cereals sold in traditional markets in Gyeongnam Province, South Korea. Seventy five samples analyzed, out of which 13 and 25 samples were found to be contaminated with T-2 (35.2–431.0 ng g<sup>-1</sup>) and HT-2 (21.1–442.7 ng g<sup>-1</sup>) toxins, respectively and 4 samples were found to be contaminated with both toxins. This study provides data on the contamination level of T-2 and HT-2 toxins in cereals in traditional market in Gyeongnam province, South Korea.

### Keywords

T-2 toxin; HT-2 toxin; Cereals; Simultaneous detection; Traditional markets