

2016-04-21

Antimicrobial and Cytotoxicity Activity of Clausena anisata, Acokanthera shemperii and Olea europaea Growing in Tanzania

Makirita, Winisia E.

European Journal of Medicinal Plants

<https://doi.org/10.9734/EJMP/2016/23635>

Provided with love from The Nelson Mandela African Institution of Science and Technology



Antimicrobial and Cytotoxicity Activity of *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea* Growing in Tanzania

Winisia E. Makirita¹, Leonard J. Chauka² and Musa Chacha^{1*}

¹*School of Life Science and Bioengineering, The Nelson Mandela African Institution of Science and Technology, P.O.Box 447, Arusha, Tanzania.*

²*Institute of Marine Science, University of Dar es Salaam, P.O.Box 668, Zanzibar, Tanzania.*

Authors' contributions

All authors worked together to achieve this work. All authors have cordially supported the work and preparation of the manuscript. Author WEM designed and supervised the study and prepared the first draft of the manuscript. Authors MC and LJC advised and guided the final draft of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJMP/2016/23635

Editor(s):

(1) Marcello Iriti, Professor of Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

Reviewers:

(1) Alba E. Vega, Universidad Nacional de San Luis, Argentina.

(2) Giusy Mandalari, University of Messina, Italy.

Complete Peer review History: <http://sciencedomain.org/review-history/14282>

Original Research Article

Received 11th December 2015
Accepted 3rd February 2016
Published 21st April 2016

ABSTRACT

Aims: To evaluate antimicrobial and cytotoxicity activities of *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea* against seven Gram negative bacteria and fungal species.

Study Design: Bioassay of antimicrobial assay was done using 96-well micro-dilution method.

Place and Duration of Study: School of Life Science and Bioengineering, Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania, from April 2014 to June 2014.

Methodology: 96-well micro dilution method was used in antimicrobial assay. Extracts were loaded in the wells of the first row, followed by serial dilution and 50 µl of the bacterial suspensions (0.5 MacFarland standard turbidity) were added in each well. The first concentration which showed no bacterial growth was considered as minimum inhibition concentration. Method developed by Meyer et al 1982 was adopted in cytotoxicity activities.

Results: All extracts indicated antibacterial activity on at least three to five of the tested seven bacteria and two fungi species with MIC value ranging 0.7812 - 12.5 mg/mL. The highest activity

*Corresponding author: E-mail: musa.chacha@nm-aist.ac.tz

was demonstrated by *Olea europaea* leaf methanolic, *Acokanthera shemperii* stem bark and *Clausena anisata* twigs ethyl acetate extracts with MIC value of 0.7812 mg/mL against *Pseudomonas aeruginosa* while the same MIC value was exhibited by *Olea europaea* stem bark methanol against *Proteus mirabilis*. However the *Olea europaea* root methanolic extract inhibited the growth of *Pseudomonas aeruginosa* and *Salmonella kisarawe* with MIC value of 0.7812 mg/mL. *Olea europaea* leaf methanolic and stem bark methanolic which demonstrated high antimicrobial activity were non toxic against brine shrimp larvae with LC₅₀ value of 369.8272 and 226.1566 µg/mL, while *Clausena anisata* twigs ethyl acetate, *Acokanthera shemperii* stem bark ethyl acetate and *Olea europaea* root methanolic extracts were toxic with LC₅₀ value of 6.21276, 67.4179 and 92.3089 µg/mL respectively.

Conclusion: This study has unveiled antimicrobial and cytotoxicity properties of *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea*.

Keywords: Antimicrobial; *Clausena anisata*; *Acokanthera shemperii*; *Olea europaea*.

1. INTRODUCTION

Antibiotics and antimicrobial agents have been used to treat infectious diseases for the last 70 years. Since 1940's antibiotics were reported to reduce the burden that the infectious diseases pose on human health [1]. However, the long term use of these drugs have given a chance to microorganism to adopt and make the drugs less effective [2]. It is known that, bacterial and some fungal species have ability to transmit and acquire resistance to drugs, which are utilized as therapeutic agents [3]. The emergence of multi-drug resistant pathogens such as *Escherichia coli*, *Klebsiella pneumoniae*, *Haemophilus* and *Candida albicans* has been a major challenge in the management of diseases caused by these pathogens [4,5]. For instance, in Tanzania, blood stream infection caused by resistant Gram negative bacteria and *C. albicans* is accounted to be a main source of mortality to children, with 34.9% mortality rate [6]. Therefore, the evidence of the rapid global spread of resistant clinical isolates necessitated the search for new class of drugs with unique mode of action [7,8]. However, there is widespread of resistance to new introduced antimicrobial agent which specify the short life expectancy for the new family of the introduced drugs [9]. Action must be taken to address and safeguard health and life in general for the poor communities which are the most vulnerable. The validation of ethnomedical information so as to develop herbal products from medicinal plants is of great interest to researchers.

Medicinal plants have been used as a primary health care by about 80% of individuals in the developing countries for many years [10]. For instance, communities in Tanzania use *Clausena anisata* leaves for treatment of oral candidiasis

[11]. This plant is also used by communities in West Africa for treatment of boils, ringworm, oral thrush and eczema [11]. Likewise the majority of sexually transmitted diseases in Tanzania and Kenya are treated by infusion of the pounded root of *Acokanthera shemperii* [12]. Moreover *Olea europaea* is extensively used to treat diarrhea, respiratory and urinary tract infections, stomach and intestinal diseases, and as mouth cleanser by different communities around the world [13]. In East-Africa infusion from *Olea europaea* bark is taken for tapeworm infestation [13]. Despite the contribution of medicinal plants for management of diseases, only limited number of medicinal plants have been scientifically validated [14]. Thus, this study reports the antimicrobial and cytotoxicity activity of *Acokanthera shemperii*, *Olea europaea* and *Clausena anisata* growing in Tanzania.

2. MATERIALS AND METHODS

2.1 Chemicals and Organisms Tested

Chloroform, ethyl acetate, methanol and dimethyl sulfoxide (DMSO) was purchased from Avantor performance materials India. Fluconazole was acquired from Lincoln Pharmaceuticals LTD, India, ciprofloxacin were bought from Micro Lab LTD, India and cyclophosphamide was bought from Khandelwa Laboratories Pvt Ltd (Mumbai), iodinitrotetrazolium chloride (INT) was purchased from SIGMA (Sigma Aldrich, St Louis, USA). Nutrient (agar and broth), sabouraud dextrose (agar and broth) were purchased from Hi Media Laboratories Pvt Ltd (Mumbai-India). Sea salt was prepared locally by evaporating water collected from the Indian Ocean, along the Dar es salaam Coast. *Klebsiella oxytoca* (clinical isolate), *Klebsiella pneumoniae* (ATCC700603), *Proteus mirabilis* (NCTC 1075), *Salmonella typhi*

(NCTC 8385), *Salmonella kisarawe* (clinical isolate), *Pseudomonas aeruginosa* (ATCC 29953), *Escherichia coli* (ATCC 25922), *Candida albicans* (ATCC 90028) and *Cryptococcus neoformans* (clinical isolate) were obtained from the Department of Microbiology, Muhimbili University of Health and Allied Sciences (MUHAS).

2.2 Sample Collection

Fruits, leaves, stem bark, twigs and roots of *C. anisata*, *A. schimperi* and *O. europaea* were collected from Monduli in Arusha region, Tanzania on 26th January 2015. Plant species were identified by a Botanist from Tropical Pesticides Research Institute (TPRI) Arusha, and the voucher specimens (CA300, AS201 and OE102) were kept at Nelson Mandela African Institute of Science and Technology, Arusha.

2.3 Extraction Process

Plant materials were under shade dried and pulverized into powder form for extraction. 800 g of the pulverized plant parts (fruits, leaves, twigs, stem bark and roots) were sequentially macerated using chloroform, ethyl acetate and methanol for 48 hrs twice for each solvent. The respective extracts were filtered through Whatman filter paper number 1 in a glass column. The crude extracts were obtained after subjecting the macerated plant materials in a rotary evaporator and the extracts were stored in refrigerator at -20°C until testing time.

2.4 Testing for Antimicrobial Activity

Extracts were tested against representative Gram negative bacteria; *S. typhi* (NCTC 8385), *P. aeruginosa* (ATCC 29953), *S. kisarawe* (clinical isolate), *E. coli* (ATCC 25922), *K. oxytoca* (clinical isolate), *K. pneumoniae* (ATCC700603), *P. mirabilis* (NCTC 1075), *C. albicans* (ATCC 90028) and *C. neoformans* (clinical isolate). Selection of the microorganism was based on their availability during the experiment. Minimum inhibitory concentrations (MICs) were obtained using microdilution method. Stock solution was prepared by dissolving 100 mg of extract in 1 mL of DMSO (100 mg/mL). Each of the 96 well microtitre plates were first preloaded with 50 µl of broth media, followed up by 50 µl of extracts in first well of each row which make up 100 µl total volume in the first wells. Subsequently 50 µl

were transferred from the first rows of each category to the second rows and the process was repeated down the columns on which the last wells at the bottom 50 µl were discarded. Gentamicin (100 µg/mL) was used as standard drug (positive control); DMSO as negative control and the row contains only broth as growth control. Thereafter 50 µl of the bacterial suspensions (0.5 MacFarland standard turbidity) were added in each well and incubated at 37 °C for 24 h. 40 µl of 0.02% p-iodonitrotetrazolium (INT) chlo-ride solution was added to each well followed by incubating for 1 h at 37°C. Bacteria growth was indicated by change in color of INT (pink color). Absence of bacterial was indicated with no change in dye's color. The first concentration which showed no bacterial growth was considered as Minimum inhibition concentration (MIC).

2.5 Brine Shrimps Lethality Test

Brine shrimps lethality test was conducted as described by Meyer et al. [15]. Stock solution was prepared by dissolving 40 mg of extracts in 1 mL DMSO (40 mg/mL). Different concentration levels (240, 120, 80, 40, 24 and 8 µg/mL) was prepared by drawing different volume from stock solution and was added in a 10 mL universal bottle containing 10 brine shrimps larvae. Volume was then adjusted to 5 mL with artificial sea water prepared by dissolving 3.8 g of sea salt in 1 L of distilled water and incubated under light for 24 hrs. Each concentration was tested in duplicate for statistical significance. Negative controls contain brine shrimp, DMSO and artificial sea water and cyclophosphamide was used as a positive control. Dead larvae were identified and the mean results of the percentage mortality were plotted against the logarithms of concentrations using the Fig P computer program. LC₁₆, LC₅₀, LC₈₄ and the 95% CI values were calculated from the Regression equation obtained from the graph.

3. RESULTS

3.1 Antimicrobial Activity

All extracts from the selected plants demonstrated antimicrobial activity on at least three to five of the tested bacteria and fungi species with minimum inhibition concentration (MIC) value ranging 0.7812-12.5 mg/mL (Table 1). The highest activity of MIC value 0.7812 mg/mL was demonstrated by

Olea europaea root methanolic extract, *Acokanthera shemperii* stem bark ethyl acetate and *Clausena anisata* twigs ethyl acetate extracts against *Pseudomonas aeruginosa*. The same MIC value was also exhibited by *O. europaea* stem bark methanolic extract against *Proteus mirabilis*. A number of extracts demonstrated antimicrobial activity with MIC value of 1.5625 mg/mL against the tested pathogens. For instance *C. anisata* leaf chloroform, ethyl acetate and methanolic extracts exhibited MIC value of 1.5625 mg/mL against *Salmonella typhi*. Similarly *C. anisata* twigs ethyl acetate indicated the same MIC value of 1.5625 mg/mL against *S. typhi*, *Escherichia coli*, *Salmonella kisarawe* and *Klebsiella oxytoca*. Likewise *C. anisata* stem bark chloroform and *A. shemperii* leaf ethyl acetate inhibited the growth of *S. typhi*, *Klebsiella oxytoca* and *P. aeruginosa* with MIC values of 1.5625 mg/mL. Another extracts that demonstrated the MIC value of 1.5625 mg/mL are *A. shemperii* stem bark chloroform and *A. shemperii* root chloroform against *P. aeruginosa*. Furthermore the MIC value of 1.5625 mg/mL was also showed by *O. europaea* leaf methanolic against *S. typhi*, *E. coli* and *Klebsiella pneumoniae*. Similarly *O. europaea* roots methanolic exhibited the same MIC value of 1.5625 mg/mL against *S. typhi*, *K. oxytoca* and *K. pneumoniae*. Minimum inhibitory concentrations ranging from 3.125-12.5 mg/mL were indicated by the remaining extracts. Despite the fact that Rios and Recio (2005), recommended the maximum concentration of interest should be less than 1 mg/mL, however the extracts with low antimicrobial activity should also be reported as it can be incorporated with other extracts to improve its biological importance.

3.2 Brine Shrimp Lethality Tests

Extracts were evaluated for lethality activity against the brine shrimp larvae and results are summarized in Table 2. According to Meyer et al [15], extracts that demonstrates LC₅₀ value greater than 100 µg/mL was considered as non-toxic and LC₅₀ value less than 100 µg/mL as toxic. *Acokanthera shemperii* root ethyl acetate, *C. anisata* leaf chloroform, *C. anisata* twigs ethyl acetate and *A. shemperii* root chloroform were more toxic with LC₅₀ less than 20 µg/mL.

4. DISCUSSION

Healing properties of medicinal plants that are used to manage infectious diseases has been

proven by several studies conducted to evaluate antimicrobial activities of medicinal plants [16,17]. Medicinal plants are known to cure infectious disease such as urinary tract infections, gastrointestinal disorders, respiratory diseases and cutaneous infections [18,19]. The antimicrobial activity of plants materials against bacteria and fungi strains is due to their chemical composition [20]. Since secondary metabolites are produced in response to fungal and bacterial challenges that plant is facing.

Secondary metabolites produced by a medicinal plant in one region may be similar or different from the same species in another region. For instance *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea* have been exploited in different parts of Africa for management of infectious diseases [21-23]. These uses prompted scientists to validate the antimicrobial properties of *C. anisata*, *A. shemperii* and *O. europaea*. For instance the study conducted on *O. europaea* leaf acetone extract inhibited the growth of *Salmonella enteritidis*, *Bacillus cereus*, *Klebsiella pneumoniae*, *Escherichia coli*, *Enterococcus faecalis*, *Streptococcus thermophilus* and *Lactobacillus bulgaricus* [24]. Likewise the study conducted in Ethiopia on the *A. schimperi* methanolic and water extracts showed the growth inhibition of *S. aureus*, *S. pyogenes*, *E. coli*, *P. aeruginosa* and *P. vulgaris* [25]. It is however interesting to observe that the current study revealed that chloroform, ethyl acetate and methanolic extracts of *C. anisata*, *A. shemperii* and *O. europaea* growing in Tanzania had antimicrobial activity with MIC value ranging between 0.7812-25 mg/mL against microbes tested in this study. Furthermore the investigations conducted on the leaf essential oil from *C. anisata* growing in India demonstrated the inhibition effects on *S. typhi* and *P. aeruginosa* with the minimum inhibitory concentration (MIC) values of 62.5 and 125 µg/mL, respectively [26]. Results of antimicrobial activity might differ due to the specifics of collection, solvent and the method used for extraction [27]. Furthermore, Rios et al. [27] proposed that the antimicrobial activity is of interest when the concentration is below 100 µg/mL, but weak compounds should be used with other compounds to improve their activities. Thus based on that study, extracts that show antimicrobial activity in this study can be used with other extracts or with other compounds when isolated to improve its activity.

Table 1. Antimicrobial activity *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea*

| Extracts | Minimum inhibition concentration MIC (mg/mL) | | | | | | | | |
|----------|--|---------------------|----------------|--------------------|-------------------|----------------------|----------------------|--------------------|----------------------|
| | <i>S. typhi</i> | <i>P. mirabilis</i> | <i>E. coli</i> | <i>S. kisarawe</i> | <i>K. oxytoca</i> | <i>K. pneumoniae</i> | <i>P. aeruginosa</i> | <i>C. albicans</i> | <i>C. neoformans</i> |
| CALC | 1.5625 | 3.125 | 6.25 | 3.125 | 3.125 | 3.125 | 3.125 | 3.125 | 1.5625 |
| CALE | 1.5625 | 3.125 | 6.25 | 3.125 | 6.25 | 3.125 | 3.125 | 6.25 | 1.5625 |
| CALM | 1.5625 | 3.125 | 12.5 | 6.25 | 6.25 | 3.125 | 3.125 | 6.25 | 3.125 |
| CATC | 6.25 | 3.125 | 12.5 | 3.125 | 6.25 | 3.125 | 3.125 | 25 | 12.5 |
| CATE | 1.5625 | 3.125 | 1.563 | 1.563 | 1.5625 | 6.25 | 0.7812 | 12.5 | 6.25 |
| CATM | 3.125 | 3.125 | 3.125 | 3.125 | 3.125 | 6.25 | 1.5625 | 12.5 | 6.25 |
| CASBC | 1.5625 | 6.25 | 3.125 | 6.25 | 1.5625 | 6.25 | 1.5625 | 12.5 | 6.25 |
| CASBM | 3.125 | 6.25 | 3.125 | 6.25 | 6.25 | 6.25 | 6.25 | 12.5 | 12.5 |
| CAFC | 3.125 | 3.125 | 6.25 | 6.25 | 6.25 | 6.25 | 3.125 | 12.5 | 12.5 |
| CAFE | 3.125 | 3.125 | 6.25 | 3.125 | 3.12 | 6.25 | 6.25 | 12.5 | 3.125 |
| CAFM | 3.125 | 3.125 | 3.12 | 6.25 | 3.12 | 12.5 | 3.12 | 12.5 | 1.5625 |
| ASLC | 12.5 | 12.5 | 12.5 | 12.5 | 6.25 | 12.5 | 6.25 | 12.5 | 3.125 |
| ASLE | 1.5625 | 12.5 | 12.5 | 6.25 | 1.5625 | 12.5 | 1.5625 | 12.5 | 3.125 |
| ASLM | 12.5 | 12.5 | 12.5 | 3.125 | 12.5 | 3.125 | 3.125 | 12.5 | 12.5 |
| ASSBC | 6.25 | 12.5 | 12.5 | 6.25 | 6.25 | 6.25 | 1.5625 | 25 | 25 |
| ASSBE | 6.25 | 6.25 | 6.25 | 6.25 | 6.25 | 6.25 | 0.78125 | 12.5 | 6.25 |
| ASSBM | 12.5 | 12.5 | 12.5 | 6.25 | 6.25 | 12.5 | 3.125 | 12.5 | 12.5 |
| ASRC | 12.5 | 12.5 | 12.5 | 12.5 | 12.5 | 12.5 | 1.5625 | 12.5 | 6.25 |
| ASRE | 12.5 | 6.25 | 12.5 | 12.5 | 6.25 | 6.25 | 3.125 | 12.5 | 3.125 |
| ASRM | 6.25 | 12.5 | 6.25 | 6.25 | 3.125 | 3.125 | 6.25 | 25 | 12.5 |
| OELC | 12.5 | 6.25 | 12.5 | 12.5 | 25 | 12.5 | 12.5 | 12.5 | 6.25 |
| OELE | 3.125 | 6.25 | 3.125 | 6.25 | 3.125 | 6.25 | 3.125 | 12.5 | 3.125 |
| OELM | 1.5625 | 3.125 | 1.562 | 6.25 | 0.7812 | 1.5625 | 6.25 | 12.5 | 6.25 |
| OESBC | 6.25 | 6.25 | 12.5 | 12.5 | 25 | 12.5 | 12.5 | 6.25 | 25 |
| OESBE | 3.125 | 12.5 | 3.125 | 6.25 | 6.25 | 3.125 | 6.25 | 25 | 12.5 |

| Extracts | Minimum inhibition concentration MIC (mg/mL) | | | | | | | | |
|----------|--|---------------------|----------------|--------------------|-------------------|----------------------|----------------------|--------------------|----------------------|
| | <i>S. typhi</i> | <i>P. mirabilis</i> | <i>E. coli</i> | <i>S. kisarawe</i> | <i>K. oxytoca</i> | <i>K. pneumoniae</i> | <i>P. aeruginosa</i> | <i>C. albicans</i> | <i>C. neoformans</i> |
| OESBM | 3.125 | 0.7812 | 3.125 | 1.5625 | 3.125 | 3.125 | 6.25 | 25 | 3.125 |
| OERC | 12.5 | 12.5 | 12.5 | 6.25 | 12.5 | 12.5 | 12.5 | 25 | 25 |
| OERE | 6.25 | 12.5 | 12.5 | 6.25 | 6.25 | 25 | 3.125 | 25 | 6.25 |
| OERM | 1.5625 | 6.25 | 3.125 | 0.7812 | 1.5625 | 1.5625 | 0.7812 | 25 | 6.25 |
| OETC | 6.25 | 3.125 | 6.25 | 6.25 | 12.5 | 6.25 | 12.5 | 25 | 6.25 |
| OETE | 3.125 | 6.25 | 6.25 | 6.25 | 6.25 | 25 | 6.25 | 25 | 6.25 |
| OETM | 3.125 | 6.25 | 3.12 | 12.5 | 6.25 | 6.12 | 12.5 | 12.5 | 25 |
| Cipro | 0.391 | 0.7812 | 0.391 | 0.7812 | 0.7812 | 0.391 | 0.391 | NA | NA |
| Fluco | NA | NA | NA | NA | NA | NA | NA | 1.5625 | 0.7812 |

Key: CALC- *C. anisata* leaf chloroform, CALE- *C. anisata* leaf ethyl acetate, CALM- *C. anisata* leaf methanolic, CAFC- *C. anisata* fruits chloroform, CAFE- *C. anisata* fruits ethyl acetate, CAFM- *C. anisata* fruits methanolic, CATC- *C. anisata* twigs chloroform, CATE- *C. anisata* twigs ethyl acetate, CATM- *C. anisata* twigs methanolic, CASBC- *C. anisata* stem bark chloroform, CASBE- *C. anisata* stem bark ethyl acetate, CASBM- *C. anisata* stem bark methanolic, ASLC- *A. shemperii* leaf chloroform, ASLE- *A. shemperii* leaf ethyl acetate, ASLM- *A. shemperii* leaf methanolic, ASSBC- *A. shemperii* stem bark chloroform, ASSBE- *A. shemperii* stem bark ethyl acetate, ASSBM- *A. shemperii* stem bark methanolic, ASRC- *A. shemperii* root chloroform, ASRE- *A. shemperii* root ethyl acetate, ASRM- *A. shemperii* root methanolic, OELC – *O. europaea* leaf chloroform, OELE- *O. europaea* leaf ethyl acetate, OELM- *O. europaea* leaf Methanol, OESBC- *O. europaea* stem bark chloroform, OESBE- *O. europaea* stem bark ethyl acetate, OESBM- *O. europaea* stem bark methanol, OETC- *O. europaea* stem bark twigs chloroform, OETE- *O. europaea* twigs ethyl acetate, OETM- *O. europaea* twigs ethyl acetate, OERC- *O. europaea* roots chloroform, OERE- *O. europaea* roots ethyl acetate, OERM- *O. europaea* roots methanol, Cipro- ciprofloxacin, Fluco- Fluconazole and NA-not applicable

Table 2. Brine shrimp lethality tests of *Clausena anisata*, *Acokanthera shemperii* and *Olea europaea*

| Extract | LC50 (µg/ml) | 95% (confidence interval) | R ² | Regression equation |
|------------------|--------------|---------------------------|----------------|-------------------------|
| CALC | 3.5761 | 1.9119-6.68865 | 0.902 | y = 34.634logx + 30.833 |
| CALE | 36.6689 | 26.7263-50.30973 | 0.9763 | y = 62.591logx - 47.911 |
| CALM | 127.7264 | 79.2052-205.9715 | 0.9851 | y = 41.428logx - 37.259 |
| CATC | 115.0821 | 94.8258-139.6653 | 0.8885 | y = 125.23logx - 208.1 |
| CATE | 6.1276 | 3.6043-10.4172 | 0.9857 | y = 45.688logx + 14.03 |
| CATM | 28.446 | 21.6340-37.4025 | 0.9699 | y = 79.221logx - 65.189 |
| CASBC | 99.0329 | 69.1918-141.7438 | 0.9738 | y = 55.209logx - 60.185 |
| CASBM | 122.419 | 71.3323-210.0927 | 0.9657 | y = 36.653logx - 26.526 |
| CAFC | 20.1162 | 13.8802-29.1540 | 0.9488 | y = 53.35logx - 19.479 |
| CAFE | 70.045 | 50.8541-96.4780 | 0.9363 | y = 61.829logx - 64.098 |
| CAFM | 1226.557 | 522.623-2878.631 | 0.9149 | y = 23.205logx - 21.673 |
| ASLC | 37.8853 | 23.113-62.0996 | 0.9211 | y = 40.059logx - 13.232 |
| ASLE | 808.289 | 370.162-1764.989 | 0.9418 | y = 25.348logx - 23.701 |
| ASLM | 56.003 | 42.043-74.5982 | 0.9162 | y = 69.046logx - 70.707 |
| ASSBC | 125.8478 | 78.668-201.3215 | 0.9741 | y = 42.135logx - 38.477 |
| ASSBE | 67.4179 | 29.425-154.4678 | 0.9021 | y = 23.878logx + 6.3325 |
| ASSBM | 222.0638 | 122.924-401.1586 | 0.8888 | y = 33.474logx - 28.546 |
| ASRC | 16.597 | 12.239-22.507 | 0.8727 | y = 79.6logx - 47.115 |
| ASRE | 14.3511 | 9.976-20.6448 | 0.9039 | y = 59.636logx - 18.992 |
| ASRM | 71.8351 | 53.9871-95.5835 | 0.904 | y = 69.308logx - 78.659 |
| OELC | 247.7954 | 85.7587- 715.991 | 0.9914 | y = 18.657logx + 5.3334 |
| OELE | 50.3363 | 29.70- 85.3109 | 0.8906 | y = 37.532logx - 13.875 |
| OELM | 369.8272 | 249.0529-549.169 | 0.947 | y = 70.81logx - 131.84 |
| OESBC | 1627.450 | 642.219-4124.125 | 0.9682 | y = 21.29logx - 18.373 |
| OESBE | 1164.063 | 468.1878-3359.85 | 0.8774 | y = 21.735logx - 16.639 |
| OESBM | 226.1566 | 141.643-361.097 | 0.9357 | y = 42.307logx - 49.608 |
| OERC | 11266.861 | 3206.30-39591.4 | 0.9274 | y = 15.752logx - 13.824 |
| OERE | 67.4264 | 36.1951-125.6060 | 0.9462 | y = 31.821logx - 8.1952 |
| OERM | 92.3089 | 52.0523-163.699 | 0.9109 | y = 34.555logx - 17.909 |
| OETC | 52.9775 | 21.1128-132.934 | 0.9616 | y = 21.518logx + 12.901 |
| OETE | 599.565 | 299.126-1201.76 | 0.8782 | y = 28.47logx - 29.085 |
| OETM | 285,991.20 | 38126.4-2145260 | 0.9019 | y = 9.8242logx - 3.6043 |
| Cyclophosphamide | 16.37 | 12.01-22.31 | 0.995 | y=69.97logx-34.936 |

In Tanzania leaves of *C. anisata* are used for management of skin fungal infections and oral candidiasis. Since the extracts from our study indicated activity against *C. albicans* and *C. neoformans*, extracts could be useful for treatment of fungal infections. Brine shrimp results indicated that some of the extracts that demonstrated antimicrobial activity are nontoxic while others are toxic. According to Meyer et al [15], extracts with LC₅₀ value less than 20 µg/mL are considered as potential anticancer agents. In this study *C. anisata* leaf chloroform, *C. anisata* twigs ethyl acetate, *A. shemperii* root chloroform and *A. shemperii* root ethyl acetate extracts which demonstrated LC₅₀ value of 3.5761, 6.1276, 16.597 and 14.3511 µg/mL respectively are potential anticancer agents. It is of high interest that our study displayed two advantages

for therapeutic industry, as compounds from these extracts are potential antimicrobial and anticancer drug leads. Thus the use of *C. anisata*, *O. europaea* and *A. shemperii* is therefore validated in this study.

5. CONCLUSION

This study revealed antimicrobial and cytotoxicity activity of *Clausena anisata*, *Olea europaea* and *Acokanthera shemperii*.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Halloran J. The overuse of antibiotics in food animals threatens public health. Consumer Union. Policy & Action For Consumer Reports. 2012. Available: <https://consumersunion.org/news/the-overuse-of-antibiotics-in-food-animals-threatens-public-health-2>
- Rachana S, Et Al. Comparative analysis of antibacterial activity of *Jatropha curcas* fruit parts. Journal of Pharmaceutical and Biomedical Sciences (Jpbms). 2012; 15(15):1-4.
- Cohen ML. Epidemiology of drug resistance: Implications for a Post-antimicrobial Era. Science. 1992. 257(5073):1050-1055.
- Akram M, Shahid M, Khan AU. Etiology and antibiotic resistance patterns of community-acquired urinary tract infections in Jnmc Hospital Aligarh, India. Annals of Clinical Microbiology and Antimicrobials, 2007;6(1):4.
- Khan AU, Musharraf A. Plasmid-mediated multiple antibiotic resistance in proteus mirabilis isolated from patients with urinary tract infection. Medical Science Monitor: International Medical Journal of Experimental and Clinical Research. 2004; 10(11):Cr598-602.
- Kumar A, et al. A Clinical resistant isolate of opportunistic fungal pathogen, candida albicans revealed more rigid membrane than its isogenic sensitive isolate. Multidisciplinary Approaches for Studying and Combating Microbial Pathogens, 2015;1.
- Cordell GA. Biodiversity and Drug Discovery— a symbiotic relationship. Phytochemistry. 2000;55(6):463-480.
- Mbwambo ZH, et al. Antibacterial, antioxidant, cytotoxic activities and preliminary phytochemical screening of extracts from *Combretum schumannii* Engl. Journal of Medicinal Plants Research. 2013;7(33):2483-2488.
- Coates A, et al. The future challenges facing the development of new antimicrobial drugs. Nature Reviews Drug Discovery. 2002;1(11):895-910.
- Nascimento GG, et al. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. Brazilian Journal of Microbiology. 2000;31(4):247-256.
- Hamza OJ, et al. Antifungal activity of some Tanzanian plants used traditionally for the treatment of fungal infections. Journal of Ethnopharmacology. 2006; 108(1):124-132.
- Record P. Prota 11 (1): Medicinal Plants/Plantes Medicinales 1. Record Display; 2012. Consultado 23/01/2012. Available: <http://database.prota.org/Protahtml/Plantago%20lanceolata>
- Hashmi MA, et al. Traditional uses, phytochemistry, and pharmacology of *Olea europaea* (Olive). Evidence-based Complementary and Alternative Medicine; 2015.
- Khan R, et al. Antimicrobial activity of five herbal extracts against multi drug resistant (Mdr) strains of bacteria and fungus of clinical origin. Molecules. 2009;14(2):586-597.
- Meyer B, et al. Brine shrimp: A convenient general bioassay for active plant constituents. Planta Medica. 1982;(45): 31-4.
- Bonjar S. Evaluation of antibacterial properties of some medicinal plants used in Iran. Journal of Ethnopharmacology. 2004;94(2):301-305.
- Lee SB, et al. The Antimicrobial activity of essential oil from *Dracocephalum foetidum* against pathogenic microorganisms. Journal of Microbiology (Seoul, Korea). 2007;45(1):53-57.
- Brantner A, Grein E. Antibacterial activity of plant extracts used externally in traditional medicine. Journal of Ethnopharmacology. 1994;44(1):35-40.
- Somchit M, et al. *In vitro* antimicrobial activity of ethanol and water extracts of *Cassia alata*. Journal of Ethnopharmacology. 2003;84(1):1-4.
- Obinna O, John A, Peter OO. The effect of extraction protocol on the phytochemical and antimicrobial activities of *Lantana camara* leaf extract found within a local environment; 2013.
- Arbab IA, et al. A review of traditional uses, phytochemical and pharmacological

- aspects of selected members of *Clausena* genus (Rutaceae). Journal of Medicinal Plants Research. 2012;6(38):5107-5118.
22. Ben Salem M, et al. Pharmacological activities of *Olea europaea* leaves. Journal of Food Processing and Preservation; 2014.
 23. Neuwinger HD. African ethnobotany: Poisons and drugs: Chemistry, pharmacology, toxicology; 1996. Crc Press.
 24. Korukluoglu M, et al. Antibacterial activity and chemical constitutions of *Olea europaea* L. leaf extracts. Journal of Food processing and preservation. 2010;34(3): 383-396.
 25. Taye B, et al. Antibacterial activities of selected medicinal plants in traditional treatment of human wounds in Ethiopia. Asian Pacific Journal of Tropical Biomedicine. 2011;1(5):370-375.
 26. Senthilkumar A, Venkatesalu V. Phytochemical Analysis and antibacterial activity of the essential oil of *Clausena anisata* (Willd.) Hook. F. Ex Benth. International Journal of Integrative Biology. 2009;5(2):116.
 27. Rios J, Recio M. Medicinal plants and antimicrobial activity. Journal of Ethnopharmacology. 2005;100(1):80-84.

© 2016 Makirita et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://sciencedomain.org/review-history/14282>