NM-AIST Repository	https://dspace.mm-aist.ac.tz

Life sciences and Bio-engineering

Masters Theses and Dissertations [LiSBE]

2022-06

A comparative analysis of determinants of low birth weight and stunting among under-five children of adolescent and non-adolescent mothers using the 2015/16 Tanzania demographic and health survey

Mtongwa, Ramadhani

NM-AIST

https://doi.org/10.58694/20.500.12479/1520 Provided with love from The Nelson Mandela African Institution of Science and Technology

A COMPARATIVE ANALYSIS OF DETERMINANTS OF LOW BIRTH WEIGHT AND STUNTING AMONG UNDER-FIVE CHILDREN OF ADOLESCENT AND NON-ADOLESCENT MOTHERS USING THE 2015/16 TANZANIA DEMOGRAPHIC AND HEALTH SURVEY

Ramadhani H. Mtongwa

A Dissertation Submitted in Partial Fulfillment of the Requirements for the Degree of Master of Science in Public Health Research of the Nelson Mandela African Institution of Science and Technology

Arusha, Tanzania

ABSTRACT

In Tanzania, adolescent pregnancies increased from 23% to 27% between 2012 and 2015. The rises have been linked to poor birth outcomes, pregnancy complications, childhood malnutrition, and maternal and child deaths. Using data from the 2015 Tanzania Demographic and Health Survey, this study investigated the factors associated with low birth weight (LBW) and under-five stunting among adolescent and non-adolescent mothers. Data from 13 266 women collected as part of the Tanzania Demographic and Health Survey in 2015/2016 were re-analyzed using STATA version 14 software while taking survey design into account. The outcome variables were low birth weight and stunting. Logistic regression models were used to identify factors that contributed to LBW and stunting in children born to adolescent and non-adolescent mothers. After controlling for potential confounders, this study discovered that non-adolescent mothers had a lower risk of having LBW babies than adolescent mothers (AOR = 0.34; 95 per cent CI: 0.22-0.50). Maternal malnutrition (AOR = 2.29; 95 percent CI: 1.43-3.67), divorce, separation, or widowhood (AOR = 1.76; 95)percent CI: 1.24-2.50), and fewer than four antenatal care (ANC) visits (AOR = 0.64; 95 percent CI: 0.49–0.83) were all associated with LBW. Stunting in children was not related to maternal age. Maternal high socioeconomic status (AOR = 0.69; 95% CI: 0.57-0.84) and maternal obesity or overweight (AOR = 0.77; 95% CI: 0.64-0.92) were associated with stunting. Stunting was found to be significantly associated with birth weight, gender, and age. Maternal age predicted LBW but not stunting. To reduce poor birth outcomes, a multi-sectoral approach is required to address childhood stunting and teenage pregnancies.

DECLARATION

I, Ramadhani H. Mtongwa, hereby declare to the Senate of the Nelson Mandela African Institution of Science and Technology that this dissertation is my original work and that it has neither been submitted nor concurrently submitted for a degree or similar award in any other institution.

Ramadhani H. Mtongwa

Name and Signature of the Candidate

The above declaration is confirmed by:

Dr. Ester Elisaria

Name and Signature of Supervisor

11th July 2022

11th July 2022 Date

Date

Dr. Abdallah Mkop

Falim

11th July 2022

Date

Name and Signature of Supervisor 2

COPYRIGHT

This dissertation/thesis is copyright material protected under the Berne Convention, the Copyright Act of 1999 and other international and national enactments, on behalf, of intellectual property. It must not be reproduced by any means, in full or in part, except for short extracts in fair dealing; for researcher private study, critical scholarly review or discourse with an acknowledgement, without the written permission of the office of Deputy Vice-Chancellor for Academic, Research and Innovation on behalf of both the author and NM-AIST.

CERTIFICATION

The undersigned certify that they have read and hereby recommend for acceptance by the Senate of the Nelson Mandela African Institution of Science and Technology the dissertation entitled "A Comparative Analysis of Determinants of Low Birth Weight and Stunting Among Under-Five Children of Adolescent and Non-Adolescent Mothers Using 2015/16 Tanzania Demographic and Health Survey" in partial fulfilment of the requirements for the Degree of Master of Science in Public Health Research of the Nelson Mandela African Institution of Science and Technology.

Dr. Ester Elisaria

Name and Signature of Supervisor 1

Date

Dr. Abdallah Mkop

lop

Name and Signature of Supervisor 2

Date

ACKNOWLEDGEMENTS

I would like to express my deepest gratitude to the following: My supervisors Dr. Ester Elisaria, my co-supervisor Dr. Abdallah Mkopi and Mr. Charles Festo, Head of data unit Ifakara Health Institute (IHI) for their excellent guidance, willingness to help and give the best suggestions throughout this work. The African Development Bank (AfDB) and (IHI) for funding my research study. The Nelson Mandela African Institution of Science and Technology for allowing me to study the course Public Health Research. The IHI for providing ethical clearance that allowed me to conduct this research without forgetting the World Health Organization (WHO) and Tanzania National Bureau of Statistics (NBS) for providing the needful dataset through the Demographic and Health Survey.

DEDICATION

I would like to dedicate this work to: My wife ms Jazila for her support and encouragement. My parents Mr. and Mrs. Hashim for their encouragement and best wishes, my brothers and sisters for their unlimited support and encouragement.

TABLE OF CONTENTS

ABS	IRACTi
DEC	LARATIONii
COP	YRIGHT iii
CER	TIFICATIONiv
ACK	NOWLEDGEMENTSv
DED	ICATIONvi
TAB	LE OF CONTENTS
LIST	OF TABLESx
LIST	OF FIGURESxi
LIST	OF ABBREVIATIONS AND SYMBOLSxii
CHA	PTER ONE1
INTR	RODUCTION1
1.1	Background of the Study1
1.2	Statement of the Problem
1.3	Rationale of the Study
1.4	Research Objectives
	1.4.1 General Objectives
	1.4.2 Specific Objectives
1.5	Research Questions
1.6	Significance of the Study
1.7	Delineation of the Study
CHA	PTER TWO
LITE	RATURE REVIEW
2.1	Overview
2.2	The Association between Mother's Age (Adolescent and Non-Adolescent Mothers) and
	Childbirth Outcomes

	2.2.1	Adolescent versus Non-Adolescent Mothers and Low Birth Weight
	2.2.2	Adolescents versus Non-Adolescents and Preterm Births7
2.3	The A Stuntin	ssociation between Mothers' Age (Adolescent and Non-Adolescent) and Under-five ng Status
	2.3.1	Adolescent Mothers versus Non-Adolescent Mothers and Stunting among under-five
2.4	Materi	nal Factors Associated with Low Birth Weight and Under-five Stunting Status among
	Adoles	scent and Non-Adolescent Mothers
	2.4.1	Maternal Education Levels and Birth Outcomes9
	2.4.2	Maternal Occupation and Birth Outcomes10
	2.4.3	Maternal Place of Residence and Birth Outcomes11
	2.4.4	Maternal Nutrition Status and Birth Outcomes11
	2.4.5	Maternal Service Utilization and Birth Outcomes
	2.4.6	Maternal Occupation and Child Stunting
	2.4.7	Maternal Education Level Child Stunting14
	2.4.8	Maternal Place of Residence and Child Stunting14
	2.4.9	Maternal Nutrition Status and Under-five Stunting15
CHA	APTER '	THREE16
MA	ΓERIAI	S AND METHODS
3.1	Data S	ource
3.2	Resear	rch Design16
3.3	Target	Population16
3.4	Sampl	ing Procedures16
3.5	Data N	Anagement and Analysis17
	3.5.1	Subjects and Sample Size17
	3.5.2	Categorization of Variables17
	3.5.3	Statistical Analysis
3.6	Ethica	l Considerations

CHA	PTER 1	FOUR	20
RES	ULTS A	AND DISCUSSION	20
4.1	Result	S	.20
	4.1.1	Association between Mother's Age (Adolescent and Non-adolescent Mothers) a Under-five stunting	und .20
	4.1.2	Prevalence of Low Birth Weight in Children from Adolescent and Non-adolescent Mothers	ent .24
	4.1.3	Prevalence of Under-five Stunting, Wasting and Underweight	.26
	4.1.4	Prevalence of stunting among under-five children according to matern characteristics	nal .26
	4.1.5	Predictors of Low Birth Weight	.28
	4.1.6	Predictors of Stunting in Under-five Children	.29
4.2	Discus	ssion	.31
	4.2.1	The association between Adolescent and Non-adolescent Mothers' Characterist and Childbirth Weight	ics .31
	4.2.2	The Association between Adolescent and Non-adolescent Maternal Characterist and Stunting in Under-five Children	ics .33
	4.2.3	Relationship between Stunting and Child's Birth Weight, Sex and Age	.34
CHA	PTER 1	FIVE	.36
CON	ICLUSI	ON AND RECOMMENDATIONS	.36
5.1	Conclu	usion	.36
5.2	Recon	nmendations	.36
REF	ERENC	CES	.37
RES	EARCH	I OUTPUTS	.50

LIST OF TABLES

Table 1:	Household and individual interviews
Table 2:	Socio-economic and Demographic Characteristics of Mothers
Table 3:	Anthropometric and nutrition status of mothers
Table 4:	Prevalence of Low Birth Weight in children from adolescent and non-adolescent mothers by maternal characteristics (N= 6386)25
Table 5:	Prevalence of under-five stunting, wasting and underweight
Table 6:	Prevalence of child stunting among adolescent and non-adolescent mothers by maternal
	characteristics (N=8852)27
Table 7:	Factors associated with Low Birth Weight (N=6385)29
Table 8:	Factors associated with stunting (N=8852)

LIST OF FIGURES

Figure 1:	Maternal level of education	21
Figure 2:	Maternal Body Mass Index (BMI)	23
Figure 3:	Gestation age at 1 st Antenatal Care Visits	24
Figure 4:	Low birth weight status of the study participants	25
Figure 5:	Stunting prevalence according to maternal age	27

LIST OF ABBREVIATIONS AND SYMBOLS

Average Birth Rate
African Development Bank
Antenatal Care
Adjusted Odds Ratio
Confidence Interval
Body Mass Index
Exclusive Breastfeeding
Demographic and Health Survey
Low Birth Weight
Maternal Health Services
Mid Upper Arm Circumference
National Bureau of Statistics
Prevention of Mother to Child Transmission
Socio Economic Status
Tanzania Demographic and Health Survey
Tanzania National Nutrition Survey
United Nations
United Nations Population Fund
United Nations Children Fund
United Republic of Tanzania
World Health Organisation
Girls not Brides
Global Action Report on Preterm Birth
Ministry of Health
Sexually Transmitted Infections
Human Immunodeficiency Virus,
Acquired Immunodeficiency Syndrome
Ifakara Health Institute Review Board

CHAPTER ONE

INTRODUCTION

1.1 Background of the Problem

Globally, about 1.2 billion people (out of 7.6 billion) are adolescents aged 10–19 years (World Health Organization [WHO], 2018b). Individuals aged 10–14 years are classified as early adolescent, while those aged 15–19 years are classified as a late adolescent (WHO, 2018c). Over 580 million adolescents are female worldwide. Asia is the leading continent with the highest number of adolescents in the world, and in sub-Saharan Africa, 23% of the population are adolescents aged 10–19 (United Nations Children Fund [UNICEF], 2019).

It is expected that from 2010 to 2030, the number of countries with more than 5 million adolescent girls will increase from 16 to 18 (United Nations Population Fund [UNFPA], 2013). The largest absolute national increases in adolescent girls will mostly happen in sub-Saharan Africa. Nigeria (9.2 million), the United Republic of Tanzania (3.7 million), the Democratic Republic of the Congo (3.3 million), Uganda (2.5 million), and Kenya (2.3 million) are the five countries with the highest absolute increases in adolescent population (UNFPA, 2013). The increase in the number of adolescents in these countries is associated with different health outcomes.

As per the 2018 population projection, Tanzania has a population of 60 million (National Bureau of Statistics [NBS], 2018). Out of them, 23% (approximately 12 million people) are adolescents aged 10–19 years, 16.7 million are children aged 0–9 years, and 29 million are non-adolescents (NBS, 2018). Recent efforts targeting adolescent empowerment have been made, but still, child marriage, early pregnancy, low quality of education, violence, abuse and exploitation continue to undermine some of their opportunities (United Republic of Tanzania [URT], 2011).

Almost one out of three girls in Tanzania gets married before her 18th birthday. By the age of 15, 5% of girls are married, and by the age of 18, 31% are married. Despite early marriage, one out of every three adolescent girls is anaemic and one out of every four is stunted (URT, 2020).

Approximately 15 million girls aged 15–19 years and 2.5 million girls under age 16 give birth each year in developing countries (UNFPA, 2015). The highest proportions of early childbearing are found in sub-Saharan Africa, where birth rates among adolescents reach over 102 births per 1000 girls aged 15–19 (UNICEF, 2019). This is higher compared to the global estimate of 44 births per 1000 girls aged 15–19 years (UNICEF, 2019). Adolescent birth rates are higher in rural areas than in urban areas among those with low education levels or living in poor households (Wado *et al.*,

2019). The average birth rate (ABR) among adolescents in rural areas with no education and in the poorest households is 1.8, 2.8 and 2.8 times higher than that of those who live in urban areas, have secondary or higher education, and are from the richest households, respectively (UNFPA, 2013). Policy enforcement has been weak in many African countries, creating a loophole for child marriages, which resulted in a low level of education and employment opportunities, early and unwanted pregnancies, poor offspring outcomes, increased maternal malnutrition and mortality (Girls Not brides [GNB], 2016).

Adolescent pregnancies are considered a global health problem that occurs in high-, middle-, and low-income countries. Adolescent pregnancies occur in marginalized communities, commonly driven by poverty and a lack of education and employment opportunities (WHO, 2018b). Adolescent pregnant girls face difficulties in sustaining their health status due to poor socio-economic and other demographic factors. Furthermore, girls are discriminated against and consequently face diminished opportunities and choices, are poorly educated, subjected to serious health risks, threatened by violence, and are prone to unintended pregnancies, resulting in early motherhood (United Nations [UN], 2011). Adolescent mothers face neglect from their family, school, workplace and community (Mangeli *et al.*, 2018). When compared to non-adolescent mothers in both high- and low-income countries, they frequently have insufficient resources, are less educated, and lack financial independence (Roux *et al.*, 2019). These circumstances are what cause adolescent pregnancies and deliveries to be accompanied by more risks as compared to non-adolescent mothers (Yussif *et al.*, 2017).

Young mothers compared to adult mothers are more likely to have low maternal weight gain and anemia during pregnancy due to competing needs for nutrients (Nabugoomu *et al.*, 2018). Due to poor nutrition outcomes, adolescent girls are likely to give birth to preterm or low-birth-weight babies with a high likelihood of malnutrition and death compared to mothers aged 20 years and above (Banke-Thomas *et al.*, 2017). Besides poor birth outcomes, pregnant adolescent girls are prone to hypertensive disorders, eclampsia, puerperal endometritis, systemic infections and cesarean delivery (Grønvik & Fossgard, 2018; Muche & Alene, 2018; Roos & Nickel, 2018).

Furthermore, the high rates of adolescent pregnancy contribute significantly to the high prevalence of malnutrition among adolescents themselves and their under-five children (Olodu *et al.*, 2019). Adolescent mothers are not only challenged by health risks, as described above, but are socially disadvantaged as well. The majority have to raise their babies as single parents because they are unable to complete their education and have limited capacity to secure a job and sustain a livelihood to support themselves and their children (Banke-Thomas *et al.*, 2017). Additionally, they lack

critical childcare knowledge, which contributes significantly to the raising of malnourished children and thus creating a vicious circle of malnutrition and poverty.

1.2 Statement of the Problem

Adolescent pregnancy is a global phenomenon with clearly known causes and serious health, social, and economic consequences for individuals, families and communities (WHO, 2019a). It is associated with high maternal and child morbidity and mortality, which eventually affects socioeconomic development (Odukogbe & Yalew, 2018; Sully & Mumah, 2019). Adolescent mothers have disproportionately high rates of poverty, food insecurity, social isolation, poor health, employment, inadequate access to health care services, and unfriendly youth service (Nabugoomu, 2018). Mustapha *et al.* (2018) noted that pregnant adolescents have difficulties accessing health services such as prenatal care, skilled attendants during birth, and prevention of mother-to-child transmission (PMTCT) services (Mustapha *et al.*, 2018).

Maternal demographic characteristics like age, education, social status, economic status, employment and living conditions have high impacts on both unborn babies and children who are under five years of age. Most of the studies done in Tanzania (Malisa, 2015; Adolf, 2014; Kiluvia, 2011; Uromi, 2014; Sik, 2015) explored an association between maternal socio-economic status and service utilization and their impacts on children's birth outcomes and nutrition status. This study intends to expand their work by exploring if the observed association is different between adolescent and non-adolescent mothers. Despite the overwhelming evidence of adolescent social and economic vulnerabilities, little is known about how maternal age contributes to low birth weight and childhood stunting in Tanzania.

1.3 Rationale of the Study

Adolescent pregnancies have become a public health challenge in Tanzania. From 2012 to 2015, adolescent pregnancies have increased from 23% to 27% in the country, with those living in the poorest conditions having a much higher prevalence (42% compared to those in the highest wealth quintile (13%) (UNICEF, 2017). The increase in teenage pregnancies has been associated with poor birth outcomes, complications during pregnancy, childhood malnutrition, and maternal and child deaths (WHO, 2019b). In Tanzania, 236 000 babies are born prematurely each year, and 11 500 children under five die due to preterm complications (Tanzania, 2017). Pregnancy during adolescence is a known risk factor for preterm birth (WHO, 2019b). Overall, it is estimated that 15% to 20% of all children born worldwide have low birth weight (LBW), representing more than 20 million births a year (WHO, 2014). The estimated prevalence of LBW in Sub-Saharan Africa is

13%, although 54% of all children are not weighed at birth (WHO, 2014). In Tanzania, the prevalence of LBW babies is 10%. Maternal age, Body Mass Index (BMI), Antenatal Care (ANC) attendances and place of residence are among the reported risk factors for LBW (Roux *et al.*, 2019; Nabugoomu *et al.*, 2018; Banke-thomas *et al.*, 2017).

On the other hand, nutrition indicators for children under five years of age in Tanzania have shown some improvement over recent years, but stunting is still widely prevalent. According to the 2015/16 Tanzania Demographic and Health Survey (TDHS) the prevalence of stunting among children under five was 34%, a decline from 42% in 2010 (TDHS, 2010), but is still unacceptably high compared to the average of middle-income countries, which is 22% (WHO, 2019). Besides, more than half of young women under the age of 19 are either pregnant or mothers. The perinatal mortality rate among young women under the age of 20 is 56 per 1000 pregnancies, which is significantly higher than the 39 per 1000 pregnancies in older women aged 20–29 years and the 32 per 1000 pregnancies in those aged 30-39 (Ministry of Health, 2015). Furthermore, the maternal mortality rate in Tanzania in 2015 was high (556 deaths per 100 000 live births), and it is likely that adolescent pregnancies contributed greatly to the burden (UNICEF, 2017). This study explored factors associated with LBW and stunting among adolescent and non-adolescent mothers using data collected as part of the Tanzania Demographic and Health Survey in 2015/2016.

1.4 Research Objectives

1.4.1 General Objectives

To compare the childhood stunting and low birth weight outcomes of under-five children born by adolescent and non-adolescent mothers and their association with maternal characteristics.

1.4.2 Specific Objectives

- To determine the association between mother's age (adolescent and non-adolescent mothers) and under-five stunting.
- (ii) To determine the association between mothers' age (adolescent and non-adolescent) and low-birth-weight.
- (iii) To determine maternal characteristics associated with under-five stunting and low birth weight.

1.5 Research Questions

- (i) What is the prevalence of low birth weight outcomes among babies born by adolescent and non-adolescent mothers?
- (ii) What is the prevalence of childhood stunting of under-five children born by adolescent and non-adolescent mothers?
- (iii) What are maternal socio-economic, demographic features and uptake of care associated with under-five stunting and low birth weight among adolescent and non-adolescent mothers?

1.6 Significance of the Study

This study enlightens the significance of adolescent reproductive health issues in Tanzania. The findings of this study indicated the vulnerability of adolescent mothers with respect to their health together with their born and unborn babies. For different reasons, the risk of low birth weight was found to be significantly higher among adolescent mothers as compared to non-adolescent mothers. LBW was also found to be a great risk factor for under-five stunting. As there are still many unpublished reports about the increase in adolescent pregnancy, especially during these times of the COVID-19 pandemic, this special group needs a lot of support and attention to avoid the unnecessary foreseen poor children's futures in terms of health and nutrition. We recommend the government, stakeholders, and communities as a whole hold hands together and protect this vulnerable group through offering emotional support and health education during pregnancies to improve immediate and later-life childbirth outcomes.

1.7 Delineation of the Study

The study had a sample size of 13 266 women from 2015/16TDHS-MIS but contained only 2904 adolescent and 10 362 non-adolescent mothers. Unequal sample size distribution can affect the statistical power of the study. Since the study was not done in the field, some important clues concerning the causal relationship between dependent factors and outcome might be missing. For example, the study revealed that male children were more stunted than females, but the reasons behind this were incomprehensive. Variability in birth weight information may introduce some bias when comparing birth weights by maternal age since only 64% of live births were measured for their weight in the five years before the survey. This may underestimate the unrecorded LBW from home deliveries, and hence the findings may require interpretation with consideration.

CHAPTER TWO

LITERATURE REVIEW

2.1 Overview

This Chapter provides an overview of maternal characteristics that are associated with low birth weight and stunting in under-five children. The section is presented to cover each research objective such as: To determine the association between mother's age (adolescent and non-adolescent mothers) and child-birth weights, to determine the association between mothers' age (adolescent and non-adolescent and non-adolescent) under-five stunting status and to determine maternal characteristics associated with low birth weight and under-five stunting status among adolescent and non-adolescent mothers

2.2 The Association between Mother's Age (Adolescent and Non-Adolescent Mothers) and Childbirth Outcomes

2.2.1 Adolescent versus Non-Adolescent Mothers and Low Birth Weight

A child born with a weight of less than 2500 g is considered a low birth weight regardless of the gestational age (WHO, 2012). Low birth weight is complex and includes preterm neonates (born before 37 weeks of gestation), small for gestational age neonates at term and the overlap between these two situations – preterm and small for gestational age neonates, who typically have the worst outcomes. Several studies have indicated an increased risk of low birth weight babies among adolescent girls but with limited comparison among different age groups (Demelash *et al.*, 2015; Endalamaw *et al.*, 2018; Desta *et al.*, 2020; Ratnasiri *et al.*, 2018; Awintuen *et al.*, 2018; Ngwira & Stanley, 2015; Kassa *et al.*, 2019). The increased likelihood of having a low birth weight is a result of competing growth and nutrition needs exerted during adolescent hood and pregnancy (Mahumud *et al.*, 2017).

A case-control study conducted by Demelash (2015) in Bale, Ethiopia with 129 cases (52 adolescents and 77 adult mothers) and 258 controls (56 adolescents and 202 adult mothers) found an increased chance of having LBW among adolescent mothers (AOR = 3.1; 95% CI (1.65-5.73); p = 0.001) compared to adult mothers. Although the study provided more evidential results, it has some weaknesses, including selection bias, as it considered only hospital delivery and didn't consider other factors such as diseases, preterm births, and multiple births, which could contribute to LBW as well as measurement and memory bias (Demelash *et al.*, 2015). Another case-control study conducted in Tigray, Ethiopia with 127 cases (46 adolescent and 81 adult mothers) and 254

controls reported an increased risk of LBW among adolescent mothers (AOR = 6.42 (95% CI = 1.93-21.42) as compared to adult mothers. Other factors related to low birth weight were having a low level of education, being a rural resident, absence of antenatal care follow-up and maternal BMI of less than 18 kg/m² (Desta *et al.*, 2020). Although several findings found a positive association between low maternal age and LBW, one study conducted in South Africa and involving 1073 women (23% adolescent mothers) found an increased risk of LBW among older women, i.e., those over 20 years of age (adjusted odds ratio [AOR] 12. 20, CI 3.90 to 38.02) compared to those 19 years of age (Tshotetsi *et al.*, 2019). The association found might be due to a small age gap and memory bias since the study was a case-control and did not collect information on the status of ANC attendance, which has an impact on birth weight.

Besides age, findings from most studies have shown that maternal socio-demographic factors and economic factors like wealth, age, level of education, occupation category, marital status and ANC attendance are associated with LBW. Adolescent mothers have a higher prevalence of LBW than adult mothers (Njim & Agbor, 2018; Tshotetsi *et al.*, 2019; DeMarco *et al.*, 2021).

2.2.2 Adolescents versus Non-Adolescents and Preterm Births

Each year, about 15 million babies are born prematurely, representing more than one in 10 of all babies born around the world. Besides that, over 1 million children die each year due to complications of preterm birth (global action report on preterm birth [GA], 2012). Common causes of preterm birth have been mentioned as multiple pregnancies, infections and chronic conditions such as diabetes and high blood pressure. However, often no cause is identified (WHO, 2019).

In one of the multi-country, facility-based, cross-sectional studies implemented from May 2010 to December 2011 by Ganchimeg (2014), which comprised 32179 adolescent and 92267 adult deliveries, they found a higher risk of preterm birth among adolescent mothers compared to adults (AOR = 1.60; 95%CI, 1.37-1.87). This study provided the clue that adolescent mothers are at a higher risk of delivering pre-term babies compared to non-adolescent mothers. Although this study gave out important clues, it was subjected to bias. Selection bias was the biggest study weakness as only city hospitals were selected to participate, leaving small hospitals located in rural and urban areas to carry out the study, which are the ones that have the most abundant numbers of adolescent and non-adolescent mothers' deliveries. The study was also subjected to information bias from participants since some of them were more likely to provide wrong responses (Ganchimeg & Ota, 2014). Almost the same odds of delivering preterm birth were reported in a cohort study by Kassa *et al.* (2019) comprised of 374 adolescent and 760 adult mothers (AOR = 1.65; 95%CI, 1.09-2.49).

Maternal age at first pregnancy, ANC attendance and preeclampsia were mentioned as the risk factors (Kassa *et al.*, 2019).

Vivid evidence was found in a randomized control clinical trial done by Mombo-ngoma *et al.* (2016) in four countries (Gabon, Benin, Mozambique and Tanzania) with infants' delivery numbers of 1223 (adolescent mothers) and 2877 (non-adolescent mothers) that took place between September 2009 and December 2013. The study found a non-significant increase in the odds of delivering preterm infants to adolescent mothers of 20 compared to adult mothers. Factors like BMI, ANC attendance, and literacy were found to not be significantly associated with preterm birth.

However, a cross-sectional study done in Kenya by Wagura *et al.* (2018) with 34 adolescents and 229 non-adolescent mothers in December, 2013 found that the odds of delivering preterm infants were significantly lower among adolescent mothers (OR = 0.236; 95% CI (0.054–1.001)) compared to non-adolescent mothers (OR = 0.834; 95% CI (0.272–2.555)). This study has weak evidence because it involved a low number of adolescents, i.e., 34, which can result in imprecise estimates. The results might also under or over represent the reality when it comes to adolescent mothers' population and their birth outcomes. Furthermore, there is a likelihood of information and selection biases since the study was conducted in a referral hospital only (Wagura *et al.*, 2018).

2.3 The Association between Mothers' Age (Adolescent and Non-Adolescent) and Underfive Stunting Status

2.3.1 Adolescent Mothers versus Non-Adolescent Mothers and Stunting among Under-five

Stunting, or being too short for one's age, is defined as a height for age that is less than minus 2 standard deviations of the World Health Organization (WHO) child growth reference standards (De Onis, 2006). It is a largely irreversible outcome of inadequate nutrition and repeated bouts of infection during the first 1000 days of a child's life. Stunting has long-term effects on individuals and societies, including diminished cognitive and physical development, reduced productive capacity, poor health and an increased risk of degenerative diseases such as diabetes (WHO, 2012). Childhood stunting is one of the most significant impediments to human development globally, affecting approximately 162 million children under the age of 5 years (WHO, 2012). Maternal age has been known to influence childhood stunting, with a high prevalence among adolescent mothers as compared to non-adolescent mothers. This fact has been reported in several studies (Wemakor *et al.*, 2018; Khan *et al.*, 2019; Maravilla *et al.*, 2020; Quarshie, 2014; Nguyen *et al.*, 2014; Branson, 2016). In a case–control study by Wemakor *et al.* (2018) involving 300 children aged 6–59 months born to teenage (n = 150) and adult (n = 150) mothers in Tamale Metropolis, Ghana, carried out

from April to June 2017, it was revealed that the children of teenage mothers, compared to those of adult mothers, were 8 times more likely to be stunted [Adjusted Odds Ratio (AOR) = 7.56; 95% confidence interval (CI) 4.20–13.63]. Although the study provided some light for further studies, it was subjected to anthropometric measurements and information bias. Also, this study failed to show a cause-effect relationship between nutrition outcomes and children's voices (Wemakor *et al.*, 2018).

According to Maravilla *et al.* (2020) the ongoing nutritional requirements of adolescent mothers due to puberty may deplete fetal nutrition, causing low birthweight, which they also found is strongly associated with stunting (Maravilla *et al.*, 2020). Like other studies discussed above, the study by Nguyen *et al.* (2014), which used the Bangladesh Demographic and Health Survey (1996–2014) and a sample size of 30 331 children, reported that children born to adolescent mothers had lower z-scores for height-for-age (mean difference: 064 SD), weight-for-age (045 SD), and a higher prevalence of stunting (18 percentage points [pp]) and underweight (12 pp) than children born to adult mothers. Adolescent mothers are at an age where they still need to provide for their growth and developmental needs. These mothers are also at risk of inadequately breastfeeding their infants due to low milk supply, which results in undernourished children (Khan *et al.*, 2019). Other maternal contributors to stunting include short stature, short birth spacing, and adolescent pregnancy, which interfere with nutrient availability to the fetus (owing to the competing demands of ongoing maternal growth) (WHO, 2012).

2.4 Maternal Factors Associated with Low Birth Weight and Under-five Stunting Status among Adolescent and Non-Adolescent Mothers

2.4.1 Maternal Education Levels and Birth Outcomes

Education enables women to make independent decisions and have good access to household resources that are important for better nutrition (Gizaw & Gebremedhin, 2018). Adolescent mothers may represent a particularly disadvantaged risk group characterized by low socioeconomic status, income, and level of education (Mekonnen *et al.*, 2019). Adolescents with high education expectations are likely to delay sexual experience and childbearing to meet those expectations. As a result, they enter into sexual affairs at an advanced age after they have gained enough knowledge and their reproductive systems are fully matured to prevent the likelihood of undesirable birth outcomes. Maternal education and parity are distal socioeconomic and reproductive factors associated with adverse pregnancy outcomes (Althabe *et al.*, 2015).

A strong association was found in a retrospective cross-sectional study done in Ghana which involved 78 adolescent mothers and 853 non-adolescent mothers by Mohammed *et al.* (2019) and assessed the relationship between the maternal level of education and the prevalence of LBW. The study revealed that mothers who attended or completed their secondary or higher education were 63% (95% CI 0.20–0.78) less likely to give birth to LBW infants when compared with uneducated mothers. However, this study recruited participants from the hospital, hence creating a likelihood of selection bias and less generalizability of the obtained findings (Mohammed *et al.*, 2019). Unlike the study conducted in Ghana, an institutional-based study conducted in East Gojjam Zone, Northwest Ethiopia with a total of 374 adolescents (15–19 years) and 760 adults (20–34 years) women, education was not a significant factor for LBW. The study revealed the rate of LBW was higher among adolescent mothers (AOR 2.14; 95% CI 1.36, 3.36, p = 0.001) as compared to nonadolescent mothers. Considering that only institutional women were involved in the study, this might have caused study bias because if those delivering outside of institutional settings were also taken into account, they could have influenced the results either positively or negatively (Kassa *et al.*, 2019).

2.4.2 Maternal Occupation and Birth Outcomes

The majority of developing countries are grappling with the growing challenge of an increasing number of women entering the labor force while balancing maternal roles (Ngwira & Mkandawire, 2003). On the other hand, there is a close link between maternal occupation and economic status. Women with good income tend to widen their food and decision choices, which in turn influences their health status and their families (Pampel *et al.*, 2010).

Different studies have reported the association between maternal occupation and low birth weight among adolescent and non-adolescent mothers. In a systematic review of a total of 30 studies having 55 085 participants by Endalamaw *et al.* (2018), it was reported that LBW among teenage mothers was due to a lack of fair jobs, which contributed to unplanned and/or unwanted pregnancies and less attention to nutritional value and healthcare services utilization as a result of low economic power. Endalamaw *et al.* (2018) found that the risk of infants born to mothers whose age was 20 years to have LBW was two times greater compared to those who were born to non-adolescent mothers. However, a case-control study conducted by Demelash *et al.* (2015), which consisted of 52 adolescent and 77 non-adolescent mothers, revealed that occupational status was not associated with LBW. Nevertheless, the study found that the odds of LBW (AOR = 3.1; 95% CI = 1.65-5.73) were higher among adolescent mothers as compared to non-adolescent mothers (Demelash *et al.*, 2015). In another study, it was reported that the wealthier group of adolescents were better engaged in maternity care compared to the coalescent ones (Gebremedhin *et al.*, 2015). Good socioeconomic status is an important factor for improving maternal health and the prevention of poor birth outcomes.

2.4.3 Maternal Place of Residence and Birth Outcomes

Maternal place of residence is one of the social determinants that influences the health status of mothers in many ways. Living in a place where there is adequate access to different social and environmental services does influence the health of mothers and their children. A study by Kassa *et al.* (2019) which comprised 374 adolescent mothers and 760 older mothers found an increased risk of low birth weight (LBW) among adolescents than in adult women (AOR 2.14; 95% CI, 1.36, 3.36, p = 0.001). The study further revealed that 29 (13.7%) of low-birth-weight babies were born to adolescent women from rural areas, while 10.7% were from adolescent women from urban areas, although the difference was not significant (Kassa *et al.*, 2019). Difficulties in accessing services like medical, health and nutrition information could be the key factors that are associated with poor birth outcomes. Inadequate antenatal care follow-ups and less use of health services are common challenges among adolescents living far from facilities (Kimario *et al.*, 2020).

2.4.4 Maternal Nutrition Status and Birth Outcomes

Overall, 10% of Tanzanian women of reproductive age (15–49 years) are considered undernourished, with a body mass index (BMI) of less than 18.5, with rural women being more affected than urban women (Ministry of Health [MoH], 2015). The proportion of pregnant women of reproductive age who were malnourished [mid-upper arm circumference (MUAC) 220 mm] was higher among adolescent girls aged 15 to 19 years than older women (Tanzania National Nutrition Survey [TNNS], 2018). The Tanzania Demographic and Health Survey also reported an increased prevalence of overweight/obesity among women of age 15–19 years, from 9% in 2010 to 11% in 2015. The triple burden of malnutrition (undernutrition, hidden hunger, and being overweight) has greater negative impacts on birth outcomes. Poor maternal nutrition is a risk factor for poor infant birth, health, and nutrition outcomes (URT, 2011). Several studies have reported the link between low maternal BMI and low birth weight among adolescent mothers compared to adult mothers living in low- and middle-income countries (Sananpanichkul *et al.*, 2015; Kurniati, 2019; Gebregzabiherher *et al.*, 2017; Bililign *et al.*, 2018; Togoobaatar & Ota, 2014; Nesara, 2018; Veghari, 2016; Dharmalingam & Navaneetham, 2009; Seid *et al.*, 2019).

A study conducted in Ethiopia by Kassa *et al.* (2019) with a total of 374 adolescents (15–19 years) and 760 adult women (20–34 years) reported an increased odd (2.14 times) of LBW among

adolescent women due to the low pregnancy weight gain (Kassa *et al.*, 2019). Inadequate weight gain during pregnancy is a risk factor for the delivery of LBW due to competing nutrients between mothers and unborn infants, and this is predominant among adolescent mothers (Gedefaw *et al.*, 2020; Mahumud *et al.*, 2017; Maravilla *et al.*, 2020; Togoobaatar & Ota, 2014).

However, a retrospective study by Sananpanichkul *et al.* (2015), which comprised 371 adolescents and 1641 non-adolescent mothers, conducted between January and June 2013 in Pakistan, found that low maternal BMI and teenage pregnancy were not significantly associated with LBW (Sananpanichkul & Rujirabanjerd, 2015).

2.4.5 Maternal Service Utilization and Birth Outcomes

Access to maternal services is a greater obstacle for young women aged 15–19 in Tanzania than for their older counterparts. It has been reported that the major barriers perceived by adolescents in accessing delivery health services include lack of money (40%), long distance to health facilities (38%), lack of transport (37%) and unfriendly services (14%) (MoH, 2015). Utilization of maternal health services (MHS) across the continuum of care, that is, antenatal, intrapartum (by skilled birth attendants) and postpartum care, is critical in reducing pregnancy-related morbidities, decreasing maternal mortality of adolescent mothers and improving birth outcomes, survival, quality of life and health of their babies (Banke-Thomas *et al.*, 2017). Unawareness of young women aged 15–19 about where to access health services has been reported as a barrier to utilization (MoH, 2015). According to Tanzania Demographic and Health Survey (TDHS, 2015), antenatal care becomes effective in the prevention of adverse pregnancy outcomes during early pregnancy and continues throughout (TDHS, 2015). A study by Carmo *et al.* (2019) reported that the use of ANC services and health facilities was positively associated with birth outcomes (Carmo *et al.*, 2019). However, an important observation from the study is the lack of equity in accessing these services.

One of the cross-sectional studies conducted by Atunyambe *et al.* (2008) with a total of 762 women (442 adolescents and 320 non-adolescents) found that adolescent mothers were significantly more disadvantaged in terms of health care seeking reproductive health services and faced more challenges during pregnancy and early motherhood compared to non-adolescent mothers. The study further reported that the mean number of times adolescent and adult mothers attended ANC was nearly the same (4.1 and 4.3, respectively) and found a strong association between adolescent mothers and less than four ANC attendances (OR = 1.52, 95% CI: 1.12-2.07). Contrary to this, the study by Tshotetsi *et al.* (2019) contained a sample of 1073 adolescent and non-adolescent mothers and found women who had fewer than 5 ANC visits were strongly associated with LBW (AOR

1.30, CI 1.06 to 1.61) as compared to non-adolescent mothers. This could be possible due to a lack of necessary care, information, and intervention that prevent the likely adverse birth outcome. Due to lack of adherence and unfriendly antenatal care, poor hospitality among health providers has been cited to be among the causes of adverse birth outcomes among adolescents.

A study by Mekonnen *et al.* (2019) reported that unfriendly maternal care contributes to poor utilization of maternal services and that the economic status of the adolescent mother is positively associated with the use of maternal health care. The study further showed that adolescents in the wealthier group were more engaged with maternity care services compared to the poor ones. The use of maternity care is critical given that young women are at higher risk of experiencing complications during pregnancy, childbirth, and the postpartum period, which may lead to an increased risk of maternal and child mortality or poor lifetime health care providers may have either positive or negative effects on their experience of accessing maternity care (Mekonnen *et al.*, 2019). For example, in Tanzania, women aged 15–19 years who completed the full phase of antenatal care from 2013 to 2018 had a 48% completion rate (WHO, 2019). Some of the reasons behind feeling shy are poor income and unfriendly health services from healthcare providers.

2.4.6 Maternal Occupation and Child Stunting

Maternal employment empowers women economically and socially and is in line with Sustainable Development Goal 8, which aims to promote economic growth and productive employment for all. Adolescent pregnancy is associated with increased school dropout rates and decreased educational achievement. Almost half of childbearing mothers fail to complete their high school education, thus increasing the probability of persistent economic and social disadvantage (Barnet *et al.*, 2004). Adolescent mothers who drop out of school and do not pursue higher education are limited in their opportunities (Mohr *et al.*, 2019).

Additionally, adolescent mothers were more likely to be involved in manual work than formal ones as they are less likely to complete secondary and high levels of education (Nguyen *et al.*, 2016). Female employment tends to increase total household income and also the part of the income that is controlled by women has a positive effect on child nutrition and health (Bethelhem, 2019). Mothers who do not work have a higher risk of stunted children compared to mothers who work due to having less time to care for their children. Adolescent mothers are less able to guarantee their children adequate food, access to safe water, and sanitary conditions, which in turn affects child growth (Wemakor *et al.*, 2018).

2.4.7 Maternal Education Level Child Stunting

Maternal education is a strong predictor of child stunting with some minimal attenuation of the association by other factors (Abuya *et al.*, 2012). Childhood stunting is more likely among mothers with no education or less than a secondary education than among mothers with at least a secondary education (Abuya *et al.*, 2012).

Most teenagers are unlikely to complete their secondary education due to early pregnancy. A study which recruited 30 331 women aged 15–49 by Phuong *et al.* (2020) highlighted that a low level of education among adolescent mothers was significantly associated with a high stunting rate among their under-five children (p=0.05) compared to adult mothers (Phuong *et al.*, 2020). The association between mothers' low level of education and an increased risk of child stunting was also reported by several others (Assefa *et al.*, 2013; Chirande *et al.*, 2015; Semali *et al.*, 2015). Educated mothers are more likely to be employed in a variety of fields, have more options, and have consistent food security in their home. However, a study by Agedew & Chane (2015) conducted in Ethiopia, comprised of 97 adolescents and 465 non-adolescent mothers, reported that having no formal education for mothers is negatively associated with the nutritional status of children (Agedew & Chane, 2015).

2.4.8 Maternal Place of Residence and Child Stunting

A residential area is one of the significant determinants of child undernutrition (Roux *et al.*, 2019). A study conducted in Tanzania found that children living in urban areas were less likely to be stunted compared to their counterparts living in rural areas (AOR = 0.56, 95%CI = 0.50-0.62, P = 0.001) (Sunguya *et al.*, 2019). This could be due to the fact that children whose mothers reside in urban settings have more access to nutrition information, which enables them to feed their children properly compared to their counterparts. Children whose parents live in rural areas are more likely to be short than those whose parents live in cities (Agedew & Chane, 2015). Possible explanations may be due to better-equipped urban healthcare systems and high access to nutritional information and healthcare services (Gebru *et al.*, 2019). In addition, a cross-sectional study that utilized the Rwanda demographic and health survey found stunting to be higher in rural settings than in urban settings (Habimana & Biracyaza, 2019). The study also found that children whose mothers were younger (under 25) (Habimana & Biracyaza, 2019). In Zambia, a study which utilized demographic and health survey data by Mzumara *et al.* (2018) found that stunting was more prevalent in children living in urban areas than those in rural areas.

were stated include decreased maternal-child contact time due to work schedules, short periods of breastfeeding, early cessation of breastfeeding and improper complementary foods, which have a negative effect on children's growth (Mzumara *et al.*, 2018).

2.4.9 Maternal Nutrition Status and Under-five Stunting

Maternal nutrition and health status are critical factors in early child development (Sarma et al., 2017). Inadequate maternal weight gain and poor dietary intake during pregnancy may result in intrauterine growth retardation, which has an impact on the nutrition status of a child in later life (Gaccioli & Lager, 2016). According to the WHO, the nutritional status of adolescents is of great importance to the health of their mothers and children (WHO, 2018d). In Tanzania, 3.4% of adolescents are wasted (MUAC 220 mm) (TNNS, 2018). Several studies have found a link between maternal nutrition status and child nutrition outcome (Berhe et al., 2019; Hall et al., 2020; Xun, 2019; Black et al., 2008). A cross-sectional study conducted in Ethiopia among 1104 under-five children and 1911 women aged 15–49 years reported an increased risk of stunting in children born from mothers who were less than 155 cm. Children whose mothers' height was less than 155 cm were two times more likely to be stunted (AOR = 2.37) and wasted (AOR = 2.17) compared to those whose height was above 155 cm. Maternal stature is a combined indicator of both genetic and environmental stress during early life and adolescence (Hall et al., 2020). Another study conducted by Dessie *et al.* (2019) with 7452 participants found children whose mothers were underweight had 1.20 times (AOR = 1.20; 95%CI: 1.06, 1.35) chances of being stunted as compared to children of mothers with normal nutritional status (Dessie et al., 2019). Low maternal body-mass index is associated with intrauterine growth restriction (Black et al., 2008) and poor nutritional status in pregnant women results in a reduced supply of nutrients to the fetus, thereby disrupting the process of organogenesis, growth and development (Fitriani et al., 2020).

CHAPTER THREE

MATERIALS AND METHODS

3.1 Data Source

This study utilized the 2015-16 Tanzania Demographic and Health Survey and Malaria Indicator Survey (2015-16 TDHS-MIS). The TDHS-MIS is a national survey done every five years to monitor and evaluate the population's health and nutrition programs. Specifically, the survey aims to provide up-to-date information on fertility and childhood mortality levels; fertility preferences; awareness, approval, and use of family planning methods; maternal and child health; and knowledge and attitudes toward HIV/AIDS and other sexually transmitted infections (STIs).

The TDHS-MIS 2015-16 is the most recent data with a sufficient sample of adolescent and nonadolescent mothers together with their children's information. It allows assessment of the association between mothers (adolescent and non-adolescent) with social-economic status, child health, and nutrition outcomes.

3.2 Research Design

The study used secondary data collected from the countrywide cross-sectional survey on population health and nutrition indicators, maternal characteristics, feeding practices, health care seeking behaviours, and child anthropometric measurements.

3.3 Target Population

The study used data for adolescents 15-19 and non-adolescent mothers aged 20-49 years and their corresponding children 0-59 months all over the country.

3.4 Sampling Procedures

The 2015 TDHS-MIS uses a two-stage stratified sampling design (TDHS, 2015). The urban and rural areas of each region formed a sampling stratum. A total of 59 sampling strata were created with each sample selected independently. Implicit stratification and proportional allocation were done at each lower-level administrative unit. The detailed sampling procedure has been documented in the 2015 DHS report (TDHS, 2015).

The 2015/2016 TDHS dataset was downloaded from the Demographic and Health Survey (DHS) website after completing the registration and fulfilment of the conditions related to the protection of sensitive data. Analysis was conducted in a sample of women who had a live birth five years preceding the survey using STATA version 14 software accounting for survey design.

3.5.1 Subjects and Sample Size

A total of 13 634 eligible women were identified from 1782 households sampled. Of interviewed women, 27% (2904) were adolescents aged 15-19 years and 73% (10 730) were non-adolescent mothers. Child anthropometric measurements were available in a total of 8852 women (600 adolescents and 8252 non-adolescent) and childbirth weight data was available in a total of 6386 women (459 adolescents and 5927 non-adolescent).

3.5.2 Categorization of Variables

Variables were categorized as per DHS guidelines with the normative category used as a reference group in regression analysis (De Onis, 2006). Maternal nutritional status was assessed using the BMI and classified as undernutrition (BMI 18.5), normal (> 18.5 BMI 25), overweight (> 25 BMI 30) and obese (BMI > 30). Stunting and childbirth weight of under-five children born by adolescents (15–19 years) and non-adolescent mothers (19–49 years) were considered primary and secondary outcomes, respectively. Manyeh *et al.* (2016) used the 2006 World Health Organization (WHO) growth standards reference points based on z-scores and 2 standard deviations (Manyeh *et al.*, 2016). A child weighed at birth below 2500 grams was termed a "low birth weight baby." Adolescent (15–19 years) and non-adolescent (20–29, 30-39 and 40–49 years) maternal ages were used. The household wealth index was developed based on the household's asset ownership using principal component analysis. The five equal categories (highest, fourth, middle, second, and lower wealth quintiles) used in the TDHS reports were further categorized into two groups (high and low socioeconomic status), with the first three being in the low socioeconomic status to reflect the actual Tanzanian condition.

3.5.3 Statistical Analysis

Frequencies and percentages were used to present results from descriptive analysis stratified by outcome variables. Univariate logistic regression models were fitted independently with stunting and birth weight as outcome variables while adjusting for socioeconomic and demographic characteristics such as age, height, weight (for both mother and child), maternal characteristics (education, place of residence), and uptake of maternal care. As anticipated, logistic regression analysis was not done separately for each maternal age group (adolescent and adult mothers) due to the small sample size, particularly in the adolescent population. A pooled analysis was done instead for each outcome variable, with maternal age as the main exposure. A forward selection procedure was applied during modelling, with variable selection based on the change in the exposure effect estimate. The procedure involved four main steps:

- (i) Descriptive analysis and preliminary investigations into the relationship between variables,
 with a focus on effect sizes and *p*-values at a 95% significant level.
- (ii) Selecting one variable at a time from a list of candidate variables (socioeconomic and demographic characteristics such as age, height, weight (for both mother and child), maternal characteristics (education, place of residence), and use of maternal care). From univariate analysis were then included in the model with and without adjustment of forced variables to help understand the effect of forced variables. The choice of the "best" predictor to be included in the model was then decided based on the change in the exposure effect estimate. Each time a new variable was added to the model, evidence of confounding and multicollinearity was assessed by comparing the effect estimates and standard errors between the "univariate" and "multivariate" models.
- (iii) Multivariable models were fitted by adding explanatory variables (socioeconomic and demographic characteristics such as age, height, weight (for both mother and child), maternal characteristics (education, place of residence), and uptake of maternal care) that were removed from the models in step "b" one at a time to help explore their effect when added to the model. Variables that resulted in positive changes in the mean square error were then included in the model. The process was repeated until all variables that provided precise estimates of exposure variables were selected.

3.6 Ethical Considerations

The protocol, study procedures, and questionnaires for DHS surveys were reviewed and approved by the international Institutional Review Board (IRB) and by the National Institute of Medical Research committee prior to the 2015 data collection. Before the interviews, written informed consent was obtained from all participants. A parent or guardian provided consent for the child or adolescent to be involved in the study. Interviews were performed as privately as possible with each eligible respondent interviewed in the absence of another person. A comparative analysis of determinants of low birth weight and stunting among under-five children of adolescent and nonadolescent mothers using the 2015/16 Tanzania demographic and health survey was again approved by the Ifakara Health Institute review board (IHI/IRB/No: 24/2020).

CHAPTER FOUR

RESULTS AND DISCUSSION

4.1 Results

4.1.1 Association between Mother's Age (Adolescent and Non-adolescent Mothers) and Under-five stunting

(i) Household Characteristics

A total of 13 360 and 1782 households in Tanzania and Zanzibar were selected for the interviews. The response rate for the interviewed households was very similar in both rural and urban residency (Table 1).

A total number of 13 634 (unweighted sample) eligible women were identified for individual interviews. Interviews were completed with 13 266 women, yielding a response rate of 97%, the number of eligible women interviewed with age 20-49 years was 10 362 and 2904 with age 15-19 years (weighed sub-sample).

Results	Tanz	ania Mainl	and	Zanzibar	Tanzania	
	Urban	Rural	Total			
Household interviews						
Household selected	3570	8008	11 578	1782	13 360	
Household occupied	3364	7639	11 003	1764	12 767	
Household interviewed	3265	7543	10 808	1755	12 563	
Household response rate	97.1	98.7	98.2	99.5	98.4	
Interviews with women age 15-19	1053	1755	2808	97	2904	
Interview with women age 20-49	3622	6432	10 055	307	10 362	

Table 1: Household and individual interviews

This table obtained from the DHS-MIS explain the distribution of interviews conducted per Tanzania mainland and Zanzibar

(ii) Maternal background characteristics

This study is aimed at identifying the association between maternal characteristics that influence birth outcome and under-five stunting. Factors like maternal level of education, occupation, marital status, wealth quintile, place of residence, service utilization and radio exposure were analyzed to determine their association with under-five birthweight and nutrition status.

About 59% of the adolescent and 63% of the non-adolescent mothers had attained primary education. Only 0.1% of the adolescent mothers and 1.8% of the non-adolescent mothers attained a higher learning education (college or university) (Fig.1). The proportion of women employed in the agriculture sector was 59% (age 15-19 years) and 55% (age 20-49), and the corresponding proportions of those working in unskilled manual labour were 18.2% and (22.6%), respectively. Few of them were employed in clerical and professional/technical/managerial work. The proportion of the study participants in the highest wealth quintile was 29.7% for adolescents and 26.4% for non-adolescent mothers. About 75% of adolescent mothers and 11.4% of non-adolescent mothers were never married. Some of them (2.3% of adolescents and 15.9% of older mothers) were divorced, separated or widowed. Only a few adolescent mothers (23.0% compared to 72.8% of non-adolescent mothers) were married or living with their partners. Most of the respondents (43.2% of teen mothers and 45.1% of older mothers) had access to listening to the radio at least once per week. However, 21.3% and 22.3% of adolescent and non-adolescent mothers, respectively, were not listening to the radio at all.



Figure 1: Maternal level of education

	Maternal Age			
Paramatar	15 – 19 years	20 years and above		
	N=2904	N=10 362		
	n (%)	n (%)		
Level of education				
No education	174 (6.0)	1773 (17.1)		
Primary education	1711 (58.9)	6500 (62.7)		
Secondary education	1017 (39.15)	1908 (18.4)		
higher	2 (0.1)	181 (1.8)		
Occupation				
Professional/technical/managerial	12 (1.0)	414 (4.8)		
Clerical	3 (0.1)	75 (1.0)		
Agricultural - self employed	835 (59.0)	4741 (55.0)		
Domestic services	173 (12.2)	581 (6.7)		
Sales and services	66 (4.7)	481 (5.6)		
skilled manual	69 (4.9)	386 (4.5)		
unskilled manual	257 (18.2)	1951 (22.6)		
Wealth quintile				
Low SES	1433 (49.4)	5415 (52.3)		
High SES	1471 (50.7)	4947 (47.7)		
Place of residence				
Urban	1083 (37.3)	3728 (36.0)		
Rural	1821 (62.7)	6636 (64.0)		
Marital status				
Never married	2170 (74.7)	1183 (11.4)		
Married or living together	668 (23.0)	7543 (72.8)		
Divorced/ Separated/ Widowed	66 (2.3)	1636 (15.8)		
Radio exposure				
not at all	619 (21.3)	2313 (22.3)		
less than once a week	1031 (35.5)	3381 (32.6)		
at least once a week	1254 (43.2)	4667 (45.1)		

Table 2: Socio-economic and Demographic Characteristics of Mothers

(iii) Anthropometric and Nutrition Status of Mothers

The study participants have an average weight of 57.5 kg for adolescent mothers and 66.3 kg for non-adolescent mothers with a mean height of 155.7cm and 156.8 cm respectively (Table 3).

Parameter	Mat 15 – 1	ternal Age 9 years	Difference Effect 20 years and above		
	Mean	95% CI	Mean	95% CI	
Weight (kg)	57.5	54.6-60.3	66.3	64.3-68.3	
Height (cm)	155.7	155.4-156.0	156.8	56.6-157.0	

 Table 3: Anthropometric and nutrition status of mothers

With regard to BMI, more adolescent mothers seemed to be underweight (16.4% CI 14.9–18.1) compared to non-adolescent mothers (6.5% C.I 5.9–7.1). The majority of the participants (71.9% of adolescents and 59.1% of non-adolescent mothers) had normal BMI. However, non-adolescent mothers were significantly affected by overweight and obesity compared to adolescent mothers (Fig. 2).





Antenatal Care attendance

Fifty-one per cent of the adolescent and 48% of non-adolescent mothers were found to attend Antenatal Care attendance (ANC) whereby 47.4% and 51.3% respectively had at least four visits. Few of them (1.6% of adolescents and 2.1 of non-adolescent mothers) did not attend ANC visits (Fig. 3).



Figure 3: Gestation age at 1stAntenatal Care Visits

4.1.2 Prevalence of Low Birth Weight in Children from Adolescent and Non-adolescent Mothers

Overall, 14.3% of adolescents and 6.3% of non-adolescent mothers gave birth to low-birth-weight babies (Table 4). Nineteen per cent of the adolescent mothers with no education had low-birth-weight babies, compared to 7.1% of the non-adolescent mothers. The urban community appears to have a higher proportion of mothers who gave birth to LBW babies, with 15.5% for adolescent mothers and 6.9% for non-adolescent mothers, when compared to those located in rural settings (13.7% and 6.0%). Twenty-two per cent of divorced, separated or widowed adolescent mothers gave birth to LBW babies, compared to 9.5% of non-adolescent mothers. Twenty-four per cent of the undernourished adolescent mothers gave birth to LBW babies, as compared to 10.4% of undernourished adult mothers (Fig. 4).



Elemente A.	I are hindl		at a t == a a	f the	at d		
HIGHTP 4.	LOW DIFFE	i weight	crarine a	M THE	STHAV	nartieit	nante
LIGUIC TO			status u		Scuur	par uch	Janus
					•/		

X 7 • 11	Adolesce	nt mothers (15-19 years.)	Non-Adolescent mothers (20+ years	
Variable	Ν	LBW children n (%)	Ν	LBW children n (%)
Overall	459	66 (14.3)	5927	376 (6.3)
Level of education				
No education	34	6 (19.1)	831	59 (7.1)
Primary education	353	52 (14.9)	3867	241 (6.2)
Secondary education or higher	71	7 (0.9)	1229	76 (6.2)
Wealth quintile				
Low SES	275	40 (14.4)	3016	186 (6.2)
High SES	184	26 (14.1)	2910	190 (6.5)
Place of residence				
Urban	148	23 (15.5)	2255	156 (6.9)
Rural	312	43 (13.7)	3672	220 (6.0)
Marital status				
Never married	128	19 (14.9)	330	25 (7.6)
Married or living together	287	37 (12.9)	4842	279 (5.8)
Divorced/ Separated/ Widowed	45	10 (22.0)	755	72 (9.5)
Maternal BMI				
Undernutrition	43	10 (23.9)	309	32 (10.4)
Normal	363	46 (12.7)	3646	232 (6.4)
Overweight or obese	52	9 (18.0)	1936	110 (5.7)
ANC visits				
Ever attended	200	31 (15.7)	1837	124 (6.7)

Table 4. Prevalence of Low Birth Weight in children from adolescent and non-adolescent

SES= Socio-economic status, BMI= Body Mass Index, ANC=Antenatal care, LBW =Low Birth Weight

218

N=Total number of children with birth weight data, n= Number of children with Low Birth Weight

Attended at least four visits

31 (14.3)

2500

106 (4.3)

4.1.3 Prevalence of Under-five Stunting, Wasting and Underweight

Generally, 31% of stunted and 10% of the severely stunted children under five years old were from adolescent mothers, while the corresponding proportions of the children from non-adolescent mothers were 34.2% and 11.5%, respectively. As with acute malnutrition, the proportion of wasting among children under five was very small in children of both adolescent (7.0%) and non-adolescent mothers (4.6%). Eleven per cent of the adolescent mother's children were underweight, as compared to 13.8% of the adult mother's children (Table 5). The table above gives a broad picture of under-five nutrition outcomes, but our study focused on stunting status among adolescent and non-adolescent mothers, where the difference between them is not significant.

	Maternal Age					
Parameter	15	– 19 years	20 years and above			
	n	%(95% CI)	Ν	%(95%CI)		
Child Nutrition Status						
Stunted						
Overall	183	30.6 (26.6-34.9)	2820	34.2 (32.6-35.8)		
Severe	59	9.9 (7.6-12.7)	945	11.5 (10.5-12.3)		
Wasted						
Overall	42	7.0 (4.9-9.9)	376	4.6 (4.1-5.1)		
Severe	13	2.2 (1.2-4.1)	97	1.2 (1.0-1.5)		
Underweight						
Overall	65	10.9 (8.3-14.1)	1138	13.8 (12.7-14.9)		
Severe	16	2.7 (1.5-4.6)	224	2.7 (2.3-3.2)		

Table 5: Prevalence of under-five stunting, wasting and underweight

4.1.4 Prevalence of stunting among under-five children according to maternal characteristics

Thirty-eight per cent of under-five children born by women who have not been to school or have a primary level of education were stunted. Over 30% of children born to mothers living in rural areas or in lower socioeconomic groups were stunted. The proportion of stunted children by marital status was very similar, ranging from 25.8% among children born to divorced or separated adolescent women to 32.3%. For non-adolescent women, it ranges from 32.2% among those never married to 39.4% among divorced or separated women. Thirty-four per cent of the under-five stunted children were born to adolescent mothers who were overweight or obese, as compared to 26.6% of adult mothers. Adult mothers who ever attended ANC visits had a high percentage (33.9%) of stunted children as compared to adolescent mothers (22.87%) as presented in Table 6.



Figure 5: Stunting prevalence according to maternal age

Table 6:	Prevalence of child stunting among adolescent and non-adolescent mothers by
	maternal characteristics (N=8852)

	Adolescent mothers		Non-Adolescent mothers (20+ vears)		
Variable	N (12	Stunted children n (%)	N	Stunted children n (%)	
Overall	600	183 (30.6)	8252	2820 (34.2)	
Level of education					
No education	79	30 (38.0)	1820	709 (38.9)	
Primary education	455	139 (30.6)	5258	1839 (35.0)	
Secondary education or higher	66	14 (21.5)	1174	272 (23.2)	
Wealth quintile					
Low SES	419	137 (32.8)	5403	2124 (39.3)	
High SES	182	46 (25.4)	2849	695 (24.4)	
Place of residence					
Urban	141	36 (25.8)	2148	532 (24.8)	
Rural	459	147 (32.0)	6104	2287 (37.5)	
Marital status					
Never married	141	38 (26.9)	333	107 (32.2)	
Married or living together	416	134 (32.3)	7042	2367 (33.6)	
Divorced/ Separated/ Widowed	43	11 (25.8)	878	345 (39.4)	
Nutrition Status (BMI)					
Undernutrition	53	16 (30.0)	552	220 (39.9)	
Normal	481	145 (30.2)	5528	2024 (36.6)	
Overweight or obese	65	22 (34.3)	2155	573 (26.6)	
ANC visits		· · ·		· · ·	
Ever attended	284	81 (28.7)	2914	989 (33.9)	
Attended at least four	260	76 (29.1)	3001	913 (30.4)	

SES= Socio-economic status, BMI= Body Mass Index, ANC=Antenatal care, N=Total number of children with anthropometric measurement, n= Number of stunted children

4.1.5 Predictors of Low Birth Weight

This analysis was conducted on a sample of 6385 women that had a live birth five years preceding the survey and weight data for their most recent child. Maternal age (adolescent and non-adolescent mothers) was considered the main exposure as explained in the method section. Overall, non-adolescent mothers have a reduced odds of giving birth to LBW babies when compared to adolescent mothers in both univariate and multivariate analyses. The odds of delivering LBW babies decreased from 0.41 (95% CI: 0.29–0.57) in crude analysis to 0.34 (95% CI: 0.22-0.50) in adjusted analysis (Table 3). Maternal BMI, marital status, and attending at least four ANC visits were significantly associated with LBW. The proportion of LBW babies was higher among mothers with primary education (66.5%), married or living with their partner (71.6%) and those with low socioeconomic status (51.1%) (Table 7).

Table 7: Factors associated with Low Birth Weight (N=6385)				
Variables	Ν	n (%)	OR (95% CI)	*AOR (95%CI)
Maternal age (Years)				
Adolescents (15-19)	459	66 (14.3)	1	1
Adults $(20 +)$	5927	376 (6.3)	0.41 (0.29-0.57)	0.34 (0.22-0.50)
Level of education				
No education	865	65 (14.8)	1	
Primary education	4220	294 (66.5)	0.91 (0.63-1.32)	
Secondary education or higher	1300	83 (18.7)	0.83 (0.55-1.26)	
Wealth quintile				
Low SES	3291	226 (51.1)	1	
High SES	3094	216 (48.9)	1.02 (0.78-1.33)	
Marital status				
Never married	458	44 (9.8)	1.62 (1.05-2.49)	1.23 (0.75-2.04)
Married or living together	5129	316 (71.6)	1	1
Divorced/ Separated/ Widowed	799	81 (18.4)	1.72 (1.23-2.42)	1.76 (1.24-2.50)
Place of residence				
Urban	2402	179 (40.4)	1	
Rural	3984	263 (59.6)	0.88 (0.67-1.16)	
Maternal BMI				
Undernutrition	352	42 (9.6)	1.84 (1.23-2.75)	2.29 (1.43-3.67)
Normal	4009	279(63.2)	1	1
Overweight or obese	1987	120 (27.2)	0.86 (0.64-1.14)	0.99 (0.71-1.38)
At least 4 ANC visits				
No	2037	155 (53.0)	1	1
Yes	2718	138 (47.0)	0.65 (0.50-0.83)	0.64 (0.49-0.83)
Child Sex				
Male	3266	197 (44.5)	1	1
Female	3120	245 (55.5)	1.33 (1.07-1.68)	1.31 (0.98-1.75)

N= Total number of respondents, n= Number of respondents with low-birth-weight babies, % =row percent, AOR=Adjusted odds ratio, OR=odds ratio, CI= 95 % Confidence interval, SES=Socioeconomic status, BMI=Body mass index, ANC=Antenatal care

* Adjusted for maternal age, marital status, maternal BMI, ANC attendance and child sex

4.1.6 Predictors of Stunting in Under-five Children

A total of 8852 women with live births in the five years preceding the survey who had their height and age measurement documented in the dataset were considered in this analysis. As for LBW analysis, maternal age was considered as the main exposure while controlling for other maternal factors, including age, education, wealth index, marital status, place of residence, occupation, nutrition status, and child characteristics (birth weight, sex, and age) in the model. Although 34.2% of children born by non-adolescent mothers were stunted compared to 30.6% of adolescent mothers, the adjusted analysis did not show any significant difference in childhood stunting between these two age groups; AOR: 0.97 (95% CI: 0.73-1.29) (Table 8).

In a multivariate analysis, socioeconomic status (SES) was significantly associated with childhood stunting (AOR = 0.69; 95% CI: 0.57-0.84), with children found in the higher socioeconomic group having a reduced odds of being stunted compared to those in the lower socioeconomic group. Children born with low birth weight were 2.4 times more likely to be stunted (95% CI: 1.80-3.20) compared to those born with normal birth weight. Female children were less likely to be stunted (AOR = 0.77; 95% CI: 0.67-0.89) compared to their male counterparts. Also, children whose age was beyond 1 year were 3 times more likely to be stunted as compared to those with less than one year (Table 8).

	Ν	n (%)	OR (95% CI)	*AOR (95% CI)
Maternal age (years)				
Adolescent (15-19)	600	183 (30.2)	1	1
Adults (20+)	8252	2820 (34.2)	1.18 (0.96-1.45)	0.97 (0.73-1.29)
Level of education				
No education	1900	739 (24.6)	1	1
Primary education	5712	1978 (65.9)	0.83 (0.73-0.94)	1.05 (0.85-1.30)
Secondary education or higher	1240	286 (9.5)	0.47 (0.38-0.58)	0.81 (0.61-1.08)
Wealth quintile				
Low SES	5822	2262 (75.3)	1	1
High SES	3031	742 (24.7)	0.51 (0.45-0.58)	0.69 (0.57-0.84)
Marital status				
Never married	475	145 (4.8)	0.87 (0.70-1.10)	1.25 (0.95-1.67)
Married or living together	7457	2501 (83.3)	1	1
Divorced/ Separated/ Widowed	921	357 (11.9)	1.25 (1.04-1.49)	1.19 (0.93-1.51)
Place of residence				
Urban	2289	569 (18.9)	1	1
Rural	6564	2435 (81.1)	1.78 (1.48-2.15)	1.21 (0.93-1.56)
Maternal BMI				
Undernutrition	605	236 (7.9)	1.13 (0.93-1.38)	1.14 (0.85-1.52)
Normal	6009	2169 (72.3)	1	1
Overweight or obese	2220	596 (19.9)	0.65 (0.56-0.75)	0.77 (0.64-0.92)
Child characteristics				
Birth weight				
Normal	5245	1568 (90.4)	1	1
Low birth weight	341	166 (9.6)	2.23 (1.67-2.98)	2.40 (1.80-3.20)
Sex				
Male	4490	1631 (54.3)	1	1
Female	4363	1372 (45.7)	0.80 (0.72-0.89)	0.77 (0.67-0.89)
Age (months)				
0-11	1989	321 (10.7)	1	1
12-23	2065	781 (26.0)	3.17 (2.66-3.76)	3.19 (2.49-4.08)
24-59	4798	1901 (63.3)	3.41 (2.94-3.96)	3.36 (2.76-4.11)

Table 8: Factors	associated wit	h stunting (N=8852)
-------------------------	----------------	---------------------

*Adjusted for the maternal level of education, wealth index, marital status, place of residence, occupation, BMI, childbirth weight, sex and age

4.2 Discussion

4.2.1 The association between Adolescent and Non-adolescent Mothers' Characteristics and Childbirth Weight

There is a higher prevalence of low birth weight (14.3%) among adolescent mothers compared to non-adolescent mothers (6.3%). The prevalence of low birth weight among adolescent mothers was

higher than the 10.32% reported in the TDHS (2015) and the Eastern and Southern Africa zone (11%) (WHO, 2014) but nearly equivalent to that reported globally (14.6%) (WHO, 2019d).

(i) Maternal age and Childbirth Weight

After adjusting for potential confounders, the findings showed that maternal age was significantly associated with LBW. Similar findings were reported in other studies (Manyeh *et al.*, 2016; Mombo-ngoma *et al.*, 2016; Abebe *et al.*, 2018). Adolescent mothers have a greater risk of delivering LBW infants compared to adult mothers. Several factors were said to predispose adolescents to giving birth to low birth weight children, including having an unplanned pregnancy, biological immaturity, and poor nutrition status during pregnancy (Manyeh *et al.*, 2016).

(ii) Maternal Attendance to Antenatal Care and Childbirth Weight

According to WHO guidelines (WHO, 2018), a minimum of four ANC visits are required for a safe pregnancy. Findings from this study have shown that mothers who attended at least four ANC visits had a lower risk of delivering LBW infants compared to those who had few visits. These findings are similar to those from a study conducted in India (Jogia & Lodhiya, 2018). Attendance at ANC is likely to reduce access to health education and other services like iron and folic acid supplements, antimalarial and deworming drugs, which contribute to the better health of the mother and unborn baby.

(iii) Maternal Marital Status and Childbirth Weight

The study found a significant increase in the risk of LBW among mothers who were divorced/separated or widowed, of which the majority were adolescents. This is parallel to the finding reported by Gizaw and Gebremedhin (2018) but in contrast to the studies conducted in Ethiopia (Talie *et al.*, 2019; Tshotetsi *et al.*, 2019), which found no significant association between marital status and LBW. Lack of social and economic support and stress among single and separated/divorced/widowed pregnant adolescent girls could be contributing factors to the delivery of LBW infants (Marimuthu *et al.*, 2018).

(iv) Maternal Nutrition Status and Childbirth Weight

There were significantly higher odds of LBW among undernourished mothers compared to those whose nutritional status was normal. Optimal nutrition is particularly critical for pregnant teens (URT, 2011). However, the prevalence of underweight was higher in adolescent mothers compared

to their older counterparts. These findings were similar to the ones reported through studies conducted in India (Sananpanichkul & Rujirabanjerd, 2015) and Ethiopia (Tyagi *et al.*, 2017). Factors like poor SES, unemployment, poor dietary intake, and poor weight gain during pregnancy are likely to contribute to the delivery of LBW infants.

4.2.2 The Association between Adolescent and Non-adolescent Maternal Characteristics and Stunting in Under-five Children

From this study, it was determined that the prevalence of stunting among under-five children was 33.9%, which, according to WHO standards (WHO, 2017), is still unacceptably high and much higher compared to 25% in some other developing countries (URT, 2019). Although 34.2% of children born by non-adolescent mothers were stunted compared to 30.6% of those born by adolescent mothers, the adjusted analysis did not show any significant difference between these two age groups, AOR: 0.97 (95% CI: 0.73–1.29).

These findings are similar to those reported in Ethiopia (Agedew & Chane, 2015) but contrary to those presented by Habyarimana *et al.* (2016) who used the Rwanda Demographic and Health Survey. The lack of a significant difference in stunting among children born to adolescent mothers compared to older women signifies that stunting is affected by several other factors beyond maternal age. In this study, maternal SES, BMI, child sex, birth weight, and age were found to be significantly associated with childhood stunting. Other studies have found that poor birth spacing, SES, and nutrition status of older mothers are the factors associated with childhood stunting (Mistry *et al.*, 2018). Improving maternal nutrition and health before and during pregnancy reduces childhood stunting and enhances maternal survival (AU, 2009). Other maternal factors that were associated with childhood stunting as generated from the multivariate analysis are discussed in the sections below:

(i) Maternal Socioeconomic Status and Childhood Stunting

A negative association was found between mothers' high socioeconomic status and childhood stunting. This is consistent with the studies done in Bangladesh, sub-Saharan Africa and Ethiopia (Kamal *et al.*, 2010; Yaya *et al.*, 2020; Birara, 2014). High household purchasing power, good sanitation, and access to health care services are among the issues that are affordable among families with high socioeconomic status, and they are contributing to good nutrition status among underfive children (Chirande *et al.*, 2015). Poor social-economic status interferes with household food

security and access to quality social and health services (Kalu & Etim, 2018; Kalkidan, 2019; Yaya *et al.*, 2020) and negatively affects the health and nutritional status.

(ii) Maternal nutrition status and childhood stunting

Maternal nutrition status was significantly associated with stunting in under-five children. Unlike the study conducted in Switzerland (Jornayvaz *et al.*, 2016), this study has found a lower risk of stunting in children born to obese women as compared to those whose mothers had normal BMI. These findings are similar to those reported in Ghana (Abubakari *et al.*, 2015), which found that infants of obese or overweight mothers were more stunted than those whose mothers had normal BMI. Obesity may be a reflection of good socioeconomic status in African settings.

4.2.3 Relationship between Stunting and Child's Birth Weight, Sex and Age

(i) Childbirth Weight and Stunting

This study found that children born with a normal weight (> 2.5 kg) were 2 times less likely to be stunted compared to those born with a low birth weight. It was further revealed that low birth weight (<2.5 kg) is a significant factor in childhood stunting. This is in line with the studies conducted in Rwanda, Indonesia and Ethiopia (Nshimyiryo *et al.*, 2019; Rachmi *et al.*, 2016, Silas *et al.*, 2018). The LBW is a predisposing factor for poor growth (Aryastami *et al.*, 2017) and is a reflection of intrauterine nutrient deprivation (Gaccioli & Lager, 2016).

(ii) Child sex and Stunting

In this study, under-five male children were found to be more stunted than females. The stunting in female children was significantly decreased by 22% compared to males. This agrees with studies conducted in Tanzania, Afghanistan and Ghana (Kejo *et al.*, 2018; Khan *et al.*, 2019; Ali *et al.*, 2017). A meta-analysis study in which data from 16 developing countries was used, reported a higher prevalence of stunting among boys (40%) than among girls (36%). In the analysis, boys from the household with the poorest social economic status were more likely to be stunted than girls in the same social-economic category (Wamani *et al.*, 2007). The justification for the disparities between boys and girls provided in a study conducted in the Mbeya region of the Tanzania mainland might support our differences. In that study, 60% of infants were stunted, and among them, more boys (65.3%) were stunted than girls (54.7%). Maternal preference for girls was highlighted as the reason for the disparities (McAuliffe *et al.*, 2011). Probably this might be related to favouring of girls during breastfeeding as reported in Nigeria whereby girls were two-fold likely to receive

exclusive breastfeeding than boys (Agho *et al.*, 2011) or early exposure to complementary foods among boys than girls as reported in Cote d'Ivoire and Senegal (Issaka *et al.*, 2015).

(iii) Child age and Stunting

This study found that stunting was significantly associated with child age. The odds of being stunted were high among children whose age was over 11 months compared to the younger ones. The odds of being stunted were high in children aged 1-2 and 2-5 years (AOR = 3.18) compared to those children younger than 1 year. These findings agree with another study conducted in Tanzania which found a positive association between child age and stunting (Makori, 2018) and are in line with TNNS (2018). Poor breastfeeding practices and improper complementary feeding are some of the factors associated with stunting among children aged over one year of age. In this study, stunting was more prevalent in children who were older than one year of age compared to the younger group. This chronic malnutrition may be due to long-term nutrient deprivation and inadequate care (Mzumara *et al.*, 2018).

CHAPTER FIVE

CONCLUSION AND RECOMMENDATION

5.1 Conclusion

After controlling for the possible confounders, this study found that adolescent mothers aged 15– 19 years were significantly at higher risk of delivering LBW babies compared to older mothers. With a national average low birth weight of 10% (TDHS, 2015), the prevalence of LBW was higher among the children of adolescent mothers (14% compared to those of non-adolescent mothers (6%). Other maternal factors that showed a significant association with LBW were frequency of ANC attendance, marital status, and BMI. Under-five stunting was independent of maternal age. The study found that non-adolescent mothers had a lower rate of under-five children who were stunted than adolescent mothers. However, after controlling for the possible confounding factors like maternal wealth index, maternal nutrition status, childbirth weight, sex and age of the child, the association was not statistically significant.

5.2 Recommendations

- (i) Along with ongoing government efforts to enforce charges against those who cause early pregnancies among school girls, there is a need to also support adolescent pregnant girls to access friendly health services and to return to school after delivery. Preventive and birth control methods must be taught in early classes to prevent unexpected pregnancies.
- (ii) This study showed the likelihood of stunting to be higher in male children than in females. Because of this, government policies, strategies, and guidelines may require specific interventions to address the needs of boys. This also needs further comprehensive research to check what causes under-five boys to be more stunted than girls.
- (iii) Nutrition interventions need to be prioritized in the regions with the highest number of stunted children and the higher prevalence of chronic malnutrition with consideration of the multi-causal nature of stunting and multi-sectoral collaboration.

REFERENCES

- Abebe, A. M., Fitie, G. W., Jember, D. A., Reda, M. M., & Wake, G. E. (2018). Teenage Pregnancy and Its Adverse Obstetric and Perinatal Outcomes at Lemlem Karl Hospital, Tigray, Ethiopia, 2018. *BioMed Research International*, 2020, 1–8.
- Abubakari, A., Kynast-wolf, G., & Jahn, A. (2015). Maternal Determinants of Birth Weight in Northern Ghana. *PLoS ONE*, *10*(8), 1–15. https://doi.org/10.1371/journal.pone.0135641
- Abuya, B. A., Ciera, J., & Kimani-murage, E. (2012). Effect of mother's education on child's nutritional status in the slums of Nairobi. *BMC Pediatrics*, 12(80), 1-10.
- Adolf, J. (2014). Socio-Economic Factors Affecting Adolescent Mothers' Struggles to Revive their Aspirations in Makete District, Tanzania. https://www.google.com
- Agedew, E., & Chane, T. (2015). Prevalence of Stunting among Children Aged 6 23 Months in Kemba Woreda, Southern Ethiopia: A Community Based Cross-Sectional Study. *Hindawi* Advances in Public, 2015, 1-7
- Ali, Z., Saaka, M., Adams, A., Kamwininaang, S. K., & Abizari, A. (2017). The effect of maternal and child factors on stunting, wasting and underweight among preschool children in Northern Ghana. *BMC Nutrition*, *3*, 1–13. https://doi.org/10.1186/s40795-017-0154-2
- Althabe, F., Moore, J. L., Gibbons, L., Berrueta, M., Goudar, S. S., Chomba, E., Derman, R. J., Patel, A., Saleem, S., Pasha, O., Esamai, F., Garces, A., Liechty, E. A., Hambidge, K. M., Krebs, N. F., Hibberd, P. L., Goldenberg, R. L., Koso-thomas, M., Carlo, W. A., ... Mcclure, E. M. (2015). Adverse maternal and perinatal outcomes in adolescent pregnancies: The Global Network's Maternal Newborn Health Registry study. *Reproductive Health*, *12*(2), 1-9. https://doi.org/10.1186/1742-4755-12-S2-S8
- Aryastami, N. K., Shankar, A., Kusumawardani, N., Besral, B., Jahari, A. B., & Achadi, E. (2017). Low birth weight was the most dominant predictor associated with stunting among children aged 12 – 23 months in Indonesia. *BMC Nutrition*, 3(16), 1-6. https:// doi. org/ 10. 1186/ s40795-017-0130-x
- AU. (2009). A snapshot of the nutrition situation in Africa acknowledgements. https://www.unicef. org/esaro/A_Snapshot_of_Nutrition_Situation_in_Africa.pdf

- Awintuen, I., Id, A., Kanmiki, E. W., Nonterah, A., Tediosi, F., Id, J. A., Welaga, P., Azongo, D., & Oduro, A. R. (2018). Socio-demographic determinants of low birth weight: Evidence from the Kassena-Nankana districts of the Upper East Region of Ghana. *PLoS ONE*, *13*(11), 1–10.
- Banke-thomas, O. E., Banke-thomas, A. O., & Ameh, C. A. (2017). Factors influencing utilisation of maternal health services by adolescent mothers in Low-and middle-income countrie: A systematic review. *BMC Pregnancy and Childbirth*, 2017, 1–14. https:// doi. org/ 10. 1186/ s12884 -017-1246-3
- Barnet, B., Arroyo, C., Devoe, M., & Duggan, A. K. (2004). Reduced School Dropout Rates Among Adolescent Mothers Receiving School-Based Prenatal Care. *Archives of Pediatrics & Adolescent Medicine*, 158(3), 262–268. https://doi.org/10.1001/ARCHPEDI.158.3.262
- Berhe, K., Seid, O., Gebremariam, Y., & Berhe, A. (2019). Risk factors of stunting (chronic undernutrition) of children aged 6 to 24 months in Mekelle City, Tigray Region, North Ethiopia: An unmatched case-control study. *Pone 14*(6), 1–11.
- Bethelhem, E. G., Esther, G., & Matin, Q. (2019). *Links between maternal employment and child nutrition in rural Tanzania*. https://www.google.com
- Bililign, N., Legesse, M., & Akibu, M. (2018). A Review of Low Birth Weight in Ethiopia: Socio-Demographic and Obstetric Risk Factors Abstract Magnitude of low birth weight. *Global Journal of Research and Review*, 5(1), 1–5. https://doi.org/10.21767/2393-8854.100033
- Birara, M. Y., Amsalu, F., & Bikes, D. (2014). Prevalence and Factors Associated with Stunting, Underweight and Wasting: A Community Based Cross-Sectional Study among Children Age 6-59 Months. *Journal of Nutritional Disorders & Therapy*, 4(2), 1-16. https:// doi. org/ 10. 4172/2161-0509.1000147
- Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., Onis, M., De Ezzati, M., Mathers, C., & Rivera, J. (2008). Series Maternal and Child Undernutrition: Maternal and child undernutrition: Global and regional exposures and health consequences. *The Lancet Series*, 2008, 5–22. https://doi.org/10.1016/S0140-6736(07)61690-0
- Branson, N. (2016). HHS Public Access. *Economic Development and Cultural Change*, 63(3), 589–616.
- Carmo, M., Id, G., Brito, M., & Costa, D. (2019). Determinants of maternal health care and birth

- Chirande, L., Charwe, D., Mbwana, H., Victor, R., Kimboka, S., Issaka, A. I., Baines, S. K., Dibley, M. J., & Agho, K. E. (2015). Determinants of stunting and severe stunting among under-fives in Tanzania: Evidence from the 2010 cross-sectional household survey. *BMC Pediatrics*, *15*(1), 1-13. https://doi.org/10.1186/s12887-015-0482-9
- De Onis, M. (2006). WHO Child Growth Standards based on length/height, weight and age. *Acta Paediatrica, International Journal of Paediatrics*, 95(450), 76–85. https:// doi. org/ 10. 1080/08035320500495548
- DeMarco, N., Twynstra, J., Ospina, M. B., Darrington, M., Whippey, C., & Seabrook, J. A. (2021). Prevalence of Low Birth Weight, Premature Birth, and Stillbirth Among Pregnant Adolescents in Canada: A Systematic Review and Meta-analysis. *Journal of Pediatric and Adolescent Gynecology*, 03, 1–8. https://doi.org/10.1016/j.jpag.2021.03.003
- Demelash, H., Motbainor, A., Nigatu, D., Gashaw, K., & Melese, A. (2015). Risk factors for low birth weight in Bale zone hospitals, South-East Ethiopia: A case-control study. *BMC Pregnancy and Childbirth*, 264 (2015), 1–10. https://doi.org/10.1186/s12884-015-0677-y
- Dessie, Z. B., Fentie, M., Abebe, Z., Ayele, T. A., & Muchie, K. F. (2019). Maternal characteristics and nutritional status among 6 – 59 months of children in Ethiopia: Further analysis of demographic and health survey. *BMC Pediatrics*, 19, 1–10. https://doi.org/10.1186/s12887-019-1459-x
- Desta, S. A., Damte, A., & Hailu, T. (2020). Maternal factors associated with low birth weight in public hospitals of Mekelle city, Ethiopia: A case-control study. *Italian Journal of Pediatrics*, 46, 1–9.
- Dharmalingam, A., & Navaneetham, K. (2009). Nutritional Status of Mothers and Low Birth Weight in India Nutritional Status of Mothers and Low Birth Weight in India. *Maternal and Child Health Journal*, 14, 290–298. https://doi.org/10.1007/s10995-009-0451-8
- Endalamaw, A., Engeda, E. H., Ekubagewargies, D. T., & Belay, G. M. (2018). Low birth weight and its associated factors in Ethiopia: A systematic review and meta-analysis. *Italian Journal of Pediatrics*, *6*, 1–12.
- Fitriani, H., Achmad, S. R., & Nurdiana, P. (2020). Risk Factors of Maternal Nutrition Status

During Pregnancy to Stunting in Toddlers Aged 12 – 59 Months. Jurnal Keperawatan Padjadjaran, 8(2), 183–191.

Global Action Report on Preterm Birth. (2012). Born Too Soon. http://www.google.com

- Gaccioli, F., & Lager, S. (2016). Placental Nutrient Transport and Intrauterine Growth Restriction. *Frontiers in Physiology*, 7, 1–8. https://doi.org/10.3389/fphys.2016.00040
- Ganchimeg, T., & Ota, E. (2014). Pregnancy and childbirth outcomes among adolescent mothers:
 A World Health Organization multicountry study Pregnancy and childbirth outcomes among adolescent mothers: A World Health Organization multicountry study. *International Journal of Obstetrics & Gynaecology*, 2014, 40–48. https://doi.org/10.1111/1471-0528.12630
- Gebregzabiherher, Y., Haftu, A., Weldemariam, S., & Gebrehiwet, H. (2017). The Prevalence and Risk Factors for Low Birth Weight among Term Newborns in Adwa General Hospital, Northern Ethiopia. *Hindawi Obstetrics and Gynecology International*, 2017, 1-8. https:// doi. org/ 10.1155/2017/2149156
- Gebremedhin, M., Ambaw, F., Admassu, E., & Berhane, H. (2015). Maternal associated factors of low birth weight: A hospital-based cross-sectional mixed study in Tigray, Northern Ethiopia. *BMC Pregnancy and Childbirth*, 2015, 1–8. https://doi.org/10.1186/s12884-015-0658-1
- Gebru, K. F., Haileselassie, W. M., Temesgen, A. H., Seid, A. O., & Mulugeta, B. A. (2019).
 Determinants of stunting among under-five children in Ethiopia: A multilevel mixed-effects analysis of 2016 Ethiopian demographic and health survey data. *BMC Pediatrics*, 19, 1–13.
- Gedefaw, G., Alemnew, B., & Demis, A. (2020). Adverse fetal outcomes and its associated factors in Ethiopia: A systematic review. *BMC Pediatrics*, 269 (2020), 1–12.
- Gizaw, B., & Gebremedhin, S. (2018). Factors associated with low birthweight in North Shewa zone, Central Ethiopia: A case-control study. *Italian Journal of Pediatrics*, 44, 1–9.
- GNB. (2016). The Role of Parliamentarians in ending Child Marriage. *African Journal Early Marriage in Africa Trends*, 2(7), 4–14.
- Grønvik, T., & Fossgard, I. (2018). Complications associated with adolescent childbearing in Sub-Saharan Africa: A systematic literature review and meta-analysis. *PLoS ONE*, *13*(9), 1–21.
- Habimana, S., & Biracyaza, E. (2019). Risk factors of stunting among children under 5 years of age

in the Eastern and Western Provinces of Rwanda: Analysis of Rwanda demographic and health survey 2014/2015. *Pediatric Health, Medicine and Therapeutics, 10,* 115–130. https://doi.org/10.2147/phmt.s222198

- Habyarimana, F., Zewotir, T., & Ramroop, S. (2016). Key determinants of malnutrition of children under five years of age in Rwanda: Simultaneous measurement of three anthropometric indices. *African Population Studies*, 30(2), 2328–2340.
- Hall, J., Walton, M., Ogtrop, F., Van-Guest, D., Black, K., & Beardsley, J. (2020). Factors influencing undernutrition among children under 5 years from growing communities in Bougainville. *BMJ Global Health*, 2020(5), 1–10. https://doi.org/10.1136/bmjgh-2020-002478
- Jogia, P. D., & Lodhiya, K. K. (2018). A cross-sectional study for utilisation of antenatal care services and its association to birth weight of babies in a tertiary care centre in Western India. *International Journal of Community Medicine and Public Health*, 5(8), 3519–3525.
- Jornayvaz, F. R., Vollenweider, P., Bochud, M., Mooser, V., Waeber, G., & Vidal, P. M. (2016). Low birth weight leads to obesity, diabetes and increased leptin levels in adults: The CoLaus study. *Cardiovascular Diabetology*, 2016(15), 1–10. https://doi.org/10.1186/s12933-016-0389-2
- Kalkidan, H. A., Waseyehun, A., & Rahma, A. (2019). Varying Relationship of Wealth of Households and Stunting of Under Five Children in Sub-Saharan Africa: A Meta-Analysis of Demographic Health Surveys. *Ecronicon Nutrition*, 14(8), 616–623.
- Kalu, R. E., & Etim, K. D. (2018). Factors associated with malnutrition among under-five children in developing countries: A review. *Global Journal of Pure and Applied Sciences*, 24(1), 69-74. https://doi.org/10.4314/gjpas.v24i1.8
- Kamal, S. M. M., Manaf, R. A., & Fire, P. (2010). Effects of Wealth on Nutritional Status of Preschool Children in Effects of Wealth on Nutritional Status of Pre-school Children in Bangladesh. *Malaysian Journal of Nutrition*, 16(2), 219–232.
- Kassa, G. M., Arowojolu, A. O., Odukogbe, A. A., & Yalew, A. W. (2018). Prevalence and determinants of adolescent pregnancy in Africa: A systematic review and Meta-analysis. *Reproductive Health*, 5 (195), 1–17.

- Kassa, G. M., Arowojolu, A. O., Odukogbe, A. A., & Yalew, A. W. (2019). Adverse neonatal outcomes of adolescent pregnancy in Northwest Ethiopia. *PloS One*, *14*(6), 1–20.
- Kebede, A. S., Muche, A. A., & Alene, A. G. (2018). Factors associated with adverse pregnancy outcome in Debre Tabor town, Northwest Ethiopia: A case-control study. *BMC Research Notes*, 820 (2018), 1–6. https://doi.org/10.1186/s13104-018-3932-2
- Kejo, D., Mosha, T. C. E., Petrucka, P., Martin, H., & Kimanya, M. E. (2018). Prevalence and predictors of undernutrition among under-five children in Arusha District, Tanzania. *Food Science & Nutrition*, 6, 2264–2272. https://doi.org/10.1002/fsn3.798
- Khan, S., Zaheer, S., & Safdar, N. F. (2019). Determinants of stunting, underweight and wasting among children < 5 years of age: Evidence from 2012-2013 Pakistan demographic and health survey. *BMC Public Health*, 19, 1–15.
- Kiluvia, F. S. (2011). Gender and sexuality notions as determinants of school pregnancies in Tanzania: A case of Iringa rural district. https://www.google.com
- Kimario, F. F., Festo, C., & Shabani, J. (2020). Determinants of Home Delivery among Women Aged 15-24 Years. *International Journal of Maternal and Child Health and AIDS*, 9(2), 191– 199. https://doi.org/10.21106/ijma.361
- Kurniati, H. F. (2019). Factors of Low Weight Babies Incidence on Teenage Pregnancy. https://www.google.com
- Mahumud, R. A., Sultana, M., & Sarker, A. R. (2017). Distribution and Determinants of Low Birth Weight in Developing Countries. *Journal of Preventive Medicine and Public Health*, 50, 18– 28.
- Makori, N., Kassim, N., Kinabo, J., & Matemu, A. (2018). Factors Associated with Stunting in Dodoma Region, Tanzania. African Journal of Food, Agriculture and Nutrition, 18(3), 13842– 13860. https://doi.org/10.18697/ajfand.83.17000
- Malisa, J. N. (2015). Factors contributing to teenage pregnancies in Tunduru district council. https://www.google.com
- Mangeli, M., Tirgari, B., Cheraghi, M. A., & Rayyani, M. (2018). *Electronic Physician*. https://www.google.com

- Manyeh, A. K., Kukula, V., Odonkor, G., Ekey, R. A., Adjei, A., Narh-bana, S., Akpakli, D. E., & Gyapong, M. (2016). Socioeconomic and demographic determinants of birth weight in southern rural Ghana: Evidence from Dodowa Health and Demographic Surveillance System. *BMC Pregnancy and Childbirth*, 160 (2016), 1–9. https://doi.org/10.1186/s12884-016-0956-2
- Maravilla, J. C., Betts, K., Adair, L., & Alati, R. (2020). Stunting of children under two from repeated pregnancy among young mothers. *Scientific Reports*, 14265 (2020), 1–9. https:// doi. org/ 10.1038/s41598-020-71106-7
- Marimuthu, Y., Sarkar, S., Sakthivel, M., Yuvaraj, K., Bharathnag, N., & Sarveswaran, G. (2018). Association of social factors with low birth weight: A narrative review. *International Journal* of Community Medicine And Public Health, 5(8), 1-5. https://bdoi.org/10.18203/2394-6040. ijcmph20183046
- Mekonnen, T., Dune, T., & Perz, J. (2019). Maternal health service utilisation of adolescent women in sub-Saharan Africa: A systematic scoping review. *BMC Pregnancy and Childbirth*, 6(19), 1–16.
- Mistry, S. K., Hossain, B., Khanam, F., Akter, F., Parvez, M., Yunus, F., Afsana, K., & Rahman, M. (2018). Individual-, maternal- and household-level factors associated with stunting among children aged 0 – 23 months in Bangladesh. *Public Health Nutrition*, 22(1), 85–94. https://doi. org/10.1017/S1368980018002926
- MoH. (2015). United Republic of Tanzania Ministry of Health and Social Welfare The National Road Map Strategic Plan To Accelerate Reduction of Maternal, Newborn and Child Deaths in Tanzania (Issue April 2008). https://www.google.com
- Mohammed, S., Bonsing, I., Yakubu, I., & Wondong, W. P. (2019). Maternal obstetric and sociodemographic determinants of low birth weight: A retrospective cross-sectional study in Ghana. *Reproductive Health*, 5(16), 1–8.
- Mohr, R., Carbajal, J., & Sharma, B. B. (2019). The Influence of Educational Attainment on Teenage Pregnancy in Low-Income Countries: A Systematic Literature Review. *Journal of Social Work in the Global Community*, 4(1), 19–31. https://doi.org/10.5590/JSWGC. 2019. 04. 1.02

- Mombo-ngoma, G., Mackanga, J. R., González, R., Ouedraogo, S., Kakolwa, M. A., Manego, R. Z., Basra, A., Rupérez, M., Cot, M., Kabanywany, A. M., Matsiegui, P., Agnandji, S. T., Vala, A., Massougbodji, A., Abdulla, S., Adegnika, A. A., Sevene, E., Macete, E., Yazdanbakhsh, M., ... Menéndez, C. (2016). Young adolescent girls are at high risk for adverse pregnancy outcomes in sub-Saharan Africa: An observational multicountry study. *BMJ Open*, 2016(6), 1-8. https://doi.org/10.1136/bmjopen-2016-011783
- Mustapha, M., Musiime, V., Bakeera-kitaka, S., Rujumba, J., & Nabukeera-barungi, N. (2018). Utilization of "prevention of mother-to-child transmission" of HIV services by adolescent and young mothers in Mulago Hospital, Uganda. https://ww.google.com
- Mzumara, B., Bwembya, P., Halwiindi, H., Mugode, R., & Banda, J. (2018). Factors associated with stunting among children below five years of age in Zambia: Evidence from the 2014 Zambia demographic and health survey. *BMC Nutrition*, *4*(51), 1–8.
- Nabugoomu, J. (2018). Adolescent Maternal Nutrition and Health in Uganda: Voices from the Community. https://www.google.com
- Nabugoomu, J., Seruwagi, G. K., Corbett, K., Kanyesigye, E., Horton, S., & Hanning, R. (2018). Needs and Barriers of Teen Mothers in Rural Eastern Uganda: Stakeholders' Perceptions Regarding Maternal / Child Nutrition and Health. https://doi.org/10.3390/ijerph15122776

NBS. (2018). National Population Projections of Tanzania. https://www.google.com

- Nesara, P. (2018). *Determinants of Low Birth Weight in a Population-Based Sample of Zimbabwe*. https://www.google.com
- Nguyen, P. H., Scott, S., Neupane, S., Tran, L. M., & Menon, P. (2016). Social, biological and programmatic factors linking adolescent pregnancy and early childhood undernutrition: A path analysis of India's 2016 National Family and Health Survey. *The Lancet Child and Adolescent Health*, 3(7), 463–473. https://doi.org/10.1016/S2352-4642(19)30110-5
- Ngwira, A., & Stanley, C. C. (2015). Determinants of Low Birth Weight in Malawi: Bayesian Geo-Additive Modelling. *PLoS Medicine*, *10*(6), 1–14. https:// doi. org/ 10. 1371/ journal. pone. 0130057
- Ngwira, N., & Mkandawire, E. (2003). Cost of Gender Disparities In Access To Socio-Economic Services. https://www.google.com

- Njim, T., & Agbor, V. N. (2018). Adolescent deliveries in rural Cameroon: Comparison of delivery outcomes between primipara and multipara adolescents. *BMC Research Notes*, *11*(1), 4–9. https://doi.org/10.1186/s13104-018-3550-z
- Nshimyiryo, A., Hedt-gauthier, B., Mutaganzwa, C., Kirk, C. M., Beck, K., Ndayisaba, A., Mubiligi, J., Kateera, F., & El-khatib, Z. (2019). Risk factors for stunting among children under five years: A cross-sectional population-based study in Rwanda using the 2015 Demographic and Health Survey. *BMC Public Health*, 19, 1–10.
- Olodu, M. D., Adeyemi, A. G., Olowookere, S. A., & Esimai, O. A. (2019). Nutritional status of under-five children born to teenage mothers in an urban setting, southwestern Nigeria. *BMC Research Notes*, 116 (2019), 1–6. https://doi.org/10.1186/s13104-019-4147-x
- Pampel, F., Patrick, K., & Justin, D. (2010). Socioeconomic Disparities in Health Behaviors. Annual Review of Sociology, 36, 349–370. https:// doi. org/ 10. 1146/ annurev. soc. 012809. 102529. Socioeconomic
- Phuong, N., Samuel, S., Long, K. P., Pramanik, A. A., Kaosar, A., & urnima, M. (2020). Why Are Adolescent Mothers More likely to Have Stunted. *Nutritional Epidemiology*, 2020, 1463.
- Quarshie, E. M. (2014). Comparative Nutritional Status of Adult and Adolescent Mothers and their Infants. http://ugspace.ug.edu.ghhttp//ugspace.ug.edu.gh
- Rachmi, C. N., Agho, K. E., Li, M., & Baur, L. A. (2016). Stunting, Underweight and Overweight in Children Aged 2 . 0 – 4 . 9 Years in Indonesia: Prevalence Trends and Associated Risk Factors. *Plos One*, 11(5), 1–17. https://doi.org/10.1371/journal.pone.0154756
- Ratnasiri, A. W. G., Parry, S. S., Arief, V. N., Delacy, I. H., Halliday, L. A., Dilibero, R. J., & Basford, K. E. (2018). Recent trends, risk factors, and disparities in low birth weight in California, 2005 2014: A retrospective study. *Maternal Health, Neonatology, and Perinatology*, 4, 1–13.
- Roux, K., Christodoulou, J., Stansert-katzen, L., Dippenaar, E., Laurenzi, C., Roux, I. M., Tomlinson, M., & Rotheram-borus, M. J. (2019). A longitudinal cohort study of rural adolescent vs adult South African mothers and their children from birth to 24 months. *BMC Pregnancy and Childbirth*, 8, 1–8.

Sananpanichkul, P., & Rujirabanjerd, S. (2015). Association between maternal body mass index

and weight gain with low birth weight in Eastern Thailand. Southeast Asian Journal of Tropical Medicine and Public Health, 46(6), 1085–1091.

- Sarma, H., Khan, J. R., Asaduzzaman, M., Uddin, F., Tarannum, S., & Hasan, M. (2017). Factors Influencing the Prevalence of Stunting among Children Aged Below Five Years in Bangladesh. *Food and Nutrition Bulletin*, 38(3), 291–301. https:// doi. org/ 10. 1177/ 0379572117710103
- Seid, S. S, Tolosa, T., & Adugna, D. (2019). Prevalence of Low Birth Weight and Associated Factor among neonate Born at Jimma Medical Center. *IMedPub Journals*, 10(1), 1–4. https://doi.org/10.21767/2172-0479.100156
- Sik, I. (2015). Early Motherhood in Tanzania: Exploring the Education, Health and Employment Challenges of Dar es Salaam's Adolescent Mothers. https://www.google.com
- Silas, L., Rantetampang, A. L., Tingginehe, R., & Mallongi, A. (2018). The Factors Affecting Stunting Child under Five Years in Sub Province Mimika. *International Journal of Science* and Healthcare Research, 3(2), 99–108.
- Sunguya, B. F., Zhu, S., Mpembeni, R., & Huang, J. (2019). Trends in prevalence and determinants of stunting in Tanzania: An analysis of Tanzania demographic health surveys (1991 – 2016). *Nutrition Journal*, 2019, 1–13.
- Talie, A., Taddele, M., & Alemayehu, M. (2019). The magnitude of Low Birth Weight and Associated Factors among Newborns Delivered in Dangla Primary Hospital, Amhara Regional State, Northwest Ethiopia, 2017. *Hindawi Journal of Pregnancy*, 2019, 1-7. https://doi.org/10.1155/2019/3587239
- Tanzania. (2017). Profile of preterm and low birth weight prevention and care. https:// www. google. com
- TDHS. (2015). Tanzania Demographic and Health Survey (TDHS). https://www.google.com
- TNNS. (2018). Tanzania National Nutrition Survey 2018 (Issue June). https://www.google.com
- Tshotetsi, L., Dzikiti, L., Hajison, P., & Id, S. F. (2019). Maternal factors contributing to low birth weight deliveries in Tshwane District, South Africa. *PloS One*, *14*(3), 1–13.
- Tyagi, S., Gurudayal, S., & Neena, B. (2017). Maternal Nutritional Status and Its Relation with

Birth Weight. International Journal of Health Sciences and Research, 7(8), 422–433.

UN. (2011). Fact Sheet: Girls and Young Women. https://www.google.com

UNFPA. (2013). Adolescent pregnancy: A Review of the Evidence. https://www.google.com

UNFPA. (2015). Facing the facts Adolescent girls and contraception. https://www.google.com

- UNICEF. (n.d.). Adolescent Demographics UNICEF DATA. Retrieved October 29, 2019, from https://data.unicef.org/topic/adolescents/demographics/
- UNICEF. (2017). Maternal and child health. https://www.google.com
- UNICEF. (2019). *Early Childbearing UNICEF DATA*. https://data.unicef.org/topic/child-health/adolescent-health/
- Uromi, S. M. (2014). *Schoolgirl pregnancies as a most critical and rapidly growing challenge in Tanzania*. https://www.google.com
- URT. (2011). Adolescence in Tanzania. https://www.google.com
- URT. (2019). United Republic of Tanzania Country overview Malnutrition burden Stunting at the subnational level. https://www.google.com
- URT. (2020). The 2019 School Malaria and Nutrition Survey (SMNS) Report. https://www.google.com
- Veghari, G. (2016). Maternal age and BMI in relation to infant birth size: A study in public health centres in the north of Iran. *International Journal of Medical Research & Health Sciences*, 5(9), 151–155.
- Wado, Y. D., Sully, E. A., & Mumah, J. N. (2019). Pregnancy and early motherhood among adolescents in five East African countries: A multi-level analysis of risk and protective factors. *BMC Pregnancy and Childbirth*, 59 (2019),1–11.
- Wagura, P., Wasunna, A., Laving, A., Wamalwa, D., & Ng, P. (2018). Prevalence and factors associated with preterm birth at Kenyatta national hospital. *BMC Pregnancy and Childbirth*, 18, 2–9.
- Wall-wieler, E., Roos, L. L., & Nickel, N. C. (2018). Adolescent Pregnancy Outcomes Among

Sisters and Mothers: A Population-Based Retrospective Cohort Study Using Linkable Administrative Data. *Public Health Reports*, *133*(1), 100–108. https:// doi. org/ 10. 1177/ 0033354917739583

- Wemakor, A., Garti, H., Azongo, T., Garti, H., & Atosona, A. (2018). Young maternal age is a risk factor for child undernutrition in Tamale Metropolis, Ghana. *BMC Research Notes*, 11, 1–5. https://doi.org/10.1186/s13104-018-3980-7
- WHO. (n.d.). Preterm birth. https://www.google.com
- WHO. (2012a). WHA Global Nutrition Targets 2025: Low Birth Weight Policy Brief. https:// www. google.com
- WHO. (2012b). WHA Global Nutrition Targets 2025: Stunting Policy Brief. https:// www. google.com
- WHO. (2014). WHA Global Nutrition Targets 2025: Low Birth Weight Policy Brief. https:// www. google. com
- WHO. (2017). Nutrition in the who African region. http://apps.who.int/bookorders.
- WHO. (2018a). 2016 WHO Antenatal Care Guidelines Malaria in Pregnancy Frequently Asked Questions (FAQ). https://www.google.com
- WHO. (2018b). *Adolescent pregnancy*. https://www.who.int/news-room/fact-sheets/ detail/ adolescent-pregnancy
- WHO. (2018c). Handout for Module A Introduction. In *Department of Child and Adolescent Health* and Development. https://www.google.com
- WHO. (2018d). *Promoting proper feeding for infants and young children*. https:// www. who. int/ nutrition/ topics/infantfeeding/en/
- WHO. (2019a). Adolescent pregnancy evidence brief. https://www.google.com

WHO. (2019b). Children, food and nutrition. https://www.google.com

WHO. (2019c). Levels and trends in child malnutrition. https://www.google.com

WHO. (2019d). Low birthweight estimates. https://www.google.com

- Xun, N. (2019). Role of Maternal Nutrition in the Health Outcomes of Mothers and their Children: A Retrospective Analysis. *Maternal Nutrition and Health Outcomes*, 25, 4430–4437. https:// doi.org/10.12659/MSM.914679
- Yaya, S., Oladimeji, O., Kolawole, E., & Bishwajit, G. (2020). Household structure, maternal characteristics and children' s stunting in Sub-Saharan Africa: Evidence from 35 countries. *International Health*, 00, 1–9. https://doi.org/10.1093/inthealth/ihz105
- Yussif, A. S., Lassey, A., Ganyaglo, G. Y. K., Kantelhardt, E. J., & Kielstein, H. (2017). The longterm effects of adolescent pregnancies in a community in Northern Ghana on subsequent pregnancies and births of the young mothers. *Reproductive Health*, 14(1), 1–7. https:// doi. org/ 10. 1186/S12978-017-0443-X/FIGURES/2

RESEARCH OUTPUTS

(i) **Publication**

Mtongwa, H. R., Festo, C., & Elisaria, E. (2021). A Comparative Analysis of Determinants of Low Birth Weight and Stunting Among Under-Five Children of Adolescent and Non-Adolescent Mothers Using 2015/16 Tanzania Demographic and Health Survey. *BMC Nutrition*, 7(64), 1-10. https://doi.org/10.1186/s40795-021-00468-6

(ii) **Poster Presentation**