The Nelson Mandela AFrican Institution of Science and Technology

https://dspace.mm-aist.ac.tz

Life sciences and Bio-engineering

Research Articles [LISBE]

2022-04-19

Spatial and seasonal group size variation of wild mammalian herbivores in multiple use landscapes of the Ngorongoro Conservation Area, Tanzania

Leweri, Cecilia

PLOS ONE

https://doi.org/10.1371/journal.pone.0267082 Provided with love from The Nelson Mandela African Institution of Science and Technology



G OPEN ACCESS

Citation: Leweri CM, Bartzke GS, Msuha MJ, Treydte AC (2022) Spatial and seasonal group size variation of wild mammalian herbivores in multiple use landscapes of the Ngorongoro Conservation Area, Tanzania. PLoS ONE 17(4): e0267082. https://doi.org/10.1371/journal.pone.0267082

Editor: Bi-Song Yue, Sichuan University, CHINA

Received: July 29, 2021

Accepted: April 2, 2022

Published: April 19, 2022

Copyright: © 2022 Leweri et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting information files.

Funding: CL received the grants from the African Development Bank and additional funds and research equipment from the Orskov foundation and IdeaWild. African Development Bank (AfDB Grant number 210015503281) Orskov foundation https://www.orskovfoundation.org/ IdeaWild https://ideawild.org/ No - The funders had no role in study design, data collection and analysis, **RESEARCH ARTICLE**

Spatial and seasonal group size variation of wild mammalian herbivores in multiple use landscapes of the Ngorongoro Conservation Area, Tanzania

Cecilia M. Leweri ^{1,2}*, Gundula S. Bartzke³, Maurus J. Msuha⁴, Anna C. Treydte^{1,5,6}

 Department of Sustainable Agriculture, Biodiversity and Ecosystem Management, School of Life Sciences and Bio-Engineering, The Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania,
Wildlife Information and Education Unit, Tanzania Wildlife Research Institute (TAWIRI), Arusha, Tanzania,
Institute of Crop Science, Biostatistics, University of Hohenheim, Stuttgart, Germany, 4 Wildlife Division, Ministry of Natural Resources and Tourism, Dodoma, Tanzania, 5 Hans Ruthenberg Institute, Ecology of Tropical Agricultural Systems, University of Hohenheim, Stuttgart, Germany, 6 Department of Physical Geography, Stockholm University, Stockholm, Sweden

* cecilia.leweri@tawiri.or.tz

Abstract

Group sizes of wild herbivores can be indicators of ecosystem health and proxies for individual and population fitness, particularly in areas where human activities have become increasingly common. We recorded 176 single- and multi-species groups of wild herbivores in human-dominated landscapes of the Ngorongoro Conservation Area (NCA) during dry and wet seasons. We analyzed how wild herbivore group sizes were affected by: (1) season, (2) distance to fully protected area (NCA crater) and to streams, (3) distance to human settlements, and (4) numbers of livestock. Group sizes were generally larger during the wet season than during the dry season and varied seasonally with distance to NCA crater, streams, and human settlements. During the wet season, larger groups were observed further away from the NCA crater whereas the opposite pattern was apparent during the dry season. Average wild herbivore group sizes increased by about three-fold with increasing distance from the streams during the dry season but were invariant to streams during the wet season. Furthermore, during the dry season, group sizes were larger close to settlements but varied little with distance to settlements during the wet season. While livestock presence did not directly affect wild herbivore group size, distance to settlements, streams and distance to the Ngorongoro crater in interaction with rainfall seasonality did. We conclude that the NCA crater functions as a key resource area for wild herbivores such as wildebeest (Connochaetes taurinus) and zebra (Equus quagga burchelli) during the dry season, highlighting the need for its full protection status in this Man and Biosphere reserve.

decision to publish, or preparation of the manuscript.

Competing interests: NO authors have competing interests The authors have declared that no competing interests exist.

Introduction

Group size and composition are the most basic elements of social organization for ungulates living in herds as they influence both foraging, migration, and other daily activities [1]. Theoretical frameworks explaining variation in group size assume that there is a trade-off between fitness relevant costs and benefits, and that individuals maintain membership in groups of optimal sizes to maximize fitness [2, 3]. In some African ungulates, for example, group sizes and their spatial distributions vary temporarily with season as rainfall governs the quantity and quality of vegetation [4, 5]. Despite the significance of long-term population monitoring and studies on population dynamics and movements [6-8], the latter have not yet addressed the effects of human presence and livestock grazing on herbivore group sizes in pastoral and protected areas for conservation planning [9].

Humans and livestock increase pressure on rangelands, and add to the complexity of their management, especially in areas where wild herbivores strongly interact with livestock on a daily basis [10, 11]. Interactions between livestock and wildlife can be either competitive or facilitative, depending on the species involved, and on the seasonal availability of resources [12, 13]. For example, wild herbivores have coexisted with domestic herbivores in few subsistence pastoral systems with abundant water points [13, 14]. However, high livestock densities can also outcompete wild herbivores [15] and reduce wild herbivore group sizes [9, 16] or lead to long-term declines in the abundance and diversity of native wildlife [17, 18]. Competition often occurs during the dry season, when grazing ranges are constricted near available water resources and when overall fodder quality is lower than during the wet season [16]. A consecutive group size reduction may impact reproductive fitness of wild herbivores and, hence, their population dynamics [2, 19]. With an increasing growth of human and livestock population, there is a pressing need for research concerning the ecology and management of wild herbivores, particularly their group sizes and behavior in response to human presence and to changing environmental conditions.

Landscape features such as elevation, seasonal streams and natural vegetation could also alter wild herbivores group sizes [20–22] since they determine the quality and structure of habitats and likely the distribution of food patches [19, 23]. Surface water for example is well documented to attract wild animals [15, 24]. Elevation is another variable that can alter herbivore group sizes as plant biomass often decreases with elevation [25, 26]. Understanding how different landscape features influence group sizes can therefore inform decisions in species conservation and management, such as identifying conservation areas [27–29].

Various studies on herbivores analyzed the long-term population trends and assessed the seasonal stability of wild herbivore communities in the Ngorongoro Conservation Area (NCA) [30]. In addition, some studies found that the exclusion of resident pastoralists and their livestock from the Ngorongoro crater resulted into both increasing and decreasing trends of some ungulate species whereas overall, wildlife biomass remained unchanged [31, 32]. However, as humans continue to impact natural habitats and livestock populations still grow within NCA, there is an increasing need to understand how this affects distributions and group sizes of wild herbivore group sizes respond to environmental and anthropogenic variables such as livestock keeping, and human settlements in the NCA, and whether this response varies between season.

We hypothesized that larger wildlife groups will be formed during the wet season than during the dry season, i.e., when forage is abundant. We also expected that larger groups will be formed closer to the streams due to the higher availability of water and food compared with areas further away from streams [35]. Furthermore, we expected that groups will be larger in areas of low competition with livestock herds. In addition, we hypothesized that wildlife groups will be larger further away from settlements (houses and livestock enclosure or temporary corrals, in which livestock are herded at night to protect them against predators).

Material and methods

Study area

We conducted our study in four wards of the Ngorongoro Conservation Area (NCA), a UNESCO World Heritage Site in Northern Tanzania (3°14'29.56"S and 35°29'16"E; Fig 1) with a total size of 8,256 km² [36]. This area extends over part of the Great Rift Valley of East Africa and contains grassland plains, savanna woodlands, forests, mountains, volcanic craters, lakes, rivers, and swampland. Ecologically, NCA is categorized into three zones; lowlands, midlands, and highlands [37], and its climatic zones span from semi-arid to montane forest climate, with average annual precipitation ranging from less than 500 mm up to 1,700 mm [38]. A description of the sources of the dataset used to produce a map in Fig 1 is provided in Table 1.

Rainfall in NCA is highly seasonal and spatially variable. The eastern slopes of the crater highlands receive average annual rainfall of about 1,200 mm/year, whereas the midlands receive about 800 mm/year and the lowlands receive only 400 mm/year [39]. Average annual temperatures lie between 2°C and 35°C [38]. We selected four wards: Nainokanoka (average elevation is 2,440 masl), Ngorongoro (1,996 masl), Endulen (1,637 masl) and Olbalbal (1,548 masl) for our field study to cover a large elevational gradient and varying distances to the fully



Fig 1. Map of Ngorongoro Conservation Area (NCA) in northern Tanzania showing locations of wild and domestic (cattle, sheep and goats) herbivore groups. In addition, we show the location of settlements that had been observed along road transects in the four wards Nainokanoka, Ngorongoro, Endulen and Olbalbal, from May 2018 to February 2019. (Source of dataset is provided in Table 1).

https://doi.org/10.1371/journal.pone.0267082.g001

Data sets	Institution	Citation and Website	
Protected Areas boundaries	Serengeti GIS and Data Center	[41] https://serengetidata.weebly.com/boundaries.html Accessed 2021-10-17	
Administrative wards	Tanzania National Bureau of Statistics	https://www.nbs.go.tz/index.php/en/census-surveys/population-and-housing-census/173- 2012-phc-shapefiles-level-three Accessed 2022-03-17	
Streams	Serengeti GIS and Data Center	[42] https://serengetidata.weebly.com/rivers-and-lakes.html Accessed 2021-10-18	
Road	Socioeconomic Data and Applications Center (SEDAC)	[43] https://sedac.ciesin.columbia.edu/data/set/groads-global-roads-open-access-v1 Accessed 2021-10-18	
Elevation raster file	USGS	[44] https://doi.org/10.5067/MEaSUREs/SRTM/SRTMGL1.003 Accessed 2022-03-18	
Settlements	Tanzania Wildlife Research Institute (TAWIRI)	https://tawiri.or.tz/ Accessed 2021-10-18	

Table 1.	Source of the datasets	used to produce th	e map of the Ngo	orongoro Conserva	tion Area, northern	Tanzania
----------	------------------------	--------------------	------------------	-------------------	---------------------	----------

https://doi.org/10.1371/journal.pone.0267082.t001

protected area, i.e., the NCA crater (Fig 1). The area covered by each elevation belt was 1,567 km² for lowlands (less than 1,600 masl): 1,819 km² for midlands (1,600–1,900 masl): and 1,789 km² for highlands (above 1,900 masl). The NCA crater fully excludes pastoralists and their livestock herds, whereas other parts of the NCA are shared by both pastoralists and wildlife. The Maasai are the dominant pastoralists in NCA, they live in settlements called boma or a homestead, which is a grouping of houses of multiple families enclosed by thorny *Acacia* branches to deter predators from entry [40].

The main economic activities in the NCA are livestock keeping and tourism [45]. The livestock species are cattle (*Bos taurus*), goats (*Capra aegagrus hircus*), sheep (*Ovis aries*) and donkeys (*Equus asinus*). The dominant wild mammalian herbivore species include plains zebra (*Equus quagga burchelli*), common eland (*Tragelaphus oryx*), blue wildebeest (*Connochaetes taurinus*), African buffalo (*Syncerus caffer*), Grant's gazelle (*Nanger granti*), and Thomson's gazelle (*Eudorcas thomsonii*). Less common species in the NCA are the giraffe (*Giraffa camelopardalis*), black rhinoceros (*Diceros bicornis*) and African elephant (*Loxodonta africana*) [46].

Data collection

Group sizes of wild herbivores were recorded during four sampling periods, two in each season (wet season: November–May, dry season: June–October) between May 2018 and February 2019. Four roads were chosen as transects. Each of those transects was visited four times during the entire study period. Transect 1 covered 55.8 km length and was distributed across an average (\pm SD) elevation of 2,097 \pm 288 masl; transect 2 covered 68.1 km and 1,659 \pm 281 masl; transect 3 covered 35.9 km, and 1,337 \pm 58 masl; and transect 4 covered 56.5 km and 2,448 \pm 90 masl. Roads were repetitively sampled using the road strip census method, in which animals were counted from the car within a certain strip width [47]. We drove the car at a constant speed of 25 km/h for 6 h each day, 3 h in the morning (07:00–10:00 h) and 3 h in the evening (15:30–18:30 h).

Observations of wild mammalian herbivore groups were restricted to distances within 250 m from the road to ensure visibility. For each sighting, we recorded the GPS coordinates and counted the number of individuals in the group of wild herbivores and livestock (defined as individuals within 50 m of each other). We used a rangefinder (Bushnell Elite 1500) to measure the perpendicular distance between the centre of each animal group and the observer [48]. The distances of all observed wild mammalian herbivore groups to the NCA crater rim (i.e., fully protected area), to the nearest settlement (i.e., human influence) and to the nearest stream (environmental factor) were obtained using QGIS version 3.6.

As the sample sizes for group observations of some species were rather low (S1 Fig) we categorized the wild herbivores into browsers (giraffe), grazers (zebra, wildebeest, and buffalo), or mixed feeders (Grant's gazelle, Thompson's gazelle and eland) [49]. We further omitted two outlying observations of an unexpectedly large Thompson's gazelle group and one elephant group. Livestock groups were categorized as either cattle or the combination of sheep and goats, due to the difficulty in distinguishing between sheep and goats in large mixed herds.

Data processing

We recorded wild herbivore group sizes across various village areas of the Ngorongoro Conservation Area (NCA) during the wet and dry season. We combined field-based techniques documenting locations and group sizes of wild herbivores and livestock with settlements locations collected by the Tanzania Wildlife Research Institute (TAWIRI) during an aerial census in the year 2016 [50]. Furthermore, we analyzed if wild herbivore group sizes were affected by (1) season, given the local climatic projections of greater rainfall variability, both within and between seasons [51, 52]; (2) landscape features such as distance to the fully protected area, i.e., the NCA crater, and distance to streams; (3) distance to human settlements; and (4) the number of livestock individuals present in proximity to the wildlife groups. A description of the range of variables used to model the group sizes of wild herbivores are listed in Table 2.

Data analyses

A generalized linear mixed model (GLMM) was applied to analyse the potential effects of season (wet vs dry), distance to the NCA crater, distance to streams (seasonal rivers in the ecosystem), elevation, livestock herds and distance to settlements on group sizes of wild herbivores. To account for repeated samples from the same transects, we nested transects in seasons and included them as a random factor. We further included the sampling date as a random factor [53]. All pair-wise correlation coefficients metric ranged from -0.4 to 0.4, indicating low levels of co-linearity [54]. We applied a zero-truncated negative binomial regression model because the observed group sizes were always larger than zero and the empirical histogram indicated that the data was strongly over-dispersed [55]. The positive negative binomial distribution was given by

$$f(y_i;k,\mu_i|y_i>0) = \frac{\frac{r(y_i+k)}{r(k) \times r(y_i+1)} \times \left(\frac{k}{\mu_i+k}\right)^k \times \left(1-\frac{k}{\mu_i+k}\right)^{y_i}}{\left(1-\left(\frac{k}{\mu_i+k}\right)^k\right)}$$

Table 2. Range of variables used to model the group sizes of wild herbivores in response to environmental variables, human settlements and livestock in the Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019.

Variable Name	Category	Data range (min-max)
Distance to streams (km)	environment	0.0-7.5
Distance to the NCA crater (km)	environment	0.3-31.7
Elevation (masl)	environment	1,288-2,654
Dry season versus wet season	environment	
Distance to the nearest settlement (km)	human	0.1-5.6
Number of cattle in proximity to wild herbivores	livestock	1-250
Number of sheep and goats in proximity to wild herbivores	livestock	1-842

https://doi.org/10.1371/journal.pone.0267082.t002

where y_i are the $i = 1, 2, ..., n_i$ observed wild herbivore group sizes, Γ is the gamma function, μ_i is the mean of the ordinary binomial distribution and k is the dispersion parameter [55, 56].

Our initial model (Eq. 1 in S2 Text) was based on the theory that variation in the environment, human activities, and competition with livestock affect the availability of resources that enable wild herbivores to form groups [57]. We incorporated interactions between season and human, environmental and livestock variables in the initial model because we expected that these covariate effects may vary seasonally and accounted for group size differences between feeding guilds (S1 Table). The perpendicular distances of animal groups to the observer were also included as an explanatory variable because herbivore group sizes could have been affected by the presence of a vehicle and closeness to roads (Eq. 1 in S2 Text).

Backwards selection of variables [58], using the drop1 function in R version 3.6.1 [59], as further used to select the most influential variables using Likelihood Ratio (LR) tests [60]. Variables were deleted from the full model starting with interaction and main effects of the variables with the highest *P*-values until all remaining variables had *P*-values of 0.056 or below. Although "statistical significance" is often interpreted as P < 0.05, we kept seasonal interaction terms with stream distances (P = 0.056) in the final model as water availability is an important variable in conservation planning [61]. Throughout the process, we kept distance to the observer as a confounding variable in the model.

During the selection procedure, seasonal interaction effects for sheep and goats were eliminated followed by the main effect of sheep and goats, seasonal interaction effects for cattle and the main effect of cattle (Eqs. 2–5 in <u>S1 Text</u> and <u>S2–S5</u> Tables). The last variables to be eliminated were seasonal elevation effects and the main effect of elevation (Eqs. 6 and 7 in <u>S2 Text</u> and <u>S6</u> and <u>S7</u> Tables).

We predicted wild herbivore group sizes based on the reduced model (Eq. 7 in S2 Text) in relation to the environmental, human and livestock variables for each feeding guild and season. Post hoc Tukey HSD Pairwise comparisons were applied for group size differences between feeding guilds. The zero-truncated negative binomial regression models were implemented via the glmmTMB R-package.

Results

We observed 176 groups of wild mammalian herbivores, with more observations (98; 56%) during the wet season than during the dry season (78; 44%). Wild herbivore group sizes were often larger in the wet season than in the dry season. Of all observed groups, 74% were formed by grazers, 18% by mixed feeders, and 8% by browsers. Grazers had similar group sizes as mixed feeders (t = -0.02, df = 161, P = 0.999) whereas browsers had group sizes that were by about 1/3 smaller than both grazers (t = -4.02, df = 161, P < 0.001) and mixed feeders (t = -3.42, df = 161, P = 0.002) (S2 Fig).

Larger groups closer to NCA crater in the dry season

Wild mammalian herbivore group sizes varied seasonally with distance away from the NCA crater (LR-test_{1, 2} = 10.5, P = 0.001). During the wet season, group sizes doubled from about 4 browsers at the NCA crater rim to 8 browsers at a distance of 32 km away from the crater (Fig 2A). Similarly, an average group size of 14 grazers and mixed feeders at the crater rim doubled to a group size of 23 grazers and mixed feeders at 32 km away from the crater rim (Fig 2A). In contrast, during the dry season, the estimated group sizes decreased about three-fold with increasing distance away from the NCA crater, i.e., from 8 browsers, 21 grazers and mixed feeders at the crater rim to 2 browsers, 6 grazers and 6 mixed feeders at 23 km distance away from the NCA crater (Fig 2B).



Fig 2. Trends in wild herbivore group sizes for browsers (giraffe), grazers (zebra, wildebeest, and buffalo) and mixed feeders (Grant's gazelle, Thompson's gazelle and eland) in relation to the fully protected area of the Ngorongoro crater (A, B), distance from streams (C, D), and settlements (E, F) during the wet season (left panels) and dry season (right panels) in the Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019.

https://doi.org/10.1371/journal.pone.0267082.g002

Larger groups further away from streams in the dry season

Wild herbivore group sizes varied across season with distance to streams (LR-test $_{1, 2} = 3.7$, P = 0.056). Most groups (67%) were observed at less than 2 km distance to streams, but group sizes did not change in relation to distance from the streams during the wet season, generally occurring from 0 to 8 km distance away from streams (Fig 2C). During the dry season, more than half of all herbivore groups (56%) were observed at less than 2 km distance to streams. But estimated group sizes increased by about three-fold with increasing distance from the streams, i.e., from 2 browsers, 7 grazers and 7 mixed feeders next to streams to 8 browsers, 24 grazers and 25 mixed feeders at 8 km away from streams (Fig 2D).

Larger groups close settlements

Wild herbivore groups responded differently to the presence of settlements in each season (LR-test _{1, 2} = 8.5, P = 0.004). During the wet season, wild herbivore group sizes were slightly higher closer to settlements than further, i.e., about 6 km, away (Fig 2E). However, during the dry season, group sizes decreased from 8 to 1 browser, 21 to 4 grazers and 22 to 6 mixed feeders with increasing distance away up to about 5 km away from settlements (Fig 2F).

Discussion

Our results on fluctuating group sizes of large wild herbivores in the Ngorongoro Conservation Area indicate that pastoralist activities and environmental variables affect the social organization of ungulates [10, 62]. These activities imply that animals have to spend more time being vigilant and have less time available for feeding [63–65]. Higher vigilance has potential consequences for reproductive fitness [66, 67], will reduce species abundance and alter community structure [68]. The tendency of increased wild herbivore group sizes we observed during the wet season agrees with results from another study [69, 70] that attributed the group size change to increased food availability.

During the wet season, wild herbivore group sizes increased further away from the NCA crater, possibly because herbivores disperse into short grass plains maintained by livestock grazing [40]. There is potential benefit from facilitation by livestock for wild herbivores, i.e., providing short-grass patches of high forage quality for growth and reproduction of herbivore populations [46, 71]. Our findings that wild herbivore group sizes increased closer to the NCA crater during the dry season is likely due to limited food availability further away from the NCA crater, triggered by competition with livestock further away from the fully protected area [72]. The NCA crater rim contains various shrubs and flowering plants [40] that may attract herbivores during the dry season [73, 74]. These spatial changes in group sizes might indicate a significant collective movement pattern towards the NCA crater potentially in search for food resources [75–77].

Moreover, the permanently flowing rivers inside the crater may attract animals in times of low rainfall, and particularly wildebeest, African buffalo, African elephant and zebra [78–81]. However, against our expectations we did not detect a humped-shaped distribution of wild herbivores with distance to water, contrary to other studies [15]. Our observed aggregation of herbivore groups close to streams in the dry season may indicate that herbivores were attracted to water. But increased competition and changes in vegetation resulting from trampling by herbivores and livestock might have influenced the herbivores' ability to form large groups [82–84].

Several studies suggest that associations between wild and domestic herbivores with similar body size and niche would lead to competitive exclusion of the wild herbivores [85, 86]. Contrary to our expectations, changes in wild herbivore group sizes were better explained by variations in the environmental and human settlements rather than by the presence of livestock. Whenever we saw livestock and wildlife together, there was no direct aggression or replacement by livestock (personal observations). On the other hand, resource use by Maasai cattle in NCA closely resembles that of resident wildlife [12] and diets of cattle and wild herbivores including impala, plain zebra, and wildebeest overlap in East African savannas [87]. However, evidence for competition between livestock and wild herbivores are scarce [9] except for few studies, which observed less abundance of wild herbivores within a radius of 10 km away from human settlements [4].

The increase in wild herbivore group sizes we observed closer to settlements implies that wild herbivores in NCA may have no choice but to aggregate close to settlements during the dry season to use water and food available at these sites. At NCA, as in other rangeland areas of eastern Africa, pastoral movement is largely governed by efforts to find the best pastures, though mostly constrained by the limited distribution of water in the dry season [88–90]. Emerging evidence also suggests a high concentration of wild herbivores in proximity to settlements to avoid predators [15].

Conclusion

Against our expectations, we found that livestock did not significantly influence wild herbivore group sizes in the multiple use landscape of the Ngorongoro Conservation Area but rather,

combined biotic and abiotic factors. Our results reveal how season, distance to protected areas, and distance to streams interactively shape herbivore group sizes in a multiple land use area. However, the observed decline in group sizes further away from the NCA crater during the dry season may suggest that wild herbivores had fewer resources (food and water) available as a result of land use by humans and their livestock. Our study is one of the few in this iconic Man and Biosphere reserve that quantified how human-driven factors might impact seasonal group sizes of wild herbivores. Our work only represents a snapshot over two seasons and we, therefore, recommend long-term monitoring of wild herbivore group sizes. This can provide a valuable indicator of temporal dynamics in wild herbivore group sizes given the ever increasing and poorly managed human and livestock population in the Ngorongoro Conservation Area. These research results can potentially guide alternative approaches to rangeland conservation practices.

Supporting information

S1 Fig. The total number of groups observed per wild and domestic mammalian herbivore species during the wet and during the dry season in the Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019. (TIF)

S2 Fig. Mean \pm SE of group sizes of wild herbivores browsers (giraffe), grazers (zebra, wildebeest, and buffalo) and mixed feeders (Grant's gazelle, Thompson's gazelle and eland) observed along road transects in the Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019. Boxes with the same letters do not differ significantly based on Tukey's HSD test at P = 0.05. (TIF)

S1 Data. Wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019. (XLS)

S1 Text. R code used to analyse the spatial and temporal distribution of wild mammalian herbivores in Ngorongoro Conservation Area from May 2018 to February 2019. (TXT)

S2 Text. Variable selection steps for analyzing wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables, and livestock. (DOCX)

S1 Table. Results of Generalized Linear Mixed Model (GLMM) of the initial model (S1 Text, Eq. 1) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables, and livestock. (Main effects of season, distances to streams, crater and settlements, sheep and goats, cattle and elevation are not shown). (XLSX)

S2 Table. Results of Generalized Linear Mixed Model (GLMM) after eliminating the seasonal interaction effect for sheep and goats (S1 Text, Eq. 2) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables, and livestock. (Main effects of season, distances to streams, crater and settlements, cattle and elevation, are not shown).

(XLSX)

S3 Table. Results of Generalized Linear Mixed Model (GLMM) after eliminating the main effect for sheep and goats (S1 Text, Eq. 3) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables, and livestock. (Main effects of season, distances to streams, crater and settlements, cattle and elevation are not shown). (XLSX)

S4 Table. Results of Generalized Linear Mixed Model (GLMM) after the eliminating the seasonal interaction effects for cattle (S1 Text, Eq. 4) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables, and livestock. (Main effects of season, distances to streams, crater and settlements and elevation are not shown).



S5 Table. Results of Generalized Linear Mixed Model (GLMM) after the eliminating the main effect for cattle (S1 Text, Eq. 5) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables. (Main effects of season, distances to streams, crater and settlements and elevation are not shown). (XLSX)

S6 Table. Results of Generalized Linear Mixed Model (GLMM) after the eliminating the seasonal interaction effects for elevation (S1 Text, Eq. 6) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from May 2018 to February 2019 in relation to feeding guilds, environmental and human variables. (Main effects of season, distances to streams, crater and settlements are not shown). (XLSX)

S7 Table. Results of Generalized Linear Mixed Model (GLMM) after the eliminating the main effect for elevation (S1 Text, Eq. 7) of wild herbivore group sizes recorded in Ngorongoro Conservation Area, northern Tanzania, from March 2018 to February 2019 in relation to feeding guilds, environmental and human variables. (Main effects of season, distances to streams, crater, and settlements are not shown). (XLSX)

Acknowledgments

We thank the Ngorongoro Conservation Area Authority (NCAA) management for permission to access the area. We thank our field assistants Juma Mkwizu (driver) and Jacob Yudah for contributing to the group size surveys and acknowledge three anonymous reviewers for constructive and helpful comments.

Author Contributions

Conceptualization: Cecilia M. Leweri, Gundula S. Bartzke, Maurus J. Msuha, Anna C. Treydte.

Data curation: Cecilia M. Leweri, Gundula S. Bartzke, Anna C. Treydte.

Formal analysis: Cecilia M. Leweri, Gundula S. Bartzke.

Funding acquisition: Cecilia M. Leweri.

Methodology: Cecilia M. Leweri, Gundula S. Bartzke, Anna C. Treydte.

Supervision: Maurus J. Msuha, Anna C. Treydte.

Writing – original draft: Cecilia M. Leweri.

Writing – review & editing: Cecilia M. Leweri, Gundula S. Bartzke, Maurus J. Msuha, Anna C. Treydte.

References

- 1. Barrette C. The size of Axis deer fluid groups in Wilpattu National Park, Sri Lanka. *Mammalia* 1991; 55: 207–220.
- Markham AC, Gesquiere LR, Alberts SC, et al. Optimal group size in a highly social mammal. Proc Natl Acad Sci U S A 2015; 112: 14882–14887. https://doi.org/10.1073/pnas.1517794112 PMID: 26504236
- Shen S-F, Akçay E, Rubenstein DR, et al. Group size and social conflict in complex societies. Am Nat 2014; 183: 301–310. https://doi.org/10.1086/674378 PMID: 24464203
- 4. Bergström R, Skarpe C. The abundance of large wild herbivores in a semi-arid savanna in relation to seasons, pans and livestock. *Afr J Ecol* 1999; 37: 12–26.
- Boone RB, Thirgood SJ, Hopcraft JGC. Serengeti wildebeest migratory patterns modeled from rainfall and new vegetation growth. *Ecology* 2006; 87: 1987–1994. https://doi.org/10.1890/0012-9658(2006) 87[1987:swmpmf]2.0.co;2 PMID: 16937638
- Boult VL, Sibly RM, Quaife T, et al. Modelling large herbivore movement decisions: beyond food availability as a predictor of ranging patterns. *Afr J Ecol* 2019; 57: 10–19.
- Pachzelt A, Rammig A, Higgins S, et al. Coupling a physiological grazer population model with a generalized model for vegetation dynamics. *Ecol Modell* 2013; 263: 92–102.
- Codling EA, Dumbrell AJ. Mathematical and theoretical ecology: linking models with ecological processes. *Interface Focus* 2012; 2: 144–149.
- **9.** Prins HH. Competition between wildlife and livestock in Africa. In: Prins HHT, Grootenhuis JG, Dolan TT (eds) *Wildlife conservation by sustainable use*. 2000, pp. 51–80.
- Baltazary A, Roskaft E, Treydte AC. Vigilance behaviour of wild herbivores when foraging with or without livestock. *Environ Nat Resour Res* 2019; 9: 64–76.
- 11. Ogutu, Owen-Smith N, Piepho HP, et al. Continuing wildlife population declines and range contraction in the Mara region of Kenya during 1977–2009. *J Zool* 2011; 285: 99–109.
- 12. Du Toit JT, Kock R, Deutsch J. *Wild rangelands: conserving wildlife while maintaining livestock in semiarid ecosystems.* Wiley-Blackwell, 2010.
- Sitters J, Heitkönig IMA, Holmgren M, et al. Herded cattle and wild grazers partition water but share forage resources during dry years in East African savannas. *Biol Conserv* 2009; 142: 738–750.
- 14. Georgiadis N, Olwero N, Ojwang' G, et al. Savanna herbivore dynamics in a livestock-dominated landscape: Dependence on land use, rainfall, density, and time. *Biol Conserv* 2007; 137: 461–472.
- Ogutu JO, Piepho HP, Reid RS, et al. Large herbivore responses to water and settlements in savannas. *Ecol Monogr* 2010; 80: 241–266.
- Butt B, Turner MD. Clarifying competition: the case of wildlife and pastoral livestock in East Africa. Pastoralism 2012; 2: 1–15.
- 17. Reid RS. Savannas of our birth: People, wildlife, and change in East Africa. University of California Press, 2012.
- Riginos C, Porensky LM, Veblen KE, et al. Lessons on the relationship between livestock husbandry and biodiversity from the Kenya Long-term Exclosure Experiment (KLEE). *Pastor Res Policy Pract* 2012; 2: 1–22.
- Rudolph K, Fichtel C, Schneider D, et al. One size fits all? Relationships among group size, health, and ecology indicate a lack of an optimal group size in a wild lemur population. *Behav Ecol Sociobiol* 2019; 73: 1–14.
- **20.** Marchand P, Garel M, Bourgoin G, et al. Combining familiarity and landscape features helps break down the barriers between movements and home ranges in a non-territorial large herbivore. *J Anim Ecol* 2017; 86: 371–383. https://doi.org/10.1111/1365-2656.12616 PMID: 27981576
- Pays O, Fortin D, Gassani J, et al. Group dynamics and landscape features constrain the exploration of herds in fusion-fission societies: The case of European roe deer. *PLoS One* 2012; 7: 1–8. <u>https://doi.org/10.1371/journal.pone.0034678</u> PMID: 22479652

- 22. Benedek AM, Sîrbu I. Dynamics of small-mammal communities along an elevational gradient. *Can J Zool* 2019; 97: 312–318.
- Strandburg-Peshkin A, Farine DR, Crofoot MC, et al. Habitat and social factors shape individual decisions and emergent group structure during baboon collective movement. *Elife* 2017; 6: 1–47. <u>https://doi.org/10.7554/eLife.19505</u> PMID: 28139196
- Adler PB, Hall SA. The development of forage production and utilization gradients around livestock watering points. In: Landscape Ecology. Springer, pp. 319–333.
- Peters MK, Hemp A, Appelhans T, et al. Predictors of elevational biodiversity gradients change from single taxa to the multi-taxa community level. *Nat Commun* 2016; 7: 13736–13736. <u>https://doi.org/10. 1038/ncomms13736 PMID: 28004657</u>
- Lee MA, Burger G, Green ER, et al. Relationships between resource availability and elevation vary between metrics creating gradients of nutritional complexity. *Oecologia* 2021; 195: 213–223. https://doi.org/10.1007/s00442-020-04824-4 PMID: 33458802
- Miller JR, Hobbs RJ. Habitat restoration—Do we know what we're doing? *Restoration Ecology* 2007; 15: 382–390.
- Stamps JA, Swaisgood RR. Someplace like home: Experience, habitat selection and conservation biology. Appl Anim Behav Sci 2007; 102: 392–409.
- Schweiger AH, Boulangeat I, Conradi T, et al. The importance of ecological memory for trophic rewilding as an ecosystem restoration approach. *Biol Rev* 2019; 94: 1–15.
- Estes RD, Atwood JL, Estes AB. Downward trends in Ngorongoro crater ungulate populations 1986– 2005: Conservation concerns and the need for ecological research. *Biol Conserv* 2006; 131: 106–120.
- Boone RB, Galvin KA, Thornton PK, et al. Cultivation and conservation in Ngorongoro Conservation Area, Tanzania. *Hum Ecol* 2006; 34: 809–828.
- Moehlman PD, Ogutu JO, Piepho H-P, et al. Long-term historical and projected herbivore population dynamics in Ngorongoro crater, Tanzania. *PLoS One* 2020; 15: e0212530. <u>https://doi.org/10.1371/journal.pone.0212530</u> PMID: 32155150
- IUCN IU for C of N. The biosphere reserve and its relationship to other protected areas | IUCN. 1979; 19.
- Homewood K, Rodgers WA. Maasailand ecology: pastoralist development and wildlife conservation in Ngorongoro, Tanzania. Cambridge University Press, 2004.
- de Boer WF, Vis MJP, de Knegt HJ, et al. Spatial distribution of lion kills determined by the water dependency of prey species. J Mammal 2010; 91: 1280–1286.
- UNESCO. Ngorongoro Conservation Area—UNESCO World Heritage Centre, <u>https://whc.unesco.org/en/list/39/</u> (1979, accessed 11 September 2019).
- Galvin KA, Thornton PK, Boone RB, et al. Ngorongoro Conservation Area, Tanzania: Fragmentation of a unique region of the greater Serengeti ecosystem. In: *Fragmentation in Semi-Arid and Arid Land-scapes*. Dordrecht: Springer Netherlands, 2008, pp. 255–279.
- Niboye EP. Vegetation cover changes in Ngorongoro Conservation Area from 1975 to 2000: The importance of remote sensing images. *Open Geogr J* 2010; 3: 15–27.
- Boone RB, Lackett JM, Galvin KA, et al. Links and broken chains: evidence of human-caused changes in land cover in remotely sensed images. *Environ Sci Policy* 2007; 10: 135–149.
- 40. Swanson LA. Ngorongoro Conservation Area: Spring of Life. University of Pennsylvania, 2007.
- Maliti H. Serengeti ecosystem boundaries including park, and protected areas digitized from 1:50,000 map sheets. SERENGETI GIS & DATA, https://serengetidata.weebly.com/boundaries.html (1996).
- 42. Maliti H, Hagen C von. Rivers and small drainages from Greater Serengeti-Mara Ecosystem digitized from 1:50,000 map sheets. *SERENGETI GIS & DATA*, https://serengetidata.weebly.com/rivers-and-lakes.html (2008, accessed 4 January 2022).
- 43. Center for International Earth Science Information Network (CIESIN)/Columbia University and ITOS (ITOS)/University of G. Global Roads Open Access Data Set, Version 1 (gROADSv1). Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC), https://sedac.ciesin.columbia.edu/data/set/groads-global-roads-open-access-v1/data-download (2013).
- 44. NASA J. NASA Shuttle Radar Topography Mission Global 1 arc second [Data set]. NASA EOSDIS Land Processes DAAC.
- Melita A, Mendlinger S. The impact of tourism revenue on the local communities' livelihood: A case study of Ngorongoro Conservation Area, Tanzania. J Serv Sci Manag 2013; 6: 117–126.
- Odadi WO, Jain M, Wieren VSE, et al. Facilitation between bovids and equids on an African savanna. Evol Ecol Res 2011; 13: 237–252.

- **47.** Hirst SM. Road-strip census techniques for wild ungulates in African woodland. *J Wildl Manage* 1969; 33: 40.
- **48.** Buckland S, Anderson D, Burnham K, et al. *Advanced distance sampling: estimating abundance of biological populations*. Oxford University Press, 2004.
- 49. Estes R, Otte D. The behavior guide to African mammals: including hoofed mammals, carnivores, primates. University of California Press, 2012.
- TAWIRI. Wildlife, livestock and bomas census in the Serengeti ecosystem, dry season, 2016. TAWIRI aerial survey report. 2016; 35.
- 51. Christensen JH, Hewitson B, Busuioc A, et al. Regional climate projections. Climate change 2007: The physical science basis. In: Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt MT and HLM (ed) Climate Change 2007: The physical science basis. Contribution of working group I to the fourth assessment report of the Intergovernmental panel on climate change. United Kingdom and New York: Cambridge University Press, Cambridge, 2007.
- McSweeney C, New M, Lizcano G, et al. The UNDP climate change country profiles: Improving the accessibility of observed and projected climate information for studies of climate change in developing countries. Am Meteorol Soc 2010; 157–166.
- Schielzeth H, Nakagawa S. Nested by design: Model fitting and interpretation in a mixed model era. Methods Ecol Evol 2013; 4: 14–24.
- 54. Agresti A, Montgomery DC, Peck EA, et al. Introduction to linear regression analysis. Wiley, 2013, pp. 1086–1089.
- 55. Zuur AF, Ieno EN, Walker NJ, et al. Mathematics for the negative binomial truncated model. In: Krickeberg W, Samet K, Tsiatis JM, et al. (eds) *Mixed effects models and extensions in ecology with R*. New York, NY, USA: Springer Science+Business Media, New York, 2009, pp. 265–265.
- 56. Zuur AF, Ieno EN, Walker NJ, et al. The negative binomial distribution. In: Krickeberg W, Samet K, Tsiatis JM, et al. (eds) *Mixed effects models and extensions in ecology with R*. New York, NY, USA: Springer Science+Business Media, New York, 2009, pp. 199–200.
- 57. du Toit JT, Cross PC, Valeix M. Managing the livestock–wildlife interface on rangelands. Springer, Cham, 2017, pp. 395–425.
- Ratner B. Variable selection methods in regression: Ignorable problem, outing notable solution. J Targeting, Meas Anal Mark 2010; 18: 65–75.
- 59. R Core Team. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna., https://www.r-project.org/ (2018, accessed 23 July 2019).
- Zuur AF, Ieno EN, Walker NJ, et al. GLM and GAM for Count Data. In: Zuur AF, Ieno EN, Walker NJ, et al. (eds) *Mixed effects models and extensions in ecology with R*. New York: Springer Science+Business Media, New York, 2009, pp. 209–243.
- Wasserstein RL, Lazar NA. The ASA statement on p-values: Context, process, and purpose. Am Stat 2016; 70: 129–133.
- Green DS, Zipkin EF, Incorvaia DC, et al. Long-term ecological changes influence herbivore diversity and abundance inside a protected area in the Mara-Serengeti ecosystem. *Glob Ecol Conserv* 2019; 20: e00697.
- Wang MY, Ruckstuhl KE, Xu WX, et al. Human activity dampens the benefits of group size on vigilance in Khulan (equus hemionus) in Western China. *PLoS One*; 11. Epub ahead of print 2016. <u>https://doi.org/10.1371/journal.pone.0146725</u> PMID: 26756993
- Ciuti S, Northrup JM, Muhly TB, et al. Effects of humans on behaviour of wildlife exceed those of natural predators in a landscape of fear. *PLoS One* 2012; 7: e50611. <u>https://doi.org/10.1371/journal.pone.</u> 0050611 PMID: 23226330
- 65. Setsaas T, Hunninck L, Jackson CR, et al. The impacts of human disturbances on the behaviour and population structure of impala (Aepyceros melampus) in the Serengeti ecosystem, Tanzania. *Glob Ecol Conserv*, 16. Epub ahead of print 2018. https://doi.org/10.1016/j.gecco.2018.e00467
- Parker KL, Barboza PS, Gillingham MP. Nutrition integrates environmental responses of ungulates. Funct Ecol 2009; 23: 57–69.
- Ogutu JO, Piepho H-P, Dublin HT. Reproductive seasonality in African ungulates in relation to rainfall. Wildl Res 2014; 41: 323.
- Bhola N, Ogutu JO, Piepho HP, et al. Comparative changes in density and demography of large herbivores in the Masai Mara Reserve and its surrounding human-dominated pastoral ranches in Kenya. *Biodivers Conserv* 2012; 21: 1509–1530.
- Bigalke RC. Observations on the behaviour and feeding habits of the springbok, Antidorcas marsupialis. Zool Africana 1972; 7: 333–359.

- 70. Jarman PJ. The social organisation of a ntelope in relation to their ecology. Behaviour 1974; 48: 215–267.
- 71. Verweij R, Verrelst J, Loth PE, et al. Grazing lawns contribute to the subsistence of mesoherbivores on dystrophic savannas. *Oikos* 2006; 114: 108–116.
- Odadi WO, Karachi MM, Abdulrazak SA, et al. African wild ungulates compete with or facilitate cattle depending on season. *Science* 2011; 334: 594. <u>https://doi.org/10.1126/science.1208653</u> PMID: 22053029
- Macandza VA, Owen-Smith N, Cain JW. Habitat and resource partitioning between abundant and relatively rare grazing ungulates. J Zool 2012; 287: 175–185.
- 74. Megaze A, Balakrishnan M, Belay G. Current population estimate and distribution of the African buffalo in Chebera Churchura National Park, Ethiopia. *Afr J Ecol* 2018; 56: 12–19.
- 75. Berdahl AM, Kao AB, Flack A, et al. Collective animal navigation and migratory culture: From theoretical models to empirical evidence. *Philos Trans R Soc B Biol Sci*; 373. Epub ahead of print 19 May 2018. https://doi.org/10.1098/rstb.2017.0009 PMID: 29581394
- 76. Fryxell JM, Berdahl AM. Fitness trade-offs of group formation and movement by Thomson's gazelles in the Serengeti ecosystem. *Philos Trans R Soc B Biol Sci*, 373. Epub ahead of print 19 May 2018. <u>https:// doi.org/10.1098/rstb.2017.0013 PMID: 29581398</u>
- 77. Bailey DW, Gross JE, Laca E a, et al. Mechanisms that result in large herbivore grazing distribution patterns. *J Range Manag* 1996; 49: 386–400.
- Veldhuis MP, Kihwele ES, Cromsigt JPGMGM, et al. Large herbivore assemblages in a changing climate: incorporating water dependence and thermoregulation. *Ecol Lett* 2019; 22: 1536–1546. <u>https://doi.org/10.1111/ele.13350 PMID: 31332945</u>
- Chamaillé-Jammes S, Valeix M, Fritz H. Managing heterogeneity in elephant distribution: Interactions between elephant population density and surface-water availability. J Appl Ecol 2007; 44: 625–633.
- **80.** Cornélis D, Benhamou S, Janeau G, et al. Spatiotemporal dynamics of forage and water resources shape space use of West African savanna buffaloes. *J Mammal* 2011; 92: 1287–1297.
- 81. Okello MM, Kenana L, Maliti H, et al. Population status and trend of water dependent grazers (buffalo and waterbuck) in the Kenya-Tanzania borderland. *Nat Resour* 2015; 06: 91–114.
- Veldhuis MP, Ritchie ME, Ogutu JO, et al. Cross-boundary human impacts compromise the Serengeti-Mara ecosystem. *Science* 2019; 363: 1424–1428. <u>https://doi.org/10.1126/science.aav0564</u> PMID: 30923217
- Davidson Z, Valeix M, Van Kesteren F, et al. Seasonal diet and prey preference of the African lion in a waterhole-driven semi-arid savanna. *PLoS One* 2013; 8: e55182. <u>https://doi.org/10.1371/journal.pone.</u> 0055182 PMID: 23405121
- van der Waal C, Kool A, Meijer SS, et al. Large herbivores may alter vegetation structure of semi-arid savannas through soil nutrient mediation. *Oecologia 2011 1654* 2011; 165: 1095–1107. https://doi.org/ 10.1007/s00442-010-1899-3 PMID: 21225433
- **85.** Voeten MM, Prins HHT. Resource partitioning between sympatric wild and domestic herbivores in the Tarangire region of Tanzania. *Oecologia* 1999; 120: 287–294. https://doi.org/10.1007/s004420050860 PMID: 28308091
- Acebes P, Traba J, Malo JE. Co-occurrence and potential for competition between wild and domestic large herbivores in a South American desert. J Arid Environ 2012; 77: 39–44.
- 87. Foufopoulos J, Altizer S, Dobson A. Interactions between wildlife and domestic livestock in the tropics. In: Vandermeer J (ed) *Tropical Agroecosystems*. CRC Press, 2002, pp. 230–244.
- Dwyer MJ, Istomin K V. Theories of nomadic movement: A new theoretical approach for understanding the movement decisions of Nenets and Komi reindeer herders. *Hum Ecol* 2008; 36: 521–533.
- Western D, Dunne T. Environmental aspects of settlement site decisions among pastoral Maasai. Hum Ecol 1979; 7: 75–98.
- Boles OJC, Lane PJ. The green, green grass of home: an archaeo-ecological approach to pastoralist settlement in central Kenya. Azania 2016; 51: 507–530.