

2023-02-16

# Chitosan-coated liposomes of *Carrisa spinarum* extract: synthesis, analysis and anti-pneumococcal potency

Rubaka, Clarence

ICE Publishing

---

<https://doi.org/10.1680/jbibn.22.00046>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

## Chitosan-coated liposomes of *Carrisa spinarum* extract: synthesis, analysis and anti-pneumococcal potency

Clarence Rubaka, Jeremiah Waweru Gathirwa, Hamisi M Malebo, Hulda Swai, Nicole Remaliah Samantha Sibuyi, Askwar Hilonga, Admire Dube

To download the complete text, click that link.

DOI: <http://dx.doi.org/10.1680/jbibr.22.00046>

### Abstract

In the present study, a chitosan (CS)-coated liposome (LipCsP-Chitosan) nanocarrier was fabricated for the delivery of *Carissa spinarum* (CsP) polyphenols to improve bioavailability and anti-pneumococcal potential against *Klebsiella pneumoniae*. LipCsP-Chitosan was synthesized by the ion gelation method and characterized by using a Malvern zetasizer and Fourier Transform Infrared (FTIR); CsP encapsulation and release kinetics were investigated. Anti-pneumococcal activity of the nanoformulations was accessed by agar-well diffusion and microdilution assays. LipCsP-chitosan exhibited a hydrodynamic size and zeta potential of  $365.22 \pm 0.70$  nm and  $+39.30 \pm 0.61$  mV, respectively. CsP had an encapsulation efficiency of 81.5%. FTIR analysis revealed the interaction of the liposomes with chitosan and the CsP. A biphasic CsP release profile followed by a sustained release pattern was observed. LipCsP-Chitosan presented a higher bioaccessibility of polyphenols in the simulated gastric phase ( $74.1\% \pm 1.3$ ) than in the simulated intestinal phase ( $63.32\% \pm 1.00$ ). LipCsP-chitosan had a relative inhibition zone diameter of  $84.33\% \pm 2.51$  when compared to CsP. At minimum inhibition concentration of 31.25 mg/mL, LipCsP-Chitosan reduced the viability of *Klebsiella pneumoniae* by  $57.45\% \pm 3.76$  after 24 h. The results obtained from the current study offer a new approach to the utilization of LipCsP-Chitosan as nanocarriers for candidate anti-pneumococcal agents.