

2022-07-08

Contribution of Illicit Drug Use to Pharmaceutical Load in the Environment: A Focus on Sub-Saharan Africa

Ripanda, Asha

Hindawi

<https://doi.org/10.1155/2022/9056476>

Provided with love from The Nelson Mandela African Institution of Science and Technology

Review Article

Contribution of Illicit Drug Use to Pharmaceutical Load in the Environment: A Focus on Sub-Saharan Africa

Asha S Ripanda ^{1,2} Mwemezi J. Rwiza ² Elias Charles Nyanza ^{3,4}
Revocatus L. Machunda ² and Said Hamadi Vuai¹

¹Department of Chemistry, College of Natural and Mathematical Sciences, University of Dodoma, P.O. Box 338, Dodoma, Tanzania

²School of Materials, Energy, Water and Environmental Sciences, The Nelson Mandela African Institution of Science and Technology, P.O. Box 447, Tengeru, Arusha, Tanzania

³School of Public Health, Catholic University of Health and Allied Sciences, P.O. Box 1464, Mwanza, Tanzania

⁴Department of Community Health Sciences, University of Calgary, Calgary, Canada

Correspondence should be addressed to Asha S Ripanda; ripandaa@nm-aist.ac.tz

Received 31 July 2021; Revised 25 October 2021; Accepted 10 May 2022; Published 8 June 2022

Academic Editor: Carla Patrícia Silva

Copyright © 2022 Asha S Ripanda et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Illicit drug abuse and addiction are universal issues requiring international cooperation and interdisciplinary and multisectoral solutions. These addictive substances are utilized for recreational purposes worldwide, including in sub-Saharan Africa. On the other hand, conventional wastewater treatment facilities such as waste stabilization ponds lack the design to remove the most recent classes of pollutants such as illicit drug abuse. As a result, effluents from these treatment schemes contaminate the entire ecosystem. Public health officials are concerned about detecting these pollutants at alarming levels in some countries, with potential undesirable effects on aquatic species and increased health hazards through exposure to contaminated waters or recycling treated or untreated effluents in agriculture. Contaminants including illicit substances enter the environment by human excreta following illegal intake, spills, or through direct dumping, such as from clandestine laboratories, when their manufacturer does not follow accepted production processes. These substances, like other pharmaceuticals, have biological activity and range from pseudopersistent to highly persistent compounds; hence, they persist in the environment while causing harm to the ecosystem. The presence of powerful pharmacological agents such as cocaine, morphine, and amphetamine in water as complex combinations can impair aquatic organisms and human health. These compounds can harm human beings and ecosystem health apart from their low environmental levels. Therefore, this article examines the presence and levels of illicit substances in ecological compartments such as wastewater, surface and ground waters in sub-Saharan Africa, and their latent impact on the ecosystem. The information on the occurrences of illicit drugs and their metabolic products in the sub-Saharan Africa environment and their contribution to pharmaceutical load is missing. In this case, it is important to research further the presence, levels, distribution, and environmental risks of exposure to human beings and the entire ecosystem.

1. Introduction

The problem of drug abuse is acute worldwide, and it threatens people's lives and ecosystem sustainability. According to the United Nations Office on Drugs and Crime (UNODC) (2020) projections, 269 million persons abused substances globally in 2018, exaggerated by 30% from 2009, and over 35 million people suffer from substance use disorders [1, 2]. Although the full impact of COVID-19 on the

drug market is not yet determined, the pandemic's border and other limitations have already resulted in drug shortages, resulting in higher costs and lower purity, which magnify drug problems. People who are socially and economically disadvantaged are more likely to develop drug use disorders. While drug usage is more prevalent in industrialized countries than in developing countries and affluent parts of society are more affected, drug use disorders are more common in socially and economically disadvantaged

individuals [1, 2]. This may be contributed by eating habits, as the effects of the drug depend on multiple factors such as genetics, health, age, dose, and stomach contents. Only one in every eight people who require drug treatment receives it.

On the other hand, only one out of every five patients in recovery clinics is a woman, despite one out of every three drug users being a woman [1, 2]. Half of the 11 million injecting drug users are infected with hepatitis C, and 1.4 million are infected with HIV. Apart from worldwide drug control measures, amphetamine seizures increased between 2009 and 2018, though global precursor control improved [2, 3]. On the other hand, traffickers and producers turn to designer compounds to synthesize amphetamine, methamphetamine, and ecstasy to avoid international restrictions [1, 2, 4]. At the same time, with stringent measures and the need to get high without legal constraints, drug abusers are shifting from controlled substances of abuse to designer drugs [1]. Traditional misused substances such as heroin and cocaine continue to be produced at some of the greatest levels ever seen in the modern history. The rise in international drug supply and petition complicates law enforcement efforts, intensifies health hazards, and makes it more difficult to prevent and treat substance use disorders [2]. The need for drug traffickers to get a higher profit causes the observed trend of increased accessibility to novel psychoactive substances (NPS). Technical synthesis of these substances is easier, at a low cost and easier conversion to varieties by substitution, which are undetectable by current standard toxicological screens [2], which help to avoid the law prohibiting their sales and use [5–8].

Illicit drugs reach the environment mainly through human excreta. Unmetabolized parent substances and their metabolites are released in high quantities into residential wastewater via urine or faeces [5, 9]. Based on conducted studies, reports of occurrences of methamphetamine, amphetamine, morphine, MDMA, and metabolites of cocaine such as benzoylecgonine and ecgonine methyl ester are reported to be the most prevalent deposits in effluents [5, 10], indicating the possibility of a regular occurrence in surface waters than other chemicals [5]. Drugs such as morphine, amphetamines, and MDMA have significant biological actions, and their presence as complex combinations in the environment may be hazardous and poses dangers to human health and the aquatic ecosystem, despite relatively low environmental concentrations [9, 11–13]. Apart from the reported effects of these illicit substances, residues, and their transformation products on the environment [9, 11–13], there are no recommended limits for permissible quantities of these compounds in surface water, tap/drinking water, or wastewater. However, more work is needed to give a clearer picture and accurate risk assessment.

Apart from all of these facts, there are 17 agreed-upon sustainable development goals to be met by 2030, including 6 goals which are as follows: improving water quality by reducing contamination, eradicating, discarding, and minimizing the release of hazardous substances and materials, halving the proportion of crude effluents, and significantly increasing recycling and safe reuse globally [14]. To achieve this aim, there must be a global focus on reducing urban river pollution

through monitoring the quality of discharge from industries and waste stabilization ponds. However, no present rule requires these developing contaminants to be determined in treated wastewater, surface water, drinking water, or the atmosphere. Although drug abuse is now recognized as a global problem [15–17], the secondary effects of these substances have not yet been evaluated, such as their contribution to pharmaceutical load in the environment. The evidence of illicit drugs in waste and river waters is mostly available in developed countries [3, 18–22]. However, there is a scarcity of information, status, and knowledge regarding Oceania, Asia, and Africa, particularly in sub-Saharan Africa and a huge part of the world. The incidence patterns, destiny, and impact on many environmental compartments remain unknown. As a result, additional research is needed to determine their occurrence in waste and surface waters in developing nations to create a safer ecology.

2. Illicit Drug Use

Substance abuse, often known as drug abuse, includes using illicit drugs in quantities or with methods that endanger the user or others; it is a type of substance abuse disorder. Society is mostly concerned with the primary effects of drug abuse; this is possibly because the evidence of the effects of addiction and its contribution to the global burden of diseases is available [23–25]. The contribution of drug abuse to the pharmaceutical load and pollution, in general, should not be ignored as these substances may pose an unknown risk to the ecosystem, which may impair its sustainability.

3. Current Status of Abuse

Initially, traditional drugs of abuse such as heroin prevailed in the illicit drug scene. Later, other products such as novel psychoactive substances (NPS) and other drugs without restrictions were introduced. NPS have not managed to replace the traditional drugs of abuse completely; therefore, they coexist, complicating addiction treatments [26, 27]. The contribution of COVID-19 pandemic and economic downturn to the negative effects of illicit drugs on the impoverished, marginalized, and vulnerable communities is yet to be determined [1, 2]. However, border restrictions are said to increase the price and decrease purity, which may magnify drug problems. These substances are used singly or as cocktails for different reasons, such as to increase effects, for example, the mixture of prescription of opioids, cannabinoids, and cathinones [28–30]. Reports of the coexistence of these substances in the black market are available [31–35], and information on the possibility of being utilized by similar methods such as inhalation, smoking, and injecting is available [36–40]. Often, the consumers of substances lack awareness of the types of substances they are using and the interrelated risks; this is often because of mislabelling and adulteration of these substances [9, 39, 41–43]. Previous studies reported [31, 44–48] that drug misuse among teenagers and older persons in sub-Saharan Africa and the potential harm it causes are expected to rise until 2050 [49]; this indicates the need for intervention.

4. Drug Abuse in Sub-Saharan Africa

Drug abuse problem is increasing in sub-Saharan Africa, with reports of youth engagement in risk behaviour, including injecting and multiple drug use [49–56]. Olawole and colleagues (2018) conducted a systematic review which included 143201 adolescents from sub-Saharan Africa, having an average age of 16 years [55]. Apart from the fact that the incidence of use of any drug among teenagers was 41.6%, a higher rate of 55.5% was observed in Central Africa. It was further observed that depressant abuse was 11.3%, followed by amphetamines at 9.4%, and heroin was the least at about 4% [55]. These results indicate that youth in sub-Saharan Africa abuse multiple drugs, including prescription drugs (PDs), traditional drugs of abuse (TDA) such as cocaine, novel psychoactive substances (NPS), and natural products, and their mixtures as reported by other researchers [47, 57, 58]. The drawbacks of abuse are seen among youth as the mean age of abusers is about 16 years. These are expected to be participating in school activities. Youths are deteriorating, and as a result, the workforce is lost. Instead of youths participating in income-generating activities, they seek addiction treatment for recovery. Therefore, there is a need for a multidisciplinary approach to tackling drug abuse issues.

5. Effects of Illicit Drug Abuse

Reports of drug problems are accompanied by vindictive social impacts such as financial difficulties and communal life, the adverse effect on the industry, education, and family life, and its contribution to crime, violence, and homelessness [59–61]. The mode of drug use also contributes to risk factors, for example, injecting drug use becomes a route through which blood-borne virus circulates to the general population. The most-reported effect of illicit substance abuse is addiction, but the possibility of effects on human beings and the ecosystem through the food chain still needs to be investigated. Figure 1 shows the roots through which human health and the entire ecosystem may be exposed to these substances.

The results of a study by Aghababaei and colleagues revealed the presence of poisonous metals and bacterial contamination in abused drugs. The study pointed out the presence of the greatest quantities of lead (Pb) ranging 63–213 g/g and chromium (Cr) ranging 427–468 g/g in opium samples [62]. Furthermore, the maximum incidence of microbial pollution was observed in opium samples, whereas heroin samples had the lowest [62]. *Clostridium tetani* was the most common microbe in the tested samples, accounting for almost half of all contaminants [62]. These results indicate the possibility of contamination risks to drug abusers, resulting from these substances being produced illicitly and, therefore, a low possibility of following good manufacturing practices.

On the other hand, the reports of combined exposure of a compound itself or its metabolites and mixtures of multiple drugs are missing globally. The effects of pharmaceuticals in the environments have been reported, such as the

development of drug-resistant strain, the possibility of giving rise to more toxic metabolites, extinction of species, mammals, aquatic ecosystem, plants, soil arthropods, and toxic effects [63–65]. The reports of poisonous effects of substance use on the environment are required. The effects of a 96-hour exposure to 500 ng/L of cocaine and 20 ng/L of benzoylecgonine or their mixture on *Mytilus galloprovincialis* were examined [12]. The results show that the oxidative status in the gills and digestive glands of the Mediterranean mussels was not altered by exposure to a single component. The dissimilarity is the exposure to a mixture of illicit drugs which modulated the antioxidant activity in the gills but not in the digestive gland of treated mussels [12]. More research studies about the toxicity of illicit drug mixtures are needed to emphasize the risk that these compounds provide to the ecosystems [61]. Apart from direct health risks to users, other illicit drugs contribute to the pollution of our environment, since some are pseudo-persistent while others are persistent. Therefore, illicit drugs persist in the environment, while they bioaccumulate, bioconcentrate, and biomagnify through the food chain. These substances have been identified in environmental samples such as soil, tap/groundwater, wastewater, urban rivers, and streams in developed countries [9, 66–75] and some African countries such as South Africa and Tunisia in some matrices. However, information about their presence in environmental compartments of sub-Saharan countries, possible harm to the ecosystem, and contribution to the pharmaceutical load is not certain. The environmental threat created by emerging contaminants and the contribution of drug abuse should not be ignored as they are associated with more risk factors leading to pollution.

6. Evaluation of Environmental Illicit Drug Load

The quantity of illicit drugs used by the inhabitants served by the wastewater treatment scheme is estimated using wastewater-based epidemiology (WBE) [76]. In this case, raw wastewater is collected and investigated for the presence of quantifiable biomarkers of drugs, and the amount of licit or illicit drugs consumed by the inhabitants served by the wastewater treatment scheme is back-calculated [76] after taking composite samples of raw wastewater and analyzing them for certain components such as illicit drugs [76]. The total consumption of a drug is calculated using this number plus a correction factor that considers the average excretion rate of a certain drug residue and the molecular mass ratio of the parent drug to its metabolite [76]. The comparison among cities is done by taking the daily values and dividing them by the number of individuals served by the treatment scheme. This value can be articulated in daily amounts (or daily doses) per thousand inhabitants [76]. The improvement of integrity and scalability of this technique is increased by adopting a standardised procedure [77]. It is important to ensure that data from different sources are more reliable and comparable. The Sewage analysis CORE group Europe (SCORE) network conducted its first Europe-wide study in 2011, and its results provided a comprehensive

TABLE 2: Comparative active chemicals (ACs) loads reported from various categories of drug.

Drug category	Daily load in g/day	References
Anti-inflammatory	100–3000	[78]
Anticonvulsant	0–100	
Anticorrosive	100–200	
Opioids	0–500	
Illicit drugs	800–3800	
Antibiotics	500–4200	

The reported chemical load from illicit drugs is comparable to anti-inflammatory and antibiotics (Table 2). However, more data are required to initiate guideline reforms to include illicit drugs in environmental assessment and monitoring activities for environmental sustainability.

Illicit drugs have been shown to contaminate aquatic ecosystems worldwide, which has been expected for many years [89–92]. Horkey and colleagues revealed fish addiction results in changes in habitat preferences, with unanticipated negative repercussions on individual and inhabitant levels [93]. This may be contributed by the dumping of illegal drugs into freshwater ecosystems or contaminated effluents. Methamphetamine is regarded as one of the most serious worldwide health threats, producing addiction and behavioural changes in brown trout *Salmo trutta* at a concentration of $1 \mu\text{gL}^{-1}$ [93]. The impact of these substances may be amplified through bioaccumulation, bioconcentration, and biomagnification through the food chain, indicating the possibility of harming ecosystems. When consumed in a food chain, these contaminated fish may cause health problems, including addiction to nonabusers. These compounds may also lead to transformational and disinfection products as other contaminants that may further impact ecosystem sustainability. Figure 2 shows the structure of methadone and its disinfection by-product.

The environmental impact of remnants of active chemicals from contraceptive pills includes the feminization of male fish downstream, resulting in a deteriorating fish population [94–96]. Few reports on misuse of oral contraceptive pills (OCPs) for topical use such as to increase hair growth and give negative findings for addictive drug screening tests are available [97]. Increased topical use of OCPs may throw the ecosystem off balance and lead to an unsustainable environment. Pharmaceutical drug methadone, which is used to mitigate addiction caused by the recreational use of opioid heroin, was observed to create N-nitrosodimethylamine (NDMA) in surface and wastewater due to the application of chlorination as a disinfection technique [88]. A probable human carcinogen reaches drinking water after formation as disinfection by-products, and its threshold in drinking water is 7 ng/L according to USA guidelines [88]. The evaluation and monitoring of these contaminants are important for the safety of the ecosystem.

8. Challenges of Sewerage Systems in Sub-Saharan Africa

It is observed that over 50% of the world's population lives in urban areas [20], individuals living in slums are increasing by about 20 million each year, and people living

in metropolitan areas lack proper sanitation. The sewage system in sub-Saharan Africa faces difficulties, since it was not intended to handle emerging contaminants [66] such as pharmaceuticals and drug abuse. Therefore, these substances remain in the wastewater after treatment and are released into the environment. Another challenge is faecal sludge management (FSM). In most settings, the sludge is utilized as an organic fertilizer, as the facility is incapable of removing or not designing the sludge also is contaminated. Urban cities in sub-Saharan Africa lack the proper FSM approach, and most cities do not prioritize FSM. FSM is considered an insignificant aspect of environmental sanitation, and discharging FS in water bodies, lagoons, and the environment is regularly considered the normal and proper way of dealing with FS. At the same time, through the utilization of sludge in the reclamation of agricultural land or fertilizer contaminants, their metabolites and transformation products [98] are circulating in the environment [73, 99–102]. These substances are toxic; hence, the possibility of causing harm to humans or the environment is rising. Another issue confronting wastewater treatment plants in sub-Saharan Africa is the lack of modern, state-of-the-art facilities for contamination identification and quantification. This issue will significantly be observed in drug of abuse as traffickers and manufacturers change precursors and drug design to avoid legal constraints concerning their manufacture, sale, and use [47, 48].

9. Global Occurrences of Illicit Drugs in Environmental Compartments

Tendencies of encountering drug abuse and other drugs in the environment are increasing, and this threat is recognized as an issue of concern. Researchers employed wastewater analysis to track the pollution produced by these compounds in the environment and assess the efficacy of treatment plants in removing them [19, 103, 104]. Sources, behaviour, and fate of chemicals in the environment as singly have been investigated broadly [105, 106]. The current research indicates the existence of these drugs in mixtures such as drug abuse and other prescription drugs, which may result in toxicity due to drug-drug interactions or combined concentration for a drug with a similar mode of action. Reports of global occurrences of illicit drugs in wastewater effluents, influents, and urban rivers are available [18, 19, 40, 42, 53, 107–115]. Table 3 provides data on the prevalence of illicit drugs around the world. As millions of people abuse amphetamine-like stimulants, marijuana, cocaine, heroin, and other substances, the volumes of illicit drugs used worldwide may be equivalent to those of prescriptive drugs. Therefore, to control and monitor our environment, drug abuse should be kept under consideration as these substances are among emerging contaminants [17, 104, 116].

In most of the areas provided in Table 3, the load increased on Saturday and Sunday. This might be because the number of residents increased in participating in

TABLE 4: Identified illicit drugs in South African wastewater.

Substance	Illicit drug (mg/day/1000 inhabitants)		References
	WWT1	WWT2	
MDMA	0–2.23	0.5–65	[124]
Methamphetamine	15–45	140–250	
Mephedrone	-	7.6	
06-MAM	-	2.09–5.54	
Cocaethylene	0–16	5–80	
Cocaine	2–6	10–40	
Benzoylcegonine	6–17	38–78	
Amphetamine	0.7–1.4	4–12	
HMMA	0–0.25	0.7–2.5	

MDMA, 3,4-methylenedioxymethamphetamine; 06-MAM, 6-monoacetylmorphine; Bdl, below detection limit; WWT, wastewater treatment.

TABLE 5: Identified illicit drugs in northern Tunisia.

Substance	Illicit drugs (mg/day/1000 inhabitants)		References
Amphetamine	LOQ–79		[125]
MDMA	LOD–51		
Methamphetamine	LOD–24		
Benzoylcegonine	LOQ–450		
Mephedrone	LOQ–102		




FIGURE 3: Structure of heroin and its metabolites which may further be changed to transformational by-products due to environmental and treatment conditions.

recreational activities. As a result, more people consume drugs, possibly more doses because there is no work, and others may come from far areas for recreation.

10. Occurrences of Illicit Drugs in African Environmental Compartments

Reports of drug abuse in South Africa are available [117–122], and the increased crime rate has been associated with drug abuse apart from other factors [118, 119]. Identification of multiple drugs, including TDA and NPS, indicates polydrug nature and the possibility of contamination of the ecosystem. This study also shows that the possibility of the availability of clandestine laboratories in the area or reported MDMA and HMMA is from clandestine sources. This is because both are chiral with conventional manufacturing methods [123]. The list of identified illicit drugs in South African wastewater is provided in Table 4.

11. Tunisia

Identification of illicit drugs in wastewater confirms their use, informs their presence in the environment, and the possibility of harm to the ecosystem. Multiple drugs indicate the polydrug nature of abuse and the existence of TDA and NPS in the black market requires intervention. The most significant aspect is that these compounds are introduced into the environment, increasing the pharmaceutical load and decreasing ecosystem safety. Table 5 provides the list of identified illicit drugs in northern Tunisia.

12. Occurrences of Illicit Drugs in Sub-Saharan Africa Environmental Compartments

In Sub-Saharan Africa, the evidence on the existence of illicit drugs, their metabolites, and transformation products is not sufficient. For example, it has been revealed that there are few reports on the presence of illegal drugs, metabolites, and transformation by-products in the environment of several sub-Saharan African countries. Officials are concerned with potential harm to human beings and the environment due to their toxicity, by itself or in combination [60, 126, 127]. The scarcity of information is contributed by the fact that some of these countries lack state-of-the-art laboratory facilities for their identification and quantification [128]. Therefore, there is a need for monitoring authorities and policymakers to ensure the availability of these types of equipment for analysis of emerging contaminants, including drug abuse, for a safer ecosystem.

13. Circulation of Drugs of Abuse in the Ecosystem

In the 1970s, countries in Europe and the United States developed nearly full sewer networks and wastewater treatment plants (WWTPs), putting a stop to the practice of directly discharging sewage into metropolitan waterways [129]. This halted the decline of many urban rivers' water quality,

gradually improving [129, 130]. At the start of rapid urbanization in the 1980s, China experienced major pollution due to a big amount of sewage being immediately released into rivers; the same was true in most other developing countries [129]. So far, most African governments, mostly in sub-Saharan Africa, have not restricted sewage discharge straight into urban rivers and other environmental compartments [100–102]. In the sewage based on the design of sewage treatment, there is the possibility of further transformation of these substances, possibly into more toxic by-products. The majority of these illicit drug residues have powerful pharmacological activity, and their existence in aquatic settings might have serious consequences for human beings and the ecosystem. Figure 3 shows the structure of the illicit drug heroin and its metabolites which may further be converted into transformational by-products due to treatment conditions or environmental conditions.

Mostly identified drugs such as ecstasy, methamphetamine, MDA, morphine, and cocaine have powerful activity, and their presence in surface waters as complex mixes with residues of innumerable drugs may result in unforeseen pharmacological interactions, with hazardous consequences on the aquatic ecosystem, as reported by other researchers [13, 18, 42, 60, 61, 73, 86, 131, 132]. In the ecosystem, these substances may bioaccumulate, bioconcentrate, and biomagnify through the food chain, as shown in Figure 1. When humans consume food, these substances may enter into a human being and create possible harm, especially to nonabusers.

14. Conclusions

Apart from the basic repercussions of drug abuse, which include chronic diseases such as cirrhosis, heart disease, blood-borne bacterial and viral infections, and mental problems indicating contribution to the global burden of diseases [23–25], the contribution of drug addiction to the degradation of the environment and biodiversity loss must be considered. Illicit drugs such as opioids were observed to be contaminated with heavy metals and microorganisms [62]. Reports of environmental and ecosystem contamination by illicit drug use are available, including addiction in fish as a result of methamphetamine exposure [93], which indicates the possibility of development of addiction to nonabusers through the food chain. Therefore, the development of an unsustainable ecosystem needs intervention. These results indicate the possibility of drug abuse contributing to contamination and exposure to contaminants and possible harm. In this case, studies on environmental effects of drug abuse need to be conducted singly and in combination to evaluate the contribution of illicit drug use to environmental contamination for conservation and sustainability of the ecosystem.

Abbreviations

AMR:	Antimicrobial resistant
ACs:	Active chemicals
BFRs:	Brominated flame retardants
BE:	Benzylecgonine

COC:	Cocaine
ECs:	Emerging contaminants
COVID-19:	Coronavirus disease-19
DOA:	Drug of abuse
FS:	Faecal sludge
FSM:	Faecal sludge management
LOQ:	Limit of quantification
LOD:	Limit of detection
MDMA:	3,4-Methylenedioxymethamphetamine
06-MAM:	6-Monoacetylmorphine
NPS:	Novel psychoactive substances
NDMA:	N-Nitrosodimethylamine
PFCs:	Perfluorochemicals
PDs:	Prescription drugs
TDA:	Traditional drug of abuse
UNODC:	United Nation Office on Drug Control
WWTPs:	Wastewater treatment plants.

Data Availability

The data used to support this study are included within the article.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Authors' Contributions

A.R, M.J.R., and S.V idealized and conceptualized the study. A.R, M.J.R, R.M, and E.N wrote the article. M.J.R, E.N, S.V, and R.M critically evaluated the study. M.J.R and S.V prepared the final draft and managed references. M.J.R., E.N, S.V, and R.M. edited the article for language. All authors have read and agreed to the submitted version of the manuscript.

Acknowledgments

The authors thank the management of the Department of Chemistry, College of Natural and Mathematical Sciences, The University of Dodoma, Dodoma, Tanzania, and the entire staff of the School of Materials, Energy, Water, and Environmental Sciences, Mandela African Institution of Science and Technology, for their challenging contribution during graduate seminars that lead to improvement of this work.

References

- [1] UNODC, "United Nations Office on Drugs and Crime, COVID-19 and the Drug Supply Chain: From Production and Trafficking to Use, UNODC Research and Trend Analysis Branch, Vienna, Austria., 2020.
- [2] J. Lobo Vicente, H. Chassaigne, M. V. Holland et al., "Systematic analytical characterization of new psychoactive substances: a case study," *Forensic Science International*, vol. 265, pp. 107–115, 2016.
- [3] K. Y. Kim and J.-E. Oh, "Evaluation of pharmaceutical abuse and illicit drug use in South Korea by wastewater-based epidemiology," *Journal of Hazardous Materials*, vol. 396, Article ID 122622, 2020.
- [4] J. Mounteney, P. Griffiths, R. Sedefov, A. Noor, J. Vicente, and R. Simon, "The drug situation in Europe: an overview of data available on illicit drugs and new psychoactive substances from European monitoring in 2015," *Addiction*, vol. 111, no. 1, pp. 34–48, 2016.
- [5] J. B. Zawilska, "Legal highs"—an emerging epidemic of novel psychoactive substances," *International Review of Neurobiology*, vol. 120, pp. 273–300, 2015.
- [6] J. B. Zawilska and J. Wojcieszak, "Designer cathinones—an emerging class of novel recreational drugs," *Forensic science international*, vol. 231, no. 1-3, pp. 42–53, 2013.
- [7] S. Rolles and D. Kushlick, "Prohibition is a key driver of the new psychoactive substances (NPS) phenomenon," *Addiction*, vol. 109, no. 10, pp. 1589–1590, 2014.
- [8] F. P. Busardò, C. Kyriakou, S. Napoletano, E. Marinelli, and S. Zaami, "Mephedrone related fatalities: a review," *European Review for Medical and Pharmacological Sciences*, vol. 19, no. 19, pp. 3777–3790, 2015.
- [9] E. Zuccato and S. Castiglioni, "Illicit drugs in the environment," *Philosophical transactions. Series A, Mathematical, physical, and engineering sciences*, vol. 367, pp. 3965–3978, 2009.
- [10] L. A. Maranho, G. V. Aguirre-Martínez, and M. L. Martín-Díaz, "Chapter 2. Adverse effects of pharmaceutical products in the marine environment: the use of non-target species to evaluate water and sediment matrices," in *Proceedings of the Ecotoxicology and Genotoxicology*, 2017.
- [11] H. M. S. B. I. Escher and E. L. Schymanski, "Tracking complex mixtures of chemicals in our changing environment," *Chemistry for Tomorrow'Searth*, vol. 367, 2020.
- [12] B. De Felice and M. Parolini, "Effects of single and combined exposure to cocaine and benzoylecgonine on the oxidative status of *Mytilus galloprovincialis*," *Environmental Toxicology and Pharmacology*, vol. 80, p. 103475, 2020.
- [13] B. I. Escher, H. M. Stapleton, and E. L. Schymanski, "Tracking complex mixtures of chemicals in our changing environment," *Science*, vol. 367, no. 6476, pp. 388–392, 2020.
- [14] UN Transforming our World, "The 2030 agenda for sustainable development," *UN Transforming our World*, vol. 70/1, 2014.
- [15] P. Griffiths, V. Mravcik, D. Lopez, and D. Klempova, "Quite a lot of smoke but very limited fire—the use of methamphetamine in Europe," *Drug and Alcohol Review*, vol. 27, no. 3, pp. 236–242, 2008.
- [16] P. Griffiths, M. Meacham, and R. McKetin, "Illicit drug trends globally," *Journal of Public Health*, 2008.
- [17] H. A. Yassa and S. T. Badea, "Patterns of drug abuse in Upper Egypt: cause or result of violence?" *Egyptian Journal of Food Science*, vol. 9, no. 1, p. 14, 2019.
- [18] B. Moslah, E. Hapeshi, A. Hedhili, A. Jrad, and D. Fatta-Kassinos, *Pharmaceuticals and Illicit Drugs in Wastewater Samples in North-Eastern Tunisia*, Environmental Science and Pollution Research, 2017.
- [19] N. Daglioglu, Evsen Yavuz Guzel, E. Y. Guzel, and S. Kilercioglu, "Assessment of illicit drugs in wastewater and estimation of drugs of abuse in adana province, Turkey," *Forensic Science International*, vol. 294, pp. 132–139, 2019.
- [20] R. Pal, M. Megharaj, K. P. Kirkbride, and R. Naidu, "Illicit drugs and the environment - a review," *The Science of the Total Environment*, vol. 463-464, no. 2013, pp. 1079–1092, 2013.

- [21] K. V. Thomas, "Comparing Illicit Drug Use in 19 European Cities through Sewage Analysis," *Science of the Total Environment*, vol. 838, 2012.
- [22] E. Zuccato and S. Castiglioni, "Illicit drugs in the environment," *Philosophical Transactions of the Royal Society Mathematical, Physical Engineering Sciences*, vol. 367, no. 1904, pp. 3965–3978, 2009.
- [23] L. Degenhardt and W. Hall, "Extent of illicit drug use and dependence, and their contribution to the global burden of disease," *The Lancet*, vol. 379, no. 9810, pp. 55–70, 2012.
- [24] R. G. Deiss, T. C. Rodwell, and R. S. Garfein, "Tuberculosis and illicit drug use: review and update," *Clinical Infectious Diseases*, vol. 48, no. 1, pp. 72–82, 2009.
- [25] W. Hall, L. Degenhardt, and N. Sindicich, *Illicit Drug Use and the Burden of Disease*, 2008.
- [26] F. P. Busardò, C. Kyriakou, S. Napoletano, E. Marinelli, and S. Zaami, "Mephedrone related fatalities: a review," *European Review for Medical and Pharmacological Sciences*, vol. 19, no. 19, pp. 3777–3790, 2015.
- [27] S. Zaami, R. Giorgetti, S. Pichini, F. Pantano, E. Marinelli, and F. P. Busardò, "Synthetic cathinones related fatalities: an update," *European Review for Medical and Pharmacological Sciences*, vol. 22, no. 1, pp. 268–274, 2018.
- [28] F. Gilani, "Novel psychoactive substances: the rising wave of 'legal highs,'" *British Journal of General Practice*, vol. 66, 2016.
- [29] C. L. German, A. E. Fleckenstein, and G. R. Hanson, "Bath salts and synthetic cathinones: an emerging designer drug phenomenon," *Life Sciences*, vol. 97, no. 1, pp. 2–8, 2014.
- [30] S. M. Somani, "Substance abuse among youth: a harsh reality," *Emergency Medicine*, vol. 6, no. 330, p. 2, 2016.
- [31] A. Kabore, E. Afriyie-Gyawu, J. Awuah et al., "Social ecological factors affecting substance abuse in Ghana (West Africa) using photovoice," *The Pan African Medical Journal*, vol. 34, p. 214, 2019.
- [32] A. Kabore, E. Afriyie-Gyawu, A. D. Walker, M. Hester, and A. R. Hansen, *Barriers and Facilitators of Substance Abuse Treatment in Ghana (West Africa): A Social-Ecological Perspective Using Photovoice*, 2016.
- [33] D. M. Wood and P. I. Dargan, "Understanding how data triangulation identifies acute toxicity of novel psychoactive drugs," *Journal of Medical Toxicology*, vol. 8, no. 3, pp. 300–303, 2012.
- [34] A. Zwartsen, L. Hondebrink, and R. H. Westerink, "Neurotoxicity screening of new psychoactive substances (NPS): effects on neuronal activity in rat cortical cultures using microelectrode arrays (MEA)," *NeuroToxicology*, vol. 66, pp. 87–97, 2018.
- [35] P. Adamowicz and A. Malczyk, "Stability of synthetic cathinones in blood and urine," *Forensic Science International*, vol. 295, pp. 36–45, 2019.
- [36] P. D. Xiqing Li and W. Zhang, "Application of wastewater-based epidemiology in China—From Wastewater Monitoring to Drug Control Efforts," *ACS Publication Most Trusted Most Cited Most Read*, 2019.
- [37] S. Koster, M. Rennen, W. Leeman et al., "A novel safety assessment strategy for non-intentionally added substances (NIAS) in carton food contact materials," *Food Additives & Contaminants: Part A*, vol. 31, no. 3, pp. 422–443, 2014.
- [38] X. Li, P. Du, and W. Zhang, "Application of wastewater-based epidemiology in China—from wastewater monitoring to drug control efforts," in *Proceedings of the Wastewater-Based Epidemiology: Estimation of Community Consumption of Drugs and Diets*, pp. 119–135, 2019.
- [39] E. Archer, E. Castrignanò, B. Kasprzyk-Hordern, and G. M. Wolfaardt, "Wastewater-based epidemiology and enantiomeric profiling for drugs of abuse in South African wastewaters," *The Science of the Total Environment*, vol. 625, pp. 792–800, 2018.
- [40] P. Du, Q. Zheng, K. V. Thomas, X. Li, and P. K. Thai, "A revised excretion factor for estimating ketamine consumption by wastewater-based epidemiology - utilising wastewater and seizure data," *Environment International*, vol. 138, Article ID 105645, 2020.
- [41] K. A. A. Kankaanpää, M. Heinonen, and K. Kuoppasalmi, "Current trends in Finnish drug abuse: wastewater based epidemiology combined with other national indicators," *The Science of the Total Environment*, vol. 2016, 2016.
- [42] R. David, "Baker and Barbara Kasprzyk-Hordern, "Multi-residue analysis of drugs of abuse in wastewater and surface water by solid-phase extraction and liquid chromatography–positive electrospray ionisation tandem mass spectrometry," *Journal of Chromatography A*, vol. 1218, no. 12, pp. 1620–1631, 2011.
- [43] A. L. N. Van Nuijs, J.-F. Mougel, I. Tarcomnicu et al., "Sewage epidemiology - a real-time approach to estimate the consumption of illicit drugs in Brussels, Belgium," *Environment International*, vol. 37, no. 3, pp. 612–621, 2011.
- [44] C. Abbo, E. S. Okello, W. Muhwezi, G. Akello, and E. Ovuga, "Alcohol, substance use and psychosocial competence of adolescents in selected secondary schools in Uganda: a cross sectional survey," *J International neuropsychiatric disease journal*, vol. 7, no. 2, Article ID 25387, 2016.
- [45] S. Tesfamariam, I. S. Anand, G. Kaleab et al., "Self-medication with over the counter drugs, prevalence of risky practice and its associated factors in pharmacy outlets of Asmara, Eritrea," *BMC Public Health*, vol. 19, no. 1, pp. 159–9, 2019.
- [46] M. Odenwald, H. Hinkel, E. Schauer et al., "The consumption of khat and other drugs in Somali combatants: a cross-sectional study," *PLoS Medicine*, vol. 4, no. 12, p. e341, 2007.
- [47] UNODC, *Global-ATS-Assessment-Exec-summary*, UNODC, Vienna, Austria, 2019.
- [48] UNODC, *UNODC World Drug Report 2020: Global Drug Use Rising; while COVID-19 Has Far Reaching Impact on Global Drug Markets*, UNODC, Vienna, Austria, 2020.
- [49] S. D. Brandt, S. Freeman, H. R. Sumnall, F. Measham, and J. Cole, "Analysis of NRG 'legal highs' in the UK: identification and formation of novel cathinones," *Drug Testing and Analysis*, vol. 3, no. 9, pp. 569–575, 2011.
- [50] A. A. Olashore, P. R. Opondo, J. A. Ogunjumo, and J. O. J. S. a. r. Ayugi, "Cannabis use disorder among first-year undergraduate students in Gaborone, Botswana," *Substance Abuse: Research and Treatment*, vol. 14, Article ID 1178221820904136, 2020.
- [51] M. Cristina, T. Marta, P. Ricardo, P. Clara, and F. J. A. Magi, "Spice drugs: cannabinoids as a new designer drugs," *Adicciones*, vol. 21, no. 3, 2009.
- [52] C. D. Smith and S. Robert, "'Designer drugs': update on the management of novel psychoactive substance misuse in the acute care setting," *Clinical Medicine*, vol. 14, no. 4, pp. 409–415, 2014.
- [53] R. Pal, M. Megharaj, K. P. Kirkbride, and R. Naidu, "Illicit drugs and the environment - a review," *The Science of the Total Environment*, vol. 463–464, pp. 1079–1092, 2013.
- [54] B. Moslah, E. Hapeshi, A. Jrad, D. Fatta-Kassinou, and A. Hedhili, "Pharmaceuticals and illicit drugs in wastewater samples in north-eastern Tunisia," *Environmental Science*

- and Pollution Research*, vol. 25, no. 19, pp. 18226–18241, 2018.
- [55] A. Olawole-Isaac, O. Ogundipe, E. O. Amoo, and D. Adeyoye, “Substance use among adolescents in sub-Saharan Africa: a systematic review and meta-analysis,” *South African Journal of Child Health*, vol. 12, pp. s79–s84, 2018.
- [56] R. Hussien, S. Ahmed, H. Awad, M. El-Setouhy, M. El-Shinawi, and J. M. J. I. J. o. E. A. C. Hirshon, “Identification of ‘Voodoo’: an emerging substance of abuse in Egypt,” *International Journal of Environmental Analytical Chemistry*, pp. 1–13, 2020.
- [57] W. Acuda, C. J. Othieno, A. Obondo, and I. B. Crome, “The epidemiology of addiction in Sub-Saharan Africa: a synthesis of reports,” *Reviews, and Original Articles*, vol. 20, no. 2, pp. 87–99, 2011.
- [58] O. A. Adedeji, J. Abdulmalik, and O. Gureje, *Contemporary Issues in Mental Health Care in Sub-saharan Africa*, Book-Builders Editions Africa, Nigeria West Africa, 2017.
- [59] A. O. C. Iroegbu, R. E. Sadiku, S. S. Ray, and Y. Hamam, “Plastics in municipal drinking water and wastewater treatment plant effluents: challenges and opportunities for South Africa—a review,” *Environmental Science and Pollution Research*, vol. 27, no. 12, pp. 12953–12966, 2020.
- [60] S. M. Richards, “Effects of pharmaceutical mixtures in aquatic microcosms,” *Environmental Toxicology & Chemistry*, vol. 23, no. 4, pp. 1035–1042, 2009.
- [61] B. I. Escher, H. M. Stapleton, and E. L. Schymanski, “Tracking complex mixtures of chemicals in our changing environment,” *Chemistry for Tomorrow’s earth*, 2020.
- [62] R. Aghababaei, I. Javadi, A. Nili-Ahmadabadi, S. Parsafar, and D. Ahmadi Moghaddam, “Occurrence of bacterial and toxic metals contamination in illegal opioid-like drugs in Iran: a significant health challenge in drug abusers,” *Daru Journal of Pharmaceutical Sciences*, vol. 26, no. 1, pp. 77–83, 2018.
- [63] B. Bonnefille, E. Gomez, F. Courant, A. Escande, and H. Fenet, “Diclofenac in the marine environment: a review of its occurrence and effects,” *Marine Pollution Bulletin*, vol. 131, pp. 496–506, 2018.
- [64] J.-F. Akoachere, N. Guelle, H. M. Dilonga, and T. K. Nkuo-Akenji, “Public health implications of contamination of Franc CFA (XAF) circulating in Buea (Cameroon) with drug resistant pathogens,” *BMC Research Notes*, vol. 7, no. 1, p. 16, 2014.
- [65] K. Abdulwahab, *Effects of Pharmaceutical Effluent on the Physico-Chemical Parameters and Distribution of Plankton in Okun Stream in Ilorin, Kwara State, Nigeria*, Kwara State University, Nigeria West Africa), 2020.
- [66] P. Sathishkumar, R. A. A. Meena, T. Palanisami, V. Ashokkumar, T. Palvannan, and F. L. Gu, “Occurrence, interactive effects and ecological risk of diclofenac in environmental compartments and biota - a review,” *The Science of the Total Environment*, vol. 698, Article ID 134057, 2020.
- [67] P. K. Olatayo, M. Ogunbanwo, A. B. Boxall et al., “High concentrations of pharmaceuticals in a Nigerian river catchment,” *Environmental Chemistry*, vol. 41, 2020.
- [68] E. N. Pius Kairigo and L.-Riina Sundberg, “Anthony gachanja and tuula tuhkanen “contamination of surface water and river sediments by antibiotic and antiretroviral drug cocktails in low and middle-income countries: occurrence, risk and mitigation strategies,” *Water*, vol. 5, no. 12, p. 1376, 2020.
- [69] J. A. Russell and A. Mehrabein, “Environmental effects of drug abuse,” *Environmental Psychology & Nonverbal Behavior*, vol. 2, no. 2, 1977.
- [70] C. D. S. P. Mayana Karoline Fontes and L. Alves Maranhão, “Review on the Occurrence and Biological Effects of Illicit Drugs in Aquatic Ecosystems,” *Environmental Science and Pollution Research*, vol. 29, 2022.
- [71] K. G. Maria Eugenia and S. Santos, “Christoph steinbach ,heike schmidt-posthaus ,eva salkova , jitka kolarova , andrea vojs stanova ,roman grabic tomas randak, “environmental concentration of methamphetamine induces pathological changes in brown trout (*Salmo trutta fario*),” *Chemosphere*, vol. 254, no. 2020, Article ID 126882, 2020.
- [72] M. S. Ângela Almeida, A. M. V. M. Soares, and R. Freitas, “Anti-inflammatory Drugs in the Marine Environment: Bioconcentration,” *Metabolism and Sub-lethal Effects in Marine Bivalves*, vol. 263, 2020.
- [73] H. Mohan, S. S. Rajput, E. B. Jadhav, M. Singh Sankh, S. J. Sonone, and R. Kumar, “Ecotoxicity, occurrence, and removal of pharmaceuticals and illicit drugs from aquatic systems,” *Biointerface Research in Applied Chemistry*, vol. 1, no. 5, pp. 12530–12546, 2021.
- [74] K. Grabicová, R. Grabic, G. Kolářová, J. Turek, and B. W. Randák, “Psychoactive pharmaceuticals in aquatic systems: a comparative assessment of environmental monitoring approaches for water and fish,” *Environmental Pollution*, vol. 261, Article ID 114150, 2020.
- [75] P. Kairigo, L.-R. S. Elijah Ngumba, G. Anthony, and T. Tuhkanen, “Contamination of surface water and river sediments by antibiotic and antiretroviral drug cocktails in low and middle-income countries: occurrence, risk and mitigation strategies,” *Water*, vol. 5, no. 12, p. 1376, 2020.
- [76] S. Castiglioni, L. Vandam, and P. Griffiths, “Assessing illicit drugs in wastewater,” *J Assessing illicit drugs in wastewater Advances in wastewater-Based drug epidemiology, EMCDDA insights*, vol. 22, pp. 1–82, 2016.
- [77] S. Castiglioni, “Estimating community drug use through wastewater-based epidemiology,” *J Assessing illicit drugs in wastewater: Advances in Wastewater-Based Drug Epidemiology, EMCDDA Insights*, vol. 22, pp. 17–33, 2016.
- [78] E. Archer, M. Volschenk, L. Brocker, and G. M. Wolfaardt, “A two-year study of emerging micro-pollutants and drugs of abuse in two Western Cape wastewater treatment works (South Africa),” *Chemosphere*, vol. 285, Article ID 131460, 2021.
- [79] O. Ganzenko, C. Trelu, N. Oturan et al., “Electro-Fenton treatment of a complex pharmaceutical mixture: mineralization efficiency and biodegradability enhancement,” *Chemosphere*, vol. 253, Article ID 126659, 2020.
- [80] A. Grossberger, Y. Hadar, T. Borch, and B. Chefetz, “Biodegradability of pharmaceutical compounds in agricultural soils irrigated with treated wastewater,” *Environmental Pollution*, vol. 185, pp. 168–177, 2014.
- [81] J. Nyirenda, A. Mwanza, and C. Lengwe, “Assessing the biodegradability of common pharmaceutical products (PPs) on the Zambian market,” *Heliyon*, vol. 6, no. 10, Article ID e05286, 2020.
- [82] A. Carucci, G. Cappai, and M. Piredda, “Biodegradability and toxicity of pharmaceuticals in biological wastewater treatment plants,” *Journal of Environmental Science and Health, Part A*, vol. 41, no. 9, pp. 1831–1842, 2006.
- [83] C. Nannou, A. Ofrydopoulou, E. Evgenidou, D. Heath, E. Heath, and D. Lambropoulou, *Antiviral Drugs in Aquatic Environment and Wastewater Treatment Plants: A Review on*

- Occurrence, Fate, Removal and Ecotoxicity*, Science of the Total Environment, 2019.
- [84] K. Lancastera, A. Ritter, k. valentine, and d. Tim Rhodessa, "A more accurate understanding of drug use": a critical analysis of wastewater analysis technology for drug policy," *International Journal of Drug Policy*, vol. 63, no. 2019, pp. 47–55, 2019.
- [85] E. Ekuadzi, G. Darko, H. Senanu Ahor et al., "Occurrence of antibiotics and antibiotic-resistant bacteria in landfill sites in Kumasi, Ghana," *Journal of Chemistry*, vol. 2019, no. 10, Article ID 6934507, 2019.
- [86] A. Celma, J. V. Sancho, N. Salgueiro-González et al., "Simultaneous determination of new psychoactive substances and illicit drugs in sewage: potential of micro-liquid chromatography tandem mass spectrometry in wastewater-based epidemiology," *Journal of Chromatography A*, vol. 1602, pp. 300–309, 2019.
- [87] M. L. Brusseau and J. F. Artiola, "Chemical contaminants," in *Proceedings of the Environmental and Pollution Science*, pp. 175–190, 2019.
- [88] D. Hanigan, E. M. Thurman, I. Ferrer et al., "Methadone contributes to N-nitrosodimethylamine formation in surface waters and wastewaters during chloramination," *Environmental Science and Technology Letters*, vol. 2, no. 6, pp. 151–157, 2015.
- [89] C. G. Daughton and T. A. Ternes, "Pharmaceuticals and personal care products in the environment: agents of subtle change?" *J Environmental Health Perspectives*, vol. 107, no. suppl 6, pp. 907–938, 1999.
- [90] E. J. Rosi-Marshall, D. Snow, S. L. Bartelt-Hunt, A. Paspalof, and J. L. Tank, "A review of ecological effects and environmental fate of illicit drugs in aquatic ecosystems," *Journal of Hazardous Materials*, vol. 282, pp. 18–25, 2015.
- [91] M. K. Fontes, L. A. Maranhão, and C. D. S. Pereira, "Review on the occurrence and biological effects of illicit drugs in aquatic ecosystems," *J Environmental Science Pollution Research*, pp. 1–37, 2020.
- [92] B. Kasprzyk-Hordern, R. M. Dinsdale, and A. J. Guwy, "The occurrence of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs in surface water in South Wales, UK," *Water Research*, vol. 42, no. 13, pp. 3498–3518, 2008.
- [93] P. Horký, R. Grabic, K. Grabicová et al., "Methamphetamine pollution elicits addiction in wild fish," *Journal of Experimental Biology*, vol. 224, no. 13, Article ID jeb242145, 2021.
- [94] W. Sanchez, W. Sremski, B. Piccini et al., "Adverse effects in wild fish living downstream from pharmaceutical manufacture discharges," *Environment International*, vol. 37, no. 8, pp. 1342–1348, 2011.
- [95] K. A. Kidd, "Collapse of a fish population after exposure to a synthetic estrogen," *Proceedings of the National Academy of Sciences*, vol. 104, no. 21, pp. 8897–8901, 2007.
- [96] S. Jarque, L. Quirós, J. O. Grimalt et al., "Background fish feminization effects in European remote sites," *Scientific Reports*, vol. 5, no. 1, pp. 11292–11296, 2015.
- [97] M. Barakat, R. a. Al-Qudah, A. Akour, N. Al-Qudah, and Y. H. Dallal Bashi, "Unforeseen uses of oral contraceptive pills: exploratory study in Jordanian community pharmacies," *PLoS One*, vol. 15, no. 12, Article ID e0244373, 2020.
- [98] J. Li, J. Gao, P. K. Thai, J. F. Mueller, Z. Yuan, and G. Jiang, "Transformation of illicit drugs and pharmaceuticals in sewer sediments," *Environmental Science & Technology*, vol. 54, no. 20, pp. 13056–13065, 2020.
- [99] T. A.-A. B. S. Choudri, Yassine Charabi, Noura Al-Nasiri "Wastewater Treatment, Reuse, and Disposal-Associated Effects on Environment and Health," Water Environment Federation, Virginia, United States, 2020.
- [100] C. N. Janeiro, A. M. Arsénio, R. Brito, and J. van Lier, *Use of (Partially) Treated Municipal Wastewater in Irrigated Agriculture*, p. 2020, Potentials and Constraints for Sub-Saharan Africa, 2020.
- [101] K. O. Oluwatobi Idowu Alawode and Ayodeji Fisayo Afolayan, "Prospects of wastewater reclamation and reuse for water scarcity mitigation and environmental pollution control in sub-Saharan Africa," *Environmental Management and Sustainable Development*, vol. 8, no. 4, 2019.
- [102] N. J. Gulamussen, A. M. Arsénio, N. P. Matsinhe, and L. C. Rietveld, "Water reclamation for industrial use in sub-Saharan Africa - a critical review," *Drinking Water Engineering and Science*, vol. 12, no. 2, pp. 45–58, 2019.
- [103] T. A. Ternes, "Occurrence of drugs in German sewage treatment plants and rivers," Dedicated to Professor Dr. Klaus Haberer on the occasion of his 70th birthday.1, *Water Research*, vol. 32, no. 11, pp. 3245–3260, 1998.
- [104] H. Zarei, Y. Salimi, E. Repo, N. D. Safaei, E. Güzel, and A. Asadi, *A Global Systematic Review and Meta-Analysis on Illicit Drug Consumption Rate through Wastewater-Based Epidemiology*, Environmental Science and Pollution Research, 2020.
- [105] T. Heberer, "Occurrence, fate, and removal of pharmaceutical residues in the aquatic environment: a review of recent research data," *Toxicology Letters*, vol. 131, no. 1-2, pp. 5–17, 2002.
- [106] A. Nikolaou, S. Meric, D. J. A. Fatta, and B. chemistry, "Occurrence patterns of pharmaceuticals in water and wastewater environments," *Analytical Bioanalytical Chemistry*, vol. 387, no. 4, pp. 1225–1234, 2007.
- [107] V. Yargeau, A. Metcalfe, and D. Chris, "Analysis of drugs of abuse in wastewater from two Canadian cities," *The Science of the Total Environment*, vol. 487, pp. 722–730, 2014.
- [108] R. Pal, M. Megharaj, and K. P. Naidu, "Illicit drugs and the environment - a review," *The Science of the Total Environment*, vol. 463-464, pp. 1079–1092, 2013.
- [109] E. E. Lubertus Bijlsma, F. . Hernández, and P. de Voogt, "Investigation of drugs of abuse and relevant metabolites in Dutch sewage water by liquid chromatography coupled to high resolution mass spectrometry," *Chemosphere*, vol. 89, 2012.
- [110] L. B. Kevin, V. Thomas, S. Castiglioni et al., "Comparing illicit drug use in 19 European cities through sewage analysis," *Science of the Total Environment*, vol. 432, 2012.
- [111] K. Lancaster, A. Ritter, k. valentine, and T. Rhodessa, "A more accurate understanding of drug use": a critical analysis of wastewater analysis technology for drug policy," *International Journal of Drug Policy*, vol. 63, no. 2019, pp. 47–55, 2019.
- [112] L. Feng, W. Zhang, and X. Li, "Monitoring of regional drug abuse through wastewater-based epidemiology-A critical review," *Science China Earth Sciences*, vol. 61, no. 3, pp. 239–255, 2018.
- [113] C. Postigo, M. L. de Alda, and D. Barceló, "Evaluation of drugs of abuse use and trends in a prison through wastewater analysis," *Environment International*, vol. 37, no. 1, pp. 49–55, 2011.
- [114] A. L. N. Castiglioni, I. Tarcomnicu, C. Postigo et al., "Illicit drug consumption estimations derived from wastewater

- analysis: a critical review," *The Science of the Total Environment*, vol. 409, no. 19, pp. 3564–3577, 2011.
- [115] K. Y. Kim, F. Y. Lai, H.-Y. Kim, P. K. Thai, J. F. Mueller, and J.-E. Oh, "The first application of wastewater-based drug epidemiology in five South Korean cities," *The Science of the Total Environment*, vol. 524-525, pp. 440–446, 2015.
- [116] J. Rice, A. M. Kannan, E. Castrignanò, K. Jagadeesan, and B. Kasprzyk-Hordern, "Wastewater-based epidemiology combined with local prescription analysis as a tool for temporal monitoring of drugs trends - a UK perspective," *The Science of the Total Environment*, vol. 735, p. 139433, 2020.
- [117] S.-Y. Liu, W.-J. Yu, Y.-R. Wang, X.-T. Shao, and D.-G. Wang, "Tracing consumption patterns of stimulants, opioids, and ketamine in China by wastewater-based epidemiology," *Environmental Science and Pollution Research*, vol. 28, no. 13, pp. 16754–16766, 2021.
- [118] F. Nyabadza, L. J. C. Coetzee, and m. m. i. medicine, "A systems dynamic model for drug abuse and drug-related crime in the Western Cape province of South Africa," *Computational and mathematical methods in medicine*, vol. 2017, 2017.
- [119] C. Hsiao, D Fry, C. L Ward et al., "Violence against children in South Africa: the cost of inaction to society and the economy," *BMJ global health*, vol. 3, no. 1, Article ID e000573, 2018.
- [120] S. Dada, N. Harker Burnhams, J. Erasmus, W. Lucas Charles Parry, A. Bhana Sandra Pretorius, and R. J. S. A. Weimann Helen Keen, "Monitoring alcohol, tobacco and other drug abuse treatment admissions in South Africa," *SACENDU*, pp. 1–72, 2018.
- [121] S. Dada, N. H. Burnhams, R. Laubscher, C. Parry, and B. J. S. A. J. o. S. Myers, "Alcohol and other drug use among women seeking substance abuse treatment in the Western Cape," *South Africa Journal of Science*, vol. 114, no. 9-10, pp. 1–7, 2018.
- [122] J. Sommer, M. Hinsberger, T. Elbert et al., "The interplay between trauma, substance abuse and appetitive aggression and its relation to criminal activity among high-risk males in South Africa," *Addictive Behaviors*, vol. 64, pp. 29–34, 2017.
- [123] E. Castrignanò, Z Yang, R Bade et al., "Enantiomeric profiling of chiral illicit drugs in a pan-European study," *Water Research*, vol. 130, pp. 151–160, 2018.
- [124] E. Archer, E. Castrignanò, B. Kasprzyk-Hordern, and G. M. Wolfaardt, "Wastewater-based epidemiology and enantiomeric profiling for drugs of abuse in South African wastewaters," *The Science of the Total Environment*, vol. 625, pp. 792–800, 2018.
- [125] B. Moslah, E. Hapeshi, A. Jrad, D. Fatta-Kassinos, A. Hedhili, and P. Research, "Pharmaceuticals and illicit drugs in wastewater samples in north-eastern Tunisia," *Environmental Science and Pollution Research*, vol. 25, no. 19, pp. 18226–18241, 2018.
- [126] S. M. Richards, "Effects of pharmaceutical mixtures in aquatic microcosms," *Environmental Toxicology & Chemistry*, vol. 23, no. 4, pp. 1035–1042, 2009.
- [127] M.-T. G. Molla-Mahmoudi, M. Shokri-Khubestani, M. Hassani, and A. H. Bahramzadeh, "Investigating the effects of pharmaceutical wastes on the environment and human health (systematic review)," *Journal of Safety Promotion and Injury Prevention*, vol. 6, 2018.
- [128] I. Kosamu, C. Kaonga, and W. Utembe, "A critical review of the status of pesticide exposure management in Malawi," *International Journal of Environmental Research and Public Health*, vol. 17, no. 18, p. 6727, 2020.
- [129] Z. Xu, J. Xu, H. Yin, W. Jin, H. Li, and Z. He, "Urban river pollution control in developing countries," *Nature Sustainability*, vol. 2, no. 3, pp. 158–160, 2019.
- [130] E. S. Bernhardt, *Synthesizing US River Restoration Efforts*, American Association for the Advancement of Science, Washington USA, 2005.
- [131] B. M. Gannon, K. I. Galindo, M. P. Mesmin, K. C. Rice, and G. T. Collins, "Reinforcing effects of binary mixtures of common bath salt constituents: studies with 3,4-methylenedioxypropylvalerone (MDPV), 3,4-methylenedioxy-methylcathinone (methylone), and caffeine in rats," *Neuropsychopharmacology*, vol. 43, no. 4, pp. 761–769, 2018.
- [132] K. N. Ellefsen, E. A. Taylor, P. Simmons, V. Willoughby, and B. J. Hall, "Multiple drug-toxicity involving novel psychoactive substances, 3-fluorophenmetrazine and U-47700," *Journal of Analytical Toxicology*, vol. 41, no. 9, pp. 765–770, 2017.