

2021-04-14

# Solvent effects on host-guest residence time and kinetics: further insights from metadynamics simulation of Toussaintine-A unbinding from chitosan nanoparticle

Shadrack, Daniel

Springer Nature

---

<https://doi.org/10.1007/s00894-021-04735-y>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

# Solvent effects on host-guest residence time and kinetics: further insights from metadynamics simulation of Toussaintine-A unbinding from chitosan nanoparticle

Daniel M Shadrack, Lucy W Kiruri, Hulda Swai, Stephen S Nyandoro

To download the complete text, click that link.

DOI: <https://doi.org/10.1007/s00894-021-04735-y>

## Abstract

Solvents play an important role in host-guest intermolecular interactions. The kinetics and residence time of Toussaintine-A (TouA) unbinding from chitosan was investigated by means of well-tempered metadynamics and thermodynamic integration using two solvents, polar aprotic (DMSO), and polar protic (water). The kinetic rates were found to be strongly dependent on the solvent polarity; hence, the unbinding rate proceeded much faster in DMSO compared to water. DMSO tends to participate less in a chemical reaction by weakening the intermolecular interaction between chitosan and TouA due to lack of acidic hydrogen resulting in a reduction of the transition state. On the other hand, water, which ought to donate hydrogen atoms, sustains a strong interaction and hence large barrier heights. Consequently, this reduces the unbinding rate and increases the residence time. Binding free energy from thermodynamic integration suggests a thermodynamic stable chitosan-TouA complex in water than in DMSO.

## Keywords:

Residence time; Solvents; Toussaintine-A; Chitosan; Metadynamics; Kinetics