

2023-07-13

# Radioactivity and dose assessment of naturally occurring radionuclides in terrestrial environments and foodstuffs: a review of Bahi district, Tanzania

Sumary, Dominic

International Journal of Environmental Health Research

---

<https://doi.org/10.1080/09603123.2023.2234299>

*Provided with love from The Nelson Mandela African Institution of Science and Technology*

# Radioactivity and dose assessment of naturally occurring radionuclides in terrestrial environments and foodstuffs: a review of Bahi district, Tanzania

Dominic Parmena Sumary, Jofrey Raymond, Musa Chacha, Frimi Paul Banzi

To download the complete text,click that link

DOI: <https://doi.org/10.1080/09603123.2023.2234299>

## Abstract

In this review, the online searchable research articles were scrutinized for the data presented in line with radioactivity and dose estimates from both terrestrial environments and foodstuffs from Bahi district and other parts of Tanzania. The data on natural gamma ray dose rates from Bahi localities were observed with variations among researchers. The observed ranges of radioactivity concentrations ( $\text{Bq kg}^{-1}$ ) in soil were  $^{226}\text{Ra}$  (28.5–57.4),  $^{232}\text{Th}$  (38.1–521.3), and  $^{40}\text{K}$  (562.9–665.0). Deep closed water wells with installed pumps from Ilindi and Bahi Mission reported radioactivity concentration of  $^{238}\text{U}$   $3.08 \text{ Bq L}^{-1}$  and Ilindi swamps reported radioactivity concentrations of  $^{226}\text{Ra}$   $15.35 \text{ Bq L}^{-1}$ , whereas radioactivity concentrations of  $^{238}\text{U}$  in cereals were within the annual tolerable limits of  $0.001\text{--}0.02 \text{ Bq kg}^{-1}$ . The quantity and accessibility of published studies, as well as the diversity of the data, point to the necessity for additional studies to be carried out in order to obtain comprehensive baseline data.

## Keywords

[Radioactivity](#),[radionuclide](#),[Uranium](#),[norms](#),[Bahi district](#)