

2007-04-18

Modeling the proteome of a Marek's disease transformed cell line: a natural animal model for CD30 overexpressing lymphomas

Buza, Joram

Proteomics

<https://doi.org/10.1002/pmic.200600946>

Provided with love from The Nelson Mandela African Institution of Science and Technology

Modeling the proteome of a Marek's disease transformed cell line: a natural animal model for CD30 overexpressing lymphomas

Joram J. Buza, Shane C. Burgess

DOI//doi.org/10.1002/pmic.200600946

Abstract

Marek's disease (MD) in the chicken, caused by the highly infectious MD α -herpesvirus (MDV), is both commercially important and a unique, naturally occurring model for human T-cell lymphomas overexpressing the Hodgkin's disease antigen, CD30. Here, we used proteomics as a basis for modeling the molecular functions and biological processes involved in MDV-induced lymphomagenesis. Proteins were extracted from an MDV-transformed cell line and were then identified using 2-D LC-ESI-MS/MS. From the resulting 3870 cellular and 21 MDV proteins we confirm the existence of 3150 "predicted" and 12 "hypothetical" chicken proteins. The UA-01 proteome is proliferative, differentiated, angiogenic, pro-metastatic and pro-immune-escape but anti-programmed cell death, -anergy, -quiescence and -senescence and is consistent with a cancer phenotype. In particular, the pro-metastatic integrin signaling pathway and the ERK/MAPK signaling pathways were the two predominant signaling pathways represented. The cytokines, cytokine receptors, and their related proteins suggest that UA-01 has a regulatory T-cell phenotype.